



CMHD Pathology Report



CMHD Pathology Core

Toronto Centre for
Phenogenomics
25 Orde St. 3rd fl.
Toronto, Ont. M5T 3H7
Tel.(416) 586-8375
Fax (416) 586-5993

contact: Dr. Susan
Newbigging
email:
newbigging@lunenfeld.ca

Principle Investigator: Dr. Jacqui White

Institute: Wellcome Trust Sanger Institute
Address: Attn: Linda Read Wellcome Trust
Genome Campus Hinxton Cambridge CB10
1SA, UK

ReportID: Report Date: May 21, 2013
Pathologist: Dr. H. Adissu

Mouse Genetics Project

Wellcome Trust Sanger
Institute
Wellcome Trust Genome
Campus
Hinxton, Cambridge
CB10 1SA
UK

email:
MGPenquiries@sanger.ac.uk
[Mouse Portal](#)
[Europhenome](#)

CMHD LabID: N13-233

Relevant History:

Phenotype

hindlimb paralysis
tremors

AnimalID: M00386485 (Male)

Histopathology Findings:

liver (MA:0000358)

Histopath Description:

severe lipidosis

Morphological Diagnosis:

Distribution: diffuse; **Severity:** severe; **MPATH Diagnosis:** steatosis MPATH:622

Definitive Diagnosis:

Hepatic lipidosis

retina (MA:0000276)

Histopath Description:


Involving one eye, there are clusters of external nuclear structures within the internal plexiform layer.

Morphological Diagnosis:

Distribution: Focal; **Severity:** mild;

Definitive Diagnosis:

Retinal dysplasia



Retina, dysplasia,
10x, HE

brown fat (MA:0000057)

Histopath Description:


There is focally extensive hyperplasia of brown fat infiltrated with low numbers of inflammatory cells (lymphocytes, macrophages and rare neutrophils)

Morphological Diagnosis:

Duration: Chronic-active; **Distribution:** Focal; **Severity:** mild; **MPATH Diagnosis:** steatitis MPATH:636

Definitive Diagnosis:

Steatitis with hyperplasia



Peripheral nerve,
normal, 40x, HE

AnimalID: M00386484 (Male)

Histopathology Findings:

liver (MA:0000358)

Histopath Description:

severe lipidosis

Morphological Diagnosis:

Distribution: diffuse; **Severity:** severe; **MPATH Diagnosis:** steatosis MPATH:622

Definitive Diagnosis:

Hepatic lipidosis

retina (MA:0000276)

Histopath Description:


Involving one eye, there are clusters of external nuclear structures within the internal plexiform layer.

Morphological Diagnosis:


Distribution: Focal; **Severity:** mild;

Definitive Diagnosis:

Retinal dysplasia



Retina, dysplasia,
10x, HE



Peripheral nerve,
normal, 40x, HE

AnimalID: M00386491 (Female)

Histopathology Findings:

liver (MA:0000358)

Histopath Description:

moderate lipidosis

Morphological Diagnosis:

Distribution: multifocal to coalescing; **Severity:** moderate; **MPATH Diagnosis:** steatosis MPATH:622

Definitive Diagnosis:

Hepatic lipidosis

retina (MA:0000276)

Histopath Description:


microphthalmia

Morphological Diagnosis:


Distribution: unilateral; **Severity:** mild;

Definitive Diagnosis:

Microphthalmia, unilateral



Eye, unilateral
microphthalmia, HE
1.25x



Peripheral nerve,
normal, 40x, HE

AnimalID: M00386490 (Female)

Histopathology Findings:

liver (MA:0000358)

Histopath Description:

severe lipidosis

Morphological Diagnosis:

Distribution: diffuse; **Severity:** severe; **MPATH Diagnosis:** steatosis MPATH:622

Definitive Diagnosis:

Hepatic lipidosis

retina (MA:0000276)

Histopath Description:


Involving one eye, there are clusters of external nuclear structures within the internal plexiform layer.

Morphological Diagnosis:

Distribution: Focal; **Severity:** mild;

Definitive Diagnosis:

Retinal dysplasia



Retina, dysplasia,
10x, HE

lymph node (MA:0000139)

Histopath Description:

The mesenteric lymph node is enlarged (greater than three-fold). There are multiple follicles with large germinal centers. The sinuses contain large numbers of mature lymphocytes.

Morphological Diagnosis:

Duration: Sub-acute; **Distribution:** Diffuse; **Severity:** moderate; **MPATH Diagnosis:** hyperplasia MPATH:134

Definitive Diagnosis:

Lymphoid hyperplasia.

Histopathology Comments:

The changes in the mesenteric lymph node are suggestive of draining of a regional inflammatory process. However, such a process was not observed in the tissues examined.


spleen (MA:0000141)

Histopath Description:

mild erythroid hyperplasia

Morphological Diagnosis:**Distribution:** multifocal; **Severity:** mild; **MPATH Diagnosis:** extramedullary hemopoiesis
MPATH:595**Definitive Diagnosis:**

Splenic erythroid hyperplasia



Peripheral nerve,
normal, 40x, HE

Report Summary and Recommendation:

All mice have ocular abnormality (one mouse with microphthalmia and three with retinal dysplasia). These abnormalities are considered background lesions in B6 strain. We did not find any lesions within central and peripheral nervous tissues and muscles to explain the tremors and hindlimb paralysis observed in this line. Image of hindlimb nerves and musculature are included. Other lesions are attributable to diet or strain background. Further review of the brain is considered (updated results will be communicated and appended)