



# CMHD Pathology Report



## CMHD Pathology Core

Toronto Centre for  
Phenogenomics  
25 Orde St. 3rd fl.  
Toronto, Ont. M5T 3H7  
Tel.(416) 586-8375  
Fax (416) 586-5993

contact: Dr. Susan  
Newbigging  
email:  
[newbigging@lunenfeld.ca](mailto:newbigging@lunenfeld.ca)

## Principle Investigator: Dr. Colin McKerlie

**ICSIG**  
Institute: Sick Kids  
Address:

ReportID: Report Date: November 23,  
2011  
Pathologist: H. Adissu

## Mouse Genetics Project

Wellcome Trust Sanger  
Institute  
Wellcome Trust Genome  
Campus  
Hinxton, Cambridge  
CB10 1SA  
UK

email:  
[MGPenquiries@sanger.ac.uk](mailto:MGPenquiries@sanger.ac.uk)  
[Mouse Portal](#)  
[Europhenome](#)

CMHD LabID: N11-388

## AnimalID: M00203045 Prdx6 hom

### Histopathology Findings:

#### retina (MA:0000276)

##### Histopath Description:

There are clusters of external nuclear structures within the layer of rods and cons.

##### Morphological Diagnosis:

**Distribution:** Focal; **Severity:** mild; **MPATH Diagnosis:** developmental and structural abnormality MPATH:55

##### Definitive Diagnosis:

Retinal dysplasia

#### eye (MA:0000261)

##### Histopath Description:

A 100 stalk of fibrous connective tissue containing a small artery in the center extends from the area of the optic disc towards the posterior vitreous. A small fragment of fibrous tissue is freely present within the vitreous anterior to this stalk (assumed to be extension of the stalk).

##### Morphological Diagnosis:

**MPATH Diagnosis:** developmental and structural abnormality MPATH:55

##### Definitive Diagnosis:

Persistent hyaloid artery

##### Histopathology Comments:

hyaloid artery remnant is a rare condition in which there remain some parts of the hyaloid artery. The posterior hyaloid vascular system of mice usually undergoes involution in the first month of life (Richard et al., 2000).

#### stomach (MA:0000353)

##### Histopath Description:

There are moderate numbers of neutrophils and a few plasma cells within the deep lamina propria and submucosa.

##### Morphological Diagnosis:

**Duration:** Chronic-active; **Distribution:** Multifocal; **Severity:** mild; **MPATH Diagnosis:** inflammation MPATH:212

##### Definitive Diagnosis:

Gastritis, suppurative

##### Histopathology Comments:

This lesion is most commonly associated with Helicobacter infection. Further investigation is suggested using histochemistry (Silver stain) or colony fecal PCR.

#### liver (MA:0000358)

##### Histopath Description:

There is diffuse hepatic lipidosis.

**Morphological Diagnosis:**

**Distribution:** Diffuse; **Severity:** moderate; **MPATH Diagnosis:** lipid deposition MPATH:42

**Definitive Diagnosis:**

Hepatic lipidosis

**Histopathology Comments:**

Hepatocellular vacuolar change of variable degree suggestive of lipidosis is present in all mice from WTSI, consistent with high lipid diet. See a representative image of diffuse hepatocellular lipidosis (Snip1 WT M00383263).

---

**AnimalID: M00203046 Prdx6 hom****Histopathology Findings:****liver (MA:0000358)****Histopath Description:**

lipid accumulation similar to M0018944 Adam 17 Het

**Morphological Diagnosis:**

**Distribution:** Diffuse; **Severity:** moderate; **MPATH Diagnosis:** lipid deposition MPATH:42

**Definitive Diagnosis:**

Hepatic lipidosis

**Histopathology Comments:**

Hepatocellular vacuolar change of variable degree suggestive of lipidosis is present in all mice from WTSI, consistent with high lipid diet. See a representative image of diffuse hepatocellular lipidosis (Snip1 WT M00383263).

**lymph node (MA:0000139)****Histopath Description:**

The mesenteric lymph node is enlarged (nearly 3x normal). There is a diffuse and marked increase in the paracortical and medullary areas and cellularity; these cells are organized as prominent chords, and they are larger and blast-like. Lymphoid follicles are increased in size and some have germinal centers.

**Morphological Diagnosis:**

**Distribution:** Diffuse; **Severity:** mild; **MPATH Diagnosis:** hyperplasia MPATH:134

**Definitive Diagnosis:**

Lymphoid hyperplasia

---

**AnimalID: M002036232 Prdx6 hom****Histopathology Findings:****liver (MA:0000358)****Histopath Description:**

The overall hepatic lobular architecture is normal. Nearly 80% of hepatocytes contain large (8-12 um in diameter) intracytoplasmic clear vacuoles (macrovesicular lipid).

**Morphological Diagnosis:**

**Distribution:** Multifocal; **Severity:** moderate; **MPATH Diagnosis:** lipid deposition MPATH:42

**Definitive Diagnosis:**

Hepatic lipidosis

**Histopathology Comments:**

Hepatocellular vacuolar change of variable degree suggestive of lipidosis is present in all mice from WTSI, consistent with high lipid diet.

**retina (MA:0000276)****Histopath Description:**

There are clusters of external nuclear structures within the layer of rods and cons.

**Morphological Diagnosis:**

**Distribution:** Focal; **Severity:** mild; **MPATH Diagnosis:** developmental and structural abnormality MPATH:55

**Definitive Diagnosis:**  
Retinal dysplasia

**lymph node (MA:0000139)**

**Histopath Description:**

The mesenteric and cervical lymph nodes are respectively enlarged nearly 4x and 2x normal and follicles are respectively prominent. Subcapsular sinuses contain large numbers of mature lymphocytes. There low numbers of neutrophils within the medullary stroma.

**Morphological Diagnosis:**

**Duration:** Sub-acute; **Distribution:** Diffuse; **Severity:** moderate; **MPATH Diagnosis:** hyperplasia MPATH:134

**Definitive Diagnosis:**

Lymphoid hyperplasia.

**Histopathology Comments:**

The changes in the mesenteric lymph node are suggestive of draining of a regional inflammatory process. However, such a process was not observed in the tissues examined.

---

**AnimalID: M002036233 Prdx6 hom**

**Histopathology Findings:**

**liver (MA:0000358)**

**Histopath Description:**

The overall hepatic lobular architecture is normal. Nearly 30% of hepatocytes contain large (8-12 um in diameter) intracytoplasmic clear vacuoles (macrovesicular lipid).

**Morphological Diagnosis:**

**Distribution:** Multifocal; **Severity:** mild; **MPATH Diagnosis:** lipid deposition MPATH:42

**Definitive Diagnosis:**

Hepatic lipidosi

**Histopathology Comments:**

Hepatocellular vacuolar change of variable degree suggestive of lipidosi is present in all mice from WTSI, consistent with high lipid diet.

**lymph node (MA:0000139)**

**Histopath Description:**

The mesenteric and cervical lymph nodes are respectively enlarged nearly 4x and 2x normal and follicles are respectively prominent. Subcapsular sinuses contain large numbers of mature lymphocytes. There low numbers of neutrophils within the medullary stroma.

**Morphological Diagnosis:**

**Duration:** Sub-acute; **Distribution:** Diffuse; **Severity:** moderate; **MPATH Diagnosis:** hyperplasia MPATH:134

**Definitive Diagnosis:**

Lymphoid hyperplasia.

**Histopathology Comments:**

The changes in the mesenteric lymph node are suggestive of draining of a regional inflammatory process. However, such a process was not observed in the tissues examined.

**brain (MA:0000168)**

**Histopath Description:**

There is a mild enlargement of the lateral ventricle.

**Morphological Diagnosis:**

**Severity:** mild; **MPATH Diagnosis:** hydrocephalus MPATH:639

**Definitive Diagnosis:**

hydrocephalus, lateral ventricle

**Histopathology Comments:**

Variable degree of hydrocephalus is observed in a proportion of wild type C57 Black 6 mice.

**Report Summary and Recommendation:**

Unique lesions in this line include retinal dysplasia in 2 of 4 mice and persistent hyaloid artery in 1 of 4 mice. Recently rd8 mutation in the Crb1 gene in C57BL/6N substrain is associated with multiple light-colored spots in the fundus of the eye that correspond histologically to retinal folds, pseudorosettes, and focal retinal dysplasia and degeneration. Hence retinal lesions in this line may be a background lesion. Other lesions are considered incidental and/or attributable to strain background.