


Gene: Prkd1

Colony prefix: DAJM

Allele: *Prkd1*^{em1(IMPC)Wtsi}

Allele type: Crispr/Cas9 mediated deletion

Allele information:



Mouse QC information

Loss of WT Allele (LOA) qPCR	pass	Mutation Sequence confirmed	pass
Mutant Specific SR-PCR	pass	Off-target analysis complete	na

Flanking sequence:

100bp 5' and 3' of the deletion

[TCAGAGGCATGGCTGGACCTAATCTATCGCTGCTGCTTTGAGTCTTGGCATT CAGTAGCAAATAAGTTCTTTCTA
CTATTTTGTGTTGGCTTCTAAACAT]
[TGGGTTATCTGACAAATGAGCTACGGATACAAATGGAGCTTTGGGATATTCAAAGTTCTACTGACACATAAGTTTC
TCATGGGATTACTGATCAACATTC]

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Report Generated on: 10-10-2017

Deletion sequence:

```
[TTGAAGACCTGATGGGCACCTTCATATCACTGGTGAGCTAAGTGTTTGTGGTATTAGCCAATTCATTCAAAT
AGCAAAATGGAGTGAGAGTCGACAGGGTAAGGATACTTGAGGGCATGGGTGTGATGATGGGTGTAGGTGGGA
ATGGCTGAGAGTCTGGGAGGAAGCACATACTAGAGATACATAGTAAAGTCCTCCTAGGTGGGAAAATGGACAAA
TTCTCCTCCGGTTTTTCATTCTTTCTTGAAACACCCTTATTTAGGTAGCCAAATCTCATGTGGAACACTACTCTACCTC
AAGTTGGTCTTAATGCATGATCCTTCTGTCTCAGCCTGACAGTATTGAGGTTACAGATGCAGGCCAGGGTTTTTCC
AAAATAAGTTTTGAAAGTTCCTGTAGACAGAGCTAAAAGCATCTCACGGTATAGCATATATGCTTTGTATTACAGTA
GCTTAGACCCAAAATTTCTTTAAAAGCAAATAATTAATCTACAGACATGGAAATCGGGATCTTACCTGACAGGA
CCACTTCAATGAGATCACCTCCTGGATATCACTTGCATTTTTACCAGCTGAAGGATGTTGTCCGGAGGCAGGAT
CGTGCCGAAAAAGCAGGATCTTATCATAGAGTCCATAGAAACCACATTCGGGGAACACTGCAAATGAGACACGAGA
GTTCAAATGTCACAATTTCCACAAATGTTTTGTCTTAAGAGAAAATTAACGAATGACCAAGGAGGTTTCTAATTCTT
TTTTTTTTTTTTCACATTTAAAGAAAGGGAAAAAAAATTCCTAGAAGTTTCTGTAGAGACATGATTAAGAATGGTACT
CAGCTCAGCAGAGTAGATGGCATAGAGATAAGAATGCATTTTCTCACCTGAAAGAGGAAATCAGCTTATAAAAAT
TTGTTACACATAACTTGACAATTTGACAAAGAAACAGGTCAGTCTGCTCTGACACACTGAAGACTAGATATGAGGC
AAGGAAGCCAAACACACTGTGGGCACAGAAGTGGAAATCTGGATACTCTGCCTCAAGACAGAACCAGGCTAGAGG
ACACACCTCTCCTTTGGCTCCCTCACTATTTATTTCCCTCGCTCTCCTTGAGGCTACAAATGCAGGGTCATATGA
ACCTGCATTACAC]
```

Guide RNAs used in initial experiment

Sequence	Chr	Chr Start	Chr End
ATATGAACCTGCATTACACTGGG	12	50490511	50490533
CCCTCGCTCTCCTTGAGGCTACA	12	50490477	50490499
AATGGAGTGAGAGTCGACAGGGG	12	50489463	50489485
CCTGATGGGCACCTTCATATCAC	12	50489389	50489411

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Report Generated on: 10-10-2017

Genotyping by end-point PCR

These mice may be genotyped through a combination of separate PCR reactions that detect the gene-specific wild type allele and a mutant allele-specific short range PCR. Interpretation of the consolidated results produces the genotype of the mice. In addition to the expected product, the mutant assay may also amplify the endogenous wild type sequence which will appear as a larger band on an agarose gel. The presence of this extra band will depend on the size of the original deletion.

PCRs primer pairs and expected size bands

Assay Type	Assay	Forward Primer	Reverse Primer	Expected Size Band (bp)
Standard PCR	Wild type	Prkd1_DF1	Prkd1_ER1	223
Standard PCR	Mutant	Prkd1_DF1	Prkd1_DR1	212

Primer sequences

Primer Name	Primer Sequence (5' > 3')
Prkd1_DF1	GGCATGGCTGGACCTAATCT
Prkd1_ER1	CACCCATGCCCTCAAGTATC
Prkd1_DR1	GCTTGACACTGGAAATGGAA

Reaction setup

Reagent	µl
DNA (~50-100 ng)	1
10x Buffer	1.5
MgCl ₂ (50 mM)	0.45
Platinum Taq (Invitrogen)	0.15
dNTPs (100 mM)	0.15
Primer 1 (10 µM)	0.3
Primer 2 (10 µM)	0.3
ddH ₂ O	11.15
Total	15

Amplification conditions

Step	Conditions	Time
1	94°C	5 min
2	94°C	30 sec
3	58°C	30 sec
4	72°C	1:30 sec
5	Go to '2' + 34 cycles	-
6	72°C	5 min
7	12°C	forever

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Report Generated on: 10-10-2017

Genotyping by loss of WT allele qPCR Assay (gene-specific assay)

The wild type loss of allele (LoA) qPCR assay uses a hydrolysis probe assay (for example Applied Biosystems TaqMan® technology) to determine the copy number of the wild type allele in a sample. Homozygotes will show no amplification, heterozygotes one copy and wild type mice will show two copies when compared to a wild type control.

The number of copies of the wildtype allele can be detected using a FAM-labelled custom qPCR TaqMan® assay. These are multiplexed with a VIC® labelled endogenous control assay (for example TaqMan® Copy Number Reference Assay, Mouse, Tfrc; Applied Biosystems part #4458366). Reference DNA controls of known genotypes should also be included to facilitate correct analysis.

Primers for LoA qPCR assay

Gene	Source	Forward Primer Seq.	Reverse Primer Seq.	Probe Primer Seq.
Prkd1	Life Technologies	CAGACATGGAAATCGGGATC TTA	CCTCCGACAACATCCT TCAG	ACAGGACCACTTCAATGAGATC ACCC

Reactions are performed in a 10µl volume using an Applied Biosystems 7900HT Fast Real-Time PCR System or Applied Biosystems Vii7 with DNA prepared using the Sample-to-SNPTM kit (Applied Biosystems) from mouse ear biopsies. GTXpress™ buffer is also used (Applied Biosystems).

Reagent	µl
2x GTXpress™ buffer	5
20x target assay	0.5
ddH2O	3
Tfrc endogenous 20x assay	0.5
DNA	1

Amplification conditions

Step	Conditions	Time
1	95°C	20 sec
2	95°C	10 sec
3	60°C	30 sec
4	Go to '2' + 34 cycles	-

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Report Generated on: 10-10-2017

Links to information and frequently asked questions

MGP mouse phenotype data:

<http://www.mousephenotype.org>

Useful publications

White, J.K., Gerdin, A.-K., Karp, N.A., Ryder, E., Buljan, M., Bussell, J.N., Salisbury, J., Clare, S., Ingham, N.J., Podrini, C., et al. (2013). Genome-wide Generation and Systematic Phenotyping of Knockout Mice Reveals New Roles for Many Genes. *Cell* 154, 452–464.

Mali P, Yang L, Esvelt KM, et al (2013) RNA-guided human genome engineering via Cas9. *Science* 339:823–6. doi: 10.1126/science.1232033

Jinek M, Chylinski K, Fonfara I, et al (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337:816–21. doi: 10.1126/science.1225829

Cong L, Ran FA, Cox D, et al (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science* 339:819–23. doi: 10.1126/science.1231143

Singh P, Schimenti JC, Bolcun-Filas E (2014) A Mouse Geneticist's Practical Guide to CRISPR Applications. *Genetics* genetics.114.169771–. doi: 10.1534/genetics.114.169771

Brandl C, Ortiz O, Röttig B, et al (2015) Creation of targeted genomic deletions using TALEN or CRISPR/Cas nuclease pairs in one-cell mouse embryos. *FEBS Open Bio* 5:26–35. doi: 10.1016/j.fob.2014.11.009

Zhou J, Wang J, Shen B, et al (2014) Dual sgRNAs facilitate CRISPR/Cas9 mediated mouse genome targeting. *FEBS J*. doi: 10.1111/febs.12735

Kraft K, Geuer S, Will AJ, et al (2015) Deletions, Inversions, Duplications: Engineering of Structural Variants using CRISPR/Cas in Mice. *Cell Rep*. doi: 10.1016/j.celrep.2015.01.016

Shen B, Zhang J, Wu H, et al (2013) Generation of gene-modified mice via Cas9/RNA-mediated gene targeting. *Cell Res* 23:720–3. doi: 10.1038/cr.2013.46

Wang H, Yang H, Shivalila CS, et al (2013) One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell* 153:910–8. doi: 10.1016/j.cell.2013.04.025

Yang H, Wang H, Shivalila CS, et al (2013) One-Step Generation of Mice Carrying Reporter and Conditional Alleles by CRISPR/Cas-Mediated Genome Engineering. *Cell* 154:1370–1379. doi: 10.1016/j.cell.2013.08.022

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Report Generated on: 10-10-2017