



**Abnormal Findings:** There are photoreceptors extending into the outer plexiform layer.

## EYE Phenotype




### Cornea:

**4/4.** Normal corneal epithelium, stroma, and endothelium.



### Anterior chamber:

**4/4.** The anterior chamber was of normal depth without cells, and the angle appeared open.



### Iris:


**4/4.** The iris showed normal pigmentation without rubeosis or pupillary membranes.

Gene: **Pfn1**

Genotype **+ / +**


Sanger Colony: MAOH

---




**Lens:**

**4/4.** No cataract was observed.




**Ciliary body:**

**4/4.** Normal stroma, pigmented and nonpigmented layers were present along with cilia.




**Vitreous:**

**4/4.** No abnormal opacities or cells.




**Retina:**

**3/4.** The retinal ganglion, inner nuclear layers are normal. There are some photoreceptors extending into the outer plexiform layer, which was seen in 3 eyes.



**Retinal pigment epithelium and Choroid:**

**4/4.** Normal pigmentation. Bruch's membrane is intact. No neovascular membranes were noted.



**Optic Nerve:**

**4/4.** The nerve is normal, but the section is not through the center as the nerve enters the eye.

**Methods.** 4 eyes from 2 male mice were enucleated by blunt dissection and fixed. Pupil-optic nerve sections were processed with hematoxylin and eosin, and standard images were captured under light microscopy for review.