

Wellcome Trust Sanger Institute Wellcome Trust Genome Campus Hinxton Cambridge CB10 1SA, U.K.

mouseinterest@sanger.ac.uk www.sanger.ac.uk

Eif2b2 (MCJP; EPD0168\_1\_D04)

Allele: Eif2b2<sup>tm2a(KOMP)Wtsi</sup>

Embryonic stem cell targeted: JM8.N4 Embryonic stem cell origin: C57BL/6N

Background used for Germ Line Transmission: C57BL/6N Taconic USA

Subsequent backcross background: C57BL/6N Taconic USA then inter cross from within Colony.

Genetic background: C57BL/6N Taconic USA;C57BL/6N

#### Coat Colour Information:

Non-Agouti (Black)

## **Breeding Performance and Lifespan:**

- Generally heterozygous mice from this colony conform to normal expectations of the background strain.
- Homozygous Lethal
- Unless timed embryonic stages are required heterozygous X wild-type matings should be used to maintain this colony.

### Bedding:

Aspen Chip derived from a Baltic supply - Supplier B&K Universal

# Diet:

Autoclavable Mouse Breeder Diet 5021 – A controlled constant-nutrient diet formulated to compensate for nutrient losses that occur during steam sterilization. Supplier Lab Diet www.labdiet.com

# **Husbandry:**

Cleaning frequency is based against cage numbers. Base changing is performed in a HEPA filtered change station which remains positive to the room environment. Gloved hands are disinfected between each cage. Diet is fed ad-libertum.

# **Housing System:**

Individual Ventilated Cages maintained at positive pressure to the room with an average of 60 HEPA filtered air changes per hour.

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Last updated: 04/02/2011



Wellcome Trust Sanger Institute Wellcome Trust Genome Campus Hinxton Cambridge CB10 1SA, U.K.

mouseinterest@sanger.ac.uk www.sanger.ac.uk

### **Further Information**

Homozygous Lethality: Please monitor www.sanger.ac.uk/mouseportal for confirmation.

As the mutant mouse strains generated by the Sanger MGP pass through the Sanger MGP primary phenotypic characterisation studies the information generated may be viewed through the Sanger Mouse Portal (<a href="www.sanger.ac.uk/mouseportal">www.sanger.ac.uk/mouseportal</a>) where a gene of interest may be searched for. A Heat Map of phenotyping by assay is also available to view. Information regarding homozygous lethality and fertility may also be sourced here if determined.

Early notification on phenotyping data may be received by subscribing to the MGP-Early-Phenotyping-Alert.

Phenotype enquiries may be made through the contact <a href="MGPEnquiries@sanger.ac.uk">MGPEnquiries@sanger.ac.uk</a> .

A further source of phenotype information is the Europhenome Mouse Phenotyping Resource (www.europhenome.org)

Information regarding availability of knockout mouse resources may be queried at the International Knockout Mouse Consortium (IKMC; www.knockoutmouse.org).

Information relating to the knockout programmes may be found at the IKMC Knowledgebase, currently in development (<a href="https://www.knockoutmouse.org/content/ikmc-prototypes">www.knockoutmouse.org/content/ikmc-prototypes</a>).

Information about targeting strategies may also be found at the IKMC website (<a href="https://www.knockoutmouse.org/about/targeting-strategies">www.knockoutmouse.org/about/targeting-strategies</a>).

### **Supplementary Notes:**

A strain with conditional potential may be mated to a FLPeR strain to delete the region of DNA flanked by FRT sites which includes the selectable cassette. Genotype confirmation of the deletion event should be carried out. The resulting conditional strain may be further progressed to incorporate Cre deleter transgenes to remove critical coding region(s) as desired.

#### References

Widespread recombinase expression using FLPeR (flipper) mice. Farley FW, Soriano P, Steffen LS, Dymecki SM. (2000). Genesis 28 (3-4),106-110.

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Last updated: 04/02/2011