

Wellcome Trust Sanger Institute Wellcome Trust Genome Campus Hinxton Cambridge CB10 1SA, U.K.

mouseinterest@sanger.ac.uk www.sanger.ac.uk

# Aldh1b1 (MGER; EPD0537\_4\_A09)

Allele: Aldh1b1<sup>tm2a(EUCOMM)Wtsi</sup>

Embryonic stem cell targeted: JM8A3.N1 Embryonic stem cell origin: C57BL/6N-A<sup>tm1Brd</sup>/a

Background used for Germ Line Transmission: C57BL/6N

Subsequent backcross background: C57BL/6N and inter cross from within Colony

Genetic background: C57BL/6N; C57BL/6N-A<sup>tm1Brd</sup>/a

#### Coat Colour Information:

Agouti and Black

#### **Breeding Performance and Lifespan:**

- Generally heterozygous mice from this colony conform to normal expectations of the background strain.
  - For maintenance of our colonies we pay particular attention to the age of the mating pairs and the resulting litters. In our experience the C57BL/6N substrain used to establish and progress this colony has shown some characteristics such as poor breeding, high preweaning mortality rates and failure to breed beyond three litters. We believe disturbance of litters has a detrimental effect on the mating pair. For our core and mutant colonies we have actively reduced our intervention with the mice. Daily observations, health checks, cleaning and cage movement is minimised in litters under 14 days of age.
- The colony was observed to produce a mouse with albino coat colour in second generation sibling x sibling breeding. This colony was subsequently investigated for the potential introduction of the genetic background of the blastocyst (C57BL/6Brd-*Tyr*<sup>c-Brd</sup>) used at microinjection into the pedigree of the propagated generations from the G1 stage of colony expansion. Evidence from tyrosinase-c mutation SNP analysis carried out for this colony suggests that the genetic background from the blastocyst was introduced into the lineage of some of the mice within the colony. The branches of the colony identified to carry the contribution from the blastocyst were discontinued and the colony reestablished from the lineage identified to be unaffected.
- Viability at Weaning Homozygous viable.

## Bedding:

Aspen Chip derived from a Baltic supply - Supplier B&K Universal

#### Diet:

Autoclavable Mouse Breeder Diet 5021 – A controlled constant-nutrient diet formulated to compensate for nutrient losses that occur during steam sterilization. Supplier Lab Diet <a href="https://www.labdiet.com">www.labdiet.com</a>

### **Husbandry:**

Cleaning frequency is based against cage occupancy and technician assessed level of soiling. Base changing is performed in a HEPA filtered change station which remains positive to the room environment. Gloved hands are disinfected between each cage. Diet is fed ad-libitum.

 $This \ technical \ data \ sheet \ and \ information \ ("Datasheet") \ is \ supplied \ by \ Genome \ Research \ Limited \ ("GRL").$ 

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Last updated: 20/05/2014



Wellcome Trust Sanger Institute Wellcome Trust Genome Campus Hinxton Cambridge CB10 1SA, U.K.

mouseinterest@sanger.ac.uk www.sanger.ac.uk

## **Housing System:**

Individual Ventilated Cages maintained at positive pressure to the room with an average of 60 HEPA filtered air changes per hour.

## **Further Information**

Whilst all reasonable effort is made to verify the mouse line and verify the individual mouse genotype at shipment, we recommend this is confirmed by the recipient.

The 3' loxP is in 3'-UTR.

No expected effects for the tm1b allele.

Potential to exhibit a phenotype for tm1c allele if the 3'loxP influences mRNA stability.

Sanger MGP mutant mouse lines are mouse lines in development; information about breeding and phenotyping characteristics may be incomplete.

As the mutant mouse strains progress through the Sanger MGP primary phenotypic characterisation, the information gathered may be viewed through the Sanger Mouse Portal (<a href="www.sanger.ac.uk/mouseportal">www.sanger.ac.uk/mouseportal</a>) and the International Mouse Phenotyping Consortium (IMPC; <a href="www.mousephenotype.org">www.mousephenotype.org</a>).

Information supplied here is current as of the date indicated below.

Please consult the Sanger MGP Mouse Resource Portal and IMPC for progressive updates on colony information such as Viability at weaning, Fertility, General Observations.

Contact <a href="mailto:MGPEnquiries@sanger.ac.uk">MGPEnquiries@sanger.ac.uk</a>

Early notification on phenotyping data may be received by subscribing to the MGP-Early-Phenotyping-Alert (<a href="https://www.sanger.ac.uk/mouseportal">www.sanger.ac.uk/mouseportal</a>).

Phenotype enquiries may be made through the contact MGPEnquiries@sanger.ac.uk.

Details of the colony quality control tests performed for a specific mouse line may be observed through the International Mouse Phenotyping Consortium (IMPC; <a href="www.mousephenotype.org">www.mousephenotype.org</a>), searching for your gene and follow the link from 'Product Details' for the mouse strain of interest.

General Descriptions of the mouse strain quality control (QC) assays. www.i-dcc.org/kb/25

General information about structure of IMPC alleles and their derivatives <a href="https://www.mousephenotype.org/martsearch\_ikmc\_project/about/targeting-strategies">www.mousephenotype.org/martsearch\_ikmc\_project/about/targeting-strategies</a>

Guidelines for converting alleles www.i-dcc.org/kb/entry/105

International Mouse Phenotyping Consortium (IMPC) Mouse Resources www.mousephenotype.org

IKMC Knowledgebase www.i-dcc.org/kb

 $This \ technical \ data \ sheet \ and \ information \ ("Datasheet") \ is \ supplied \ by \ Genome \ Research \ Limited \ ("GRL").$ 

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Last updated: 20/05/2014



Wellcome Trust Sanger Institute Wellcome Trust Genome Campus Hinxton Cambridge CB10 1SA, U.K.

mouseinterest@sanger.ac.uk www.sanger.ac.uk

#### **Publications**

Bradley A, Anastassiadis K, Ayadi A, Battey JF, Bell C, Birling M-C, Bottomley J, Brown SD, Bürger A, Bult CJ, Bushell W, Collins FS, Desaintes C, Doe B, Economides A, Eppig JT, Finnell RH, Fletcher C, Fray M, Frendewey D, *et al.* (2012) The mammalian gene function resource: the international knockout mouse consortium. *Mamm. Genome*, **23**, 580-586.

Liang, Q., Conte, N., Skarnes, W. C. and Bradley, A. (2008). Extensive genomic copy number variation in embryonic stem cells. *Proc. Nat. Acad. Sci.*, **105** (11), 17453-17456.

Pettitt SJ, Liang Q, Rairdan XY, Moran JL, Prosser HM, Beier DR, Lloyd KC, Bradley A & Skarnes WC (2009) Agouti C57BL/6N embryonic stem cells for mouse genetic resources. *Nature methods*, **6**, 493-495.

Ryder, E., Doe, B., Gleeson, D., Houghton, R., Dalvi, P., Grau, E., ... Ramirez-Solis, R. (2013). Rapid conversion of EUCOMM/KOMP-CSD alleles in mouse embryos using a cell-permeable Cre recombinase. *Transgenic research*. 23(1), 177–185.

Ryder E, Gleeson D, Sethi D, Vyas S, Miklejewska E, Dalvi P, Habib B, Cook R, Hardy M, Jhaveri K, Bottomley J, Wardle-Jones H, Bussell JN, Houghton R, Salisbury J, Skarnes WC; Sanger Mouse Genetics Project, Ramirez-Solis R. (2013). Molecular characterization of mutant mouse strains generated from the EUCOMM/KOMP-CSD ES cell resource. *Mamm. Genome*, **24**, 286–294.

Skarnes, W.C., Rosen, B., West, A.P., Koutsourakis, M., Bushell, W., Iyer, V., Mujica, A.O., Thomas, M., Harrow, J., Cox, T. *et al.* (2011) A conditional knockout resource for the genomewide study of mouse gene function. *Nature*, **474**, 337-342.

White, J. K., Gerdin, A.-K., Karp, N. A., Ryder, E., Buljan, M., Bussell, J. N., Salisbury, J., *et al.* (2013). Genome-wide Generation and Systematic Phenotyping of Knockout Mice Reveals New Roles for Many Genes. *Cell*, **154**(2), 452–464.

#### Additional Useful Publications

Birling M.C., Gofflot F. and Warot X. (2009). Site-specific recombinases for manipulation of the mouse genome. *Methods Mol. Biol.*, **561**, 245-263. Review.

Farley FW, Soriano P, Steffen LS, Dymecki SM. (2000). Widespread recombinase expression using FLPeR (flipper) mice. *Genesis*, **28** (3-4), 106-110.

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").

Although reasonable care is taken in the preparation of this Datasheet, GRL gives no warranties express or implied for any use of the Datasheet or for the accuracy of the Datasheet. GRL assumes no responsibility or liability for any decisions based upon the Datasheet. Without limiting the foregoing the Datasheet was prepared for mice supplied directly from GRL and where copies of this Datasheet are available from third party repositories or distribution centres ("Third Parties") GRL shall not be liable for any inconsistency between the mouse strain supplied by the Third Party and the Datasheet howsoever arising.

Last updated: 20/05/2014