

References

- (2004) Finishing the euchromatic sequence of the human genome. *Nature* 431(7011): 931-945.
- (2004) The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science* 306(5696): 636-640.
- Adlam M, Siu G (2003) Hierarchical interactions control CD4 gene expression during thymocyte development. *Immunity* 18(2): 173-184.
- Affolter M, Montagne J, Walldorf U, Groppe J, Kloter U et al. (1994) The Drosophila SRF homolog is expressed in a subset of tracheal cells and maps within a genomic region required for tracheal development. *Development* 120(4): 743-753.
- Agalioti T, Chen G, Thanos D (2002) Deciphering the transcriptional histone acetylation code for a human gene. *Cell* 111(3): 381-392.
- Agger K, Cloos PA, Christensen J, Pasini D, Rose S et al. (2007) UTX and JMJD3 are histone H3K27 demethylases involved in HOX gene regulation and development. *Nature* 449(7163): 731-734.
- Ahmad K, Henikoff S (2001) Centromeres are specialized replication domains in heterochromatin. *J Cell Biol* 153(1): 101-110.
- Ahmad K, Henikoff S (2002) The histone variant H3.3 marks active chromatin by replication-independent nucleosome assembly. *Mol Cell* 9(6): 1191-1200.
- Ahn SH, Cheung WL, Hsu JY, Diaz RL, Smith MM et al. (2005) Sterile 20 kinase phosphorylates histone H2B at serine 10 during hydrogen peroxide-induced apoptosis in *S. cerevisiae*. *Cell* 120(1): 25-36.
- Akasaka T, van Lohuizen M, van der Lugt N, Mizutani-Koseki Y, Kanno M et al. (2001) Mice doubly deficient for the Polycomb Group genes *Mel18* and *Bmi1* reveal synergy and requirement for maintenance but not initiation of Hox gene expression. *Development* 128(9): 1587-1597.
- Allen GC, Spiker S, Thompson WF (2000) Use of matrix attachment regions (MARs) to minimize transgene silencing. *Plant Mol Biol* 43(2-3): 361-376.
- Allfrey VG, Faulkner R, Mirsky AE (1964) Acetylation and Methylation of Histones and Their Possible Role in the Regulation of Rna Synthesis. *Proc Natl Acad Sci U S A* 51: 786-794.
- Amati B, Gasser SM (1990) Drosophila scaffold-attached regions bind nuclear scaffolds and can function as ARS elements in both budding and fission yeasts. *Mol Cell Biol* 10(10): 5442-5454.
- Anderson CL, Abraham GN (1980) Characterization of the Fc receptor for IgG on a human macrophage cell line, U937. *J Immunol* 125(6): 2735-2741.
- Angelov D, Molla A, Perche PY, Hans F, Cote J et al. (2003) The histone variant macroH2A interferes with transcription factor binding and SWI/SNF nucleosome remodeling. *Mol Cell* 11(4): 1033-1041.
- Aoyagi S, Narlikar G, Zheng C, Sif S, Kingston RE et al. (2002) Nucleosome remodeling by the human SWI/SNF complex requires transient global disruption of histone-DNA interactions. *Mol Cell Biol* 22(11): 3653-3662.
- Aplan PD, Begley CG, Bertness V, Nussmeier M, Ezquerro A et al. (1990) The SCL gene is formed from a transcriptionally complex locus. *Mol Cell Biol* 10(12): 6426-6435.

- Aronow BJ, Silbiger RN, Dusing MR, Stock JL, Yager KL et al. (1992) Functional analysis of the human adenosine deaminase gene thymic regulatory region and its ability to generate position-independent transgene expression. *Mol Cell Biol* 12(9): 4170-4185.
- Attema JL, Papathanasiou P, Forsberg EC, Xu J, Smale ST et al. (2007) Epigenetic characterization of hematopoietic stem cell differentiation using miniChIP and bisulfite sequencing analysis. *Proc Natl Acad Sci U S A* 104(30): 12371-12376.
- Ayer DE, Lawrence QA, Eisenman RN (1995) Mad-Max transcriptional repression is mediated by ternary complex formation with mammalian homologs of yeast repressor Sin3. *Cell* 80(5): 767-776.
- Azuara V, Perry P, Sauer S, Spivakov M, Jorgensen HF et al. (2006) Chromatin signatures of pluripotent cell lines. *Nat Cell Biol* 8(5): 532-538.
- Babu MM, Luscombe NM, Aravind L, Gerstein M, Teichmann SA (2004) Structure and evolution of transcriptional regulatory networks. *Curr Opin Struct Biol* 14(3): 283-291.
- Bailey TL, Elkan C (1995) The value of prior knowledge in discovering motifs with MEME. *Proc Int Conf Intell Syst Mol Biol* 3: 21-29.
- Bajic VB, Tan SL, Suzuki Y, Sugano S (2004) Promoter prediction analysis on the whole human genome. *Nat Biotechnol* 22(11): 1467-1473.
- Bajic VB, Brent MR, Brown RH, Frankish A, Harrow J et al. (2006) Performance assessment of promoter predictions on ENCODE regions in the EGASP experiment. *Genome Biol* 7 Suppl 1: S3 1-13.
- Balhoff JP, Wray GA (2005) Evolutionary analysis of the well characterized endo16 promoter reveals substantial variation within functional sites. *Proc Natl Acad Sci U S A* 102(24): 8591-8596.
- Banerji J, Rusconi S, Schaffner W (1981) Expression of a beta-globin gene is enhanced by remote SV40 DNA sequences. *Cell* 27(2 Pt 1): 299-308.
- Banerji J, Olson L, Schaffner W (1983) A lymphocyte-specific cellular enhancer is located downstream of the joining region in immunoglobulin heavy chain genes. *Cell* 33(3): 729-740.
- Bannister AJ, Zegerman P, Partridge JF, Miska EA, Thomas JO et al. (2001) Selective recognition of methylated lysine 9 on histone H3 by the HP1 chromo domain. *Nature* 410(6824): 120-124.
- Bannister AJ, Kouzarides T (2005) Reversing histone methylation. *Nature* 436(7054): 1103-1106.
- Bannister AJ, Schneider R, Myers FA, Thorne AW, Crane-Robinson C et al. (2005) Spatial distribution of di- and tri-methyl lysine 36 of histone H3 at active genes. *J Biol Chem* 280(18): 17732-17736.
- Barski A, Cuddapah S, Cui K, Roh TY, Schones DE et al. (2007) High-resolution profiling of histone methylations in the human genome. *Cell* 129(4): 823-837.
- Battaglioli E, Andres ME, Rose DW, Chenoweth JG, Rosenfeld MG et al. (2002) REST repression of neuronal genes requires components of the hSWI.SNF complex. *J Biol Chem* 277(43): 41038-41045.
- Becker PB (2002) Nucleosome sliding: facts and fiction. *EMBO J* 21(18): 4749-4753.
- Begley CG, Green AR (1999) The SCL gene: from case report to critical hematopoietic regulator. *Blood* 93(9): 2760-2770.

- Bell AC, West AG, Felsenfeld G (1999) The protein CTCF is required for the enhancer blocking activity of vertebrate insulators. *Cell* 98(3): 387-396.
- Bell AC, Felsenfeld G (2000) Methylation of a CTCF-dependent boundary controls imprinted expression of the *Igf2* gene. *Nature* 405(6785): 482-485.
- Bell AC, West AG, Felsenfeld G (2001) Insulators and boundaries: versatile regulatory elements in the eukaryotic. *Science* 291(5503): 447-450.
- Bentley DR (2006) Whole-genome re-sequencing. *Curr Opin Genet Dev* 16(6): 545-552.
- Berger SL (2002) Histone modifications in transcriptional regulation. *Curr Opin Genet Dev* 12(2): 142-148.
- Berger JA, Hautaniemi S, Jarvinen AK, Edgren H, Mitra SK et al. (2004) Optimized LOWESS normalization parameter selection for DNA microarray data. *BMC Bioinformatics* 5: 194.
- Bernard O, Azogui O, Lecointe N, Mugneret F, Berger R et al. (1992) A third *tal-1* promoter is specifically used in human T cell leukemias. *J Exp Med* 176(4): 919-925.
- Bernstein BE, Humphrey EL, Erlich RL, Schneider R, Bouman P et al. (2002) Methylation of histone H3 Lys 4 in coding regions of active genes. *Proc Natl Acad Sci U S A* 99(13): 8695-8700.
- Bernstein BE, Liu CL, Humphrey EL, Perlstein EO, Schreiber SL (2004) Global nucleosome occupancy in yeast. *Genome Biol* 5(9): R62.
- Bernstein BE, Kamal M, Lindblad-Toh K, Bekiranov S, Bailey DK et al. (2005) Genomic maps and comparative analysis of histone modifications in human and mouse. *Cell* 120(2): 169-181.
- Bernstein BE, Mikkelsen TS, Xie X, Kamal M, Huebert DJ et al. (2006) A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* 125(2): 315-326.
- Bernstein BE, Meissner A, Lander ES (2007) The mammalian epigenome. *Cell* 128(4): 669-681.
- Bhatia M (2007) Hematopoiesis from human embryonic stem cells. *Ann N Y Acad Sci* 1106: 219-222.
- Bieda M, Xu X, Singer MA, Green R, Farnham PJ (2006) Unbiased location analysis of E2F1-binding sites suggests a widespread role for E2F1 in the human genome. *Genome Res* 16(5): 595-605.
- Biggin MD (1999) Ultraviolet cross-linking assay to measure sequence-specific DNA binding in vivo. *Methods Enzymol* 304: 496-515.
- Bird A (2002) DNA methylation patterns and epigenetic memory. *Genes Dev* 16(1): 6-21.
- Birney E, Durbin R (2000) Using GeneWise in the *Drosophila* annotation experiment. *Genome Res* 10(4): 547-548.
- Birney E, Stamatoyannopoulos JA, Dutta A, Guigo R, Gingeras TR et al. (2007) Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 447(7146): 799-816.
- Birve A, Sengupta AK, Beuchle D, Larsson J, Kennison JA et al. (2001) *Su(z)12*, a novel *Drosophila* Polycomb group gene that is conserved in vertebrates and plants. *Development* 128(17): 3371-3379.

- Bitoun E, Oliver PL, Davies KE (2007) The mixed-lineage leukemia fusion partner AF4 stimulates RNA polymerase II transcriptional elongation and mediates coordinated chromatin remodeling. *Hum Mol Genet* 16(1): 92-106.
- Bjerkvig R, Tysnes BB, Aboody KS, Najbauer J, Terzis AJ (2005) Opinion: the origin of the cancer stem cell: current controversies and new insights. *Nat Rev Cancer* 5(11): 899-904.
- Blais A, Tsikitis M, Acosta-Alvear D, Sharan R, Kluger Y et al. (2005) An initial blueprint for myogenic differentiation. *Genes Dev* 19(5): 553-569.
- Blanchette M, Tompa M (2003) FootPrinter: A program designed for phylogenetic footprinting. *Nucleic Acids Res* 31(13): 3840-3842.
- Bode J, Benham C, Knopp A, Mielke C (2000) Transcriptional augmentation: modulation of gene expression by scaffold/matrix-attached regions (S/MAR elements). *Crit Rev Eukaryot Gene Expr* 10(1): 73-90.
- Boffelli D, McAuliffe J, Ovcharenko D, Lewis KD, Ovcharenko I et al. (2003) Phylogenetic shadowing of primate sequences to find functional regions of the human genome. *Science* 299(5611): 1391-1394.
- Bosher JM, Totty NF, Hsuan JJ, Williams T, Hurst HC (1996) A family of AP-2 proteins regulates c-erbB-2 expression in mammary carcinoma. *Oncogene* 13(8): 1701-1707.
- Botuyan MV, Lee J, Ward IM, Kim JE, Thompson JR et al. (2006) Structural basis for the methylation state-specific recognition of histone H4-K20 by 53BP1 and Crb2 in DNA repair. *Cell* 127(7): 1361-1373.
- Boyer LA, Latek RR, Peterson CL (2004) The SANT domain: a unique histone-tail-binding module? *Nat Rev Mol Cell Biol* 5(2): 158-163.
- Boyer LA, Lee TI, Cole MF, Johnstone SE, Levine SS et al. (2005) Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 122(6): 947-956.
- Boyer LA, Plath K, Zeitlinger J, Brambrink T, Medeiros LA et al. (2006) Polycomb complexes repress developmental regulators in murine embryonic stem cells. *Nature* 441(7091): 349-353.
- Bracken AP, Dietrich N, Pasini D, Hansen KH, Helin K (2006) Genome-wide mapping of Polycomb target genes unravels their roles in cell fate transitions. *Genes Dev* 20(9): 1123-1136.
- Bradley A (1990) Embryonic stem cells: proliferation and differentiation. *Curr Opin Cell Biol* 2(6): 1013-1017.
- Brasset E, Vaury C (2005) Insulators are fundamental components of the eukaryotic genomes. *Heredity* 94(6): 571-576.
- Briggs SD, Bryk M, Strahl BD, Cheung WL, Davie JK et al. (2001) Histone H3 lysine 4 methylation is mediated by Set1 and required for cell growth and rDNA silencing in *Saccharomyces cerevisiae*. *Genes Dev* 15(24): 3286-3295.
- Brown KE, Baxter J, Graf D, Merckenschlager M, Fisher AG (1999) Dynamic repositioning of genes in the nucleus of lymphocytes preparing for cell division. *Mol Cell* 3(2): 207-217.

- Bruce AW, Donaldson IJ, Wood IC, Yerbury SA, Sadowski MI et al. (2004) Genome-wide analysis of repressor element 1 silencing transcription factor/neuron-restrictive silencing factor (REST/NRSF) target genes. *Proc Natl Acad Sci U S A* 101(28): 10458-10463.
- Bryne JC, Valen E, Tang MH, Marstrand T, Winther O et al. (2008) JASPAR, the open access database of transcription factor-binding profiles: new content and tools in the 2008 update. *Nucleic Acids Res* 36(Database issue): D102-106.
- Buck MJ, Nobel AB, Lieb JD (2005) ChIPOTle: a user-friendly tool for the analysis of ChIP-chip data. *Genome Biol* 6(11): R97.
- Bulger M, Groudine M (1999) Looping versus linking: toward a model for long-distance gene activation. *Genes Dev* 13(19): 2465-2477.
- Bulger M, van Doorninck JH, Saitoh N, Telling A, Farrell C et al. (1999) Conservation of sequence and structure flanking the mouse and human beta-globin loci: the beta-globin genes are embedded within an array of odorant receptor genes. *Proc Natl Acad Sci U S A* 96(9): 5129-5134.
- Burcin M, Arnold R, Lutz M, Kaiser B, Runge D et al. (1997) Negative protein 1, which is required for function of the chicken lysozyme gene silencer in conjunction with hormone receptors, is identical to the multivalent zinc finger repressor CTCF. *Mol Cell Biol* 17(3): 1281-1288.
- Burge C, Karlin S (1997) Prediction of complete gene structures in human genomic DNA. *J Mol Biol* 268(1): 78-94.
- Burgess-Beusse B, Farrell C, Gaszner M, Litt M, Mutskov V et al. (2002) The insulation of genes from external enhancers and silencing chromatin. *Proc Natl Acad Sci U S A* 99 Suppl 4: 16433-16437.
- Cai HN, Shen P (2001) Effects of cis arrangement of chromatin insulators on enhancer-blocking activity. *Science* 291(5503): 493-495.
- Cao R, Wang L, Wang H, Xia L, Erdjument-Bromage H et al. (2002) Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science* 298(5595): 1039-1043.
- Cao R, Zhang Y (2004) The functions of E(Z)/EZH2-mediated methylation of lysine 27 in histone H3. *Curr Opin Genet Dev* 14(2): 155-164.
- Capelson M, Corces VG (2005) The ubiquitin ligase dTopors directs the nuclear organization of a chromatin insulator. *Mol Cell* 20(1): 105-116.
- Carninci P, Kasukawa T, Katayama S, Gough J, Frith MC et al. (2005) The transcriptional landscape of the mammalian genome. *Science* 309(5740): 1559-1563.
- Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA et al. (2005) Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell* 122(1): 33-43.
- Carroll JS, Meyer CA, Song J, Li W, Geistlinger TR et al. (2006) Genome-wide analysis of estrogen receptor binding sites. *Nat Genet* 38(11): 1289-1297.

- Carrozza MJ, Li B, Florens L, Suganuma T, Swanson SK et al. (2005) Histone H3 methylation by Set2 directs deacetylation of coding regions by Rpd3S to suppress spurious intragenic transcription. *Cell* 123(4): 581-592.
- Cawley S, Bekiranov S, Ng HH, Kapranov P, Sekinger EA et al. (2004) Unbiased mapping of transcription factor binding sites along human chromosomes 21 and 22 points to widespread regulation of noncoding RNAs. *Cell* 116(4): 499-509.
- Celeste A, Fernandez-Capetillo O, Kruhlak MJ, Pilch DR, Staudt DW et al. (2003) Histone H2AX phosphorylation is dispensable for the initial recognition of DNA breaks. *Nat Cell Biol* 5(7): 675-679.
- Chadwick BP, Willard HF (2001) Histone H2A variants and the inactive X chromosome: identification of a second macroH2A variant. *Hum Mol Genet* 10(10): 1101-1113.
- Chakalova L, Carter D, Debrand E, Goyenechea B, Horton A et al. (2005) Developmental regulation of the beta-globin gene locus. *Prog Mol Subcell Biol* 38: 183-206.
- Chambeyron S, Da Silva NR, Lawson KA, Bickmore WA (2005) Nuclear re-organisation of the Hoxb complex during mouse embryonic development. *Development* 132(9): 2215-2223.
- Chen D, Ma H, Hong H, Koh SS, Huang SM et al. (1999) Regulation of transcription by a protein methyltransferase. *Science* 284(5423): 2174-2177.
- Cho DH, Thienes CP, Mahoney SE, Analau E, Filippova GN et al. (2005) Antisense transcription and heterochromatin at the DM1 CTG repeats are constrained by CTCF. *Mol Cell* 20(3): 483-489.
- Chow CM, Georgiou A, Szutorisz H, Maia e Silva A, Pombo A et al. (2005) Variant histone H3.3 marks promoters of transcriptionally active genes during mammalian cell division. *EMBO Rep* 6(4): 354-360.
- Christensen J, Agger K, Cloos PA, Pasini D, Rose S et al. (2007) RBP2 belongs to a family of demethylases, specific for tri- and dimethylated lysine 4 on histone 3. *Cell* 128(6): 1063-1076.
- Chung JH, Whiteley M, Felsenfeld G (1993) A 5' element of the chicken beta-globin domain serves as an insulator in human erythroid cells and protects against position effect in *Drosophila*. *Cell* 74(3): 505-514.
- Cloos PA, Christensen J, Agger K, Maiolica A, Rappsilber J et al. (2006) The putative oncogene GASC1 demethylates tri- and dimethylated lysine 9 on histone H3. *Nature* 442(7100): 307-311.
- Cockerill PN (2000) Identification of DNaseI hypersensitive sites within nuclei. *Methods Mol Biol* 130: 29-46.
- Collins FS, Green ED, Guttmacher AE, Guyer MS (2003) A vision for the future of genomics research. *Nature* 422(6934): 835-847.
- Cora D, Herrmann C, Dieterich C, Di Cunto F, Provero P et al. (2005) Ab initio identification of putative human transcription factor binding sites by comparative genomics. *BMC Bioinformatics* 6: 110.
- Cosgrove MS (2007) Histone proteomics and the epigenetic regulation of nucleosome mobility. *Expert Rev Proteomics* 4(4): 465-478.

- Costanzi C, Stein P, Worrada DM, Schultz RM, Pehrson JR (2000) Histone macroH2A1 is concentrated in the inactive X chromosome of female preimplantation mouse embryos. *Development* 127(11): 2283-2289.
- Craig JM (2005) Heterochromatin--many flavours, common themes. *Bioessays* 27(1): 17-28.
- Crawford GE, Holt IE, Mullikin JC, Tai D, Blakesley R et al. (2004) Identifying gene regulatory elements by genome-wide recovery of DNase hypersensitive sites. *Proc Natl Acad Sci U S A* 101(4): 992-997.
- Crawford GE, Davis S, Scacheri PC, Renaud G, Halawi MJ et al. (2006) DNase-chip: a high-resolution method to identify DNase I hypersensitive sites using tiled microarrays. *Nat Methods* 3(7): 503-509.
- Crawford GE, Holt IE, Whittle J, Webb BD, Tai D et al. (2006 b) Genome-wide mapping of DNase hypersensitive sites using massively parallel signature sequencing (MPSS). *Genome Res* 16(1): 123-131.
- Cuthbert GL, Daujat S, Snowden AW, Erdjument-Bromage H, Hagiwara T et al. (2004) Histone deimination antagonizes arginine methylation. *Cell* 118(5): 545-553.
- Czermin B, Melfi R, McCabe D, Seitz V, Imhof A et al. (2002) Drosophila enhancer of Zeste/ESC complexes have a histone H3 methyltransferase activity that marks chromosomal Polycomb sites. *Cell* 111(2): 185-196.
- Dahl JA, Collas P (2007) Q2ChIP, a quick and quantitative chromatin immunoprecipitation assay, unravels epigenetic dynamics of developmentally regulated genes in human carcinoma cells. *Stem Cells* 25(4): 1037-1046.
- Damelin M, Simon I, Moy TI, Wilson B, Komili S et al. (2002) The genome-wide localization of Rsc9, a component of the RSC chromatin-remodeling complex, changes in response to stress. *Mol Cell* 9(3): 563-573.
- Dannenbergh JH, David G, Zhong S, van der Torre J, Wong WH et al. (2005) mSin3A corepressor regulates diverse transcriptional networks governing normal and neoplastic growth and survival. *Genes Dev* 19(13): 1581-1595.
- Davuluri RV, Grosse I, Zhang MQ (2001) Computational identification of promoters and first exons in the human genome. *Nat Genet* 29(4): 412-417.
- De Gobbi M, Viprakasit V, Hughes JR, Fisher C, Buckle VJ et al. (2006) A regulatory SNP causes a human genetic disease by creating a new transcriptional promoter. *Science* 312(5777): 1215-1217.
- De Gobbi M, Anguita E, Hughes J, Sloane-Stanley JA, Sharpe JA et al. (2007) Tissue-specific histone modification and transcription factor binding in alpha globin gene expression. *Blood* 110(13): 4503-4510.
- De la Serna IL, Ohkawa Y, Imbalzano AN (2006) Chromatin remodelling in mammalian differentiation: lessons from ATP-dependent remodellers. *Nat Rev Genet* 7(6): 461-473.

- De Nadal E, Zapater M, Alepuz PM, Sumoy L, Mas G et al. (2004) The MAPK Hog1 recruits Rpd3 histone deacetylase to activate osmoresponsive genes. *Nature* 427(6972): 370-374.
- De Napoles M, Mermoud JE, Wakao R, Tang YA, Endoh M et al. (2004) Polycomb group proteins Ring1A/B link ubiquitylation of histone H2A to heritable gene silencing and X inactivation. *Dev Cell* 7(5): 663-676.
- Defossez PA, Kelly KF, Filion GJ, Perez-Torrado R, Magdinier F et al. (2005) The human enhancer blocker CTC-binding factor interacts with the transcription factor Kaiso. *J Biol Chem* 280(52): 43017-43023.
- Dekker J, Rippe K, Dekker M, Kleckner N (2002) Capturing chromosome conformation. *Science* 295(5558): 1306-1311.
- Delabesse E, Ogilvy S, Chapman MA, Piltz SG, Gottgens B et al. (2005) Transcriptional regulation of the SCL locus: identification of an enhancer that targets the primitive erythroid lineage in vivo. *Mol Cell Biol* 25(12): 5215-5225.
- Denell RE, Frederick RD (1983) Homoeosis in *Drosophila*: a description of the Polycomb lethal syndrome. *Dev Biol* 97(1): 34-47.
- Dermitzakis ET, Clark AG (2002) Evolution of transcription factor binding sites in Mammalian gene regulatory regions: conservation and turnover. *Mol Biol Evol* 19(7): 1114-1121.
- Dhalluin C, Carlson JE, Zeng L, He C, Aggarwal AK et al. (1999) Structure and ligand of a histone acetyltransferase bromodomain. *Nature* 399(6735): 491-496.
- Dhami P (2005) The SCL gene and transcriptional control of haematopoiesis. PhD Thesis, University of Cambridge, UK.
- Dhami P, Coffey AJ, Abbs S, Vermeesch JR, Dumanski JP et al. (2005) Exon array CGH: detection of copy-number changes at the resolution of individual exons in the human genome. *Am J Hum Genet* 76(5): 750-762.
- Dhami P, A.W. Bruce, et al. (2008) Re-defining the regulatory landscape at the human SCL (TAL-1) locus. Submitted.
- Di Duca M, Oleggini R, Sanna-Cherchi S, Pasquali L, Di Donato A et al. (2006) Cis and trans regulatory elements in NPHS2 promoter: implications in proteinuria and progression of renal diseases. *Kidney Int* 70(7): 1332-1341.
- Dion MF, Altschuler SJ, Wu LF, Rando OJ (2005) Genomic characterization reveals a simple histone H4 acetylation code. *Proc Natl Acad Sci U S A* 102(15): 5501-5506.
- Donohoe ME, Zhang LF, Xu N, Shi Y, Lee JT (2007) Identification of a Ctfc cofactor, Yy1, for the X chromosome binary switch. *Mol Cell* 25(1): 43-56.
- Dostie J, Richmond TA, Arnaout RA, Selzer RR, Lee WL et al. (2006) Chromosome Conformation Capture Carbon Copy (5C): a massively parallel solution for mapping interactions between genomic elements. *Genome Res* 16(10): 1299-1309.

- Dou Y, Milne TA, Tackett AJ, Smith ER, Fukuda A et al. (2005) Physical association and coordinate function of the H3 K4 methyltransferase MLL1 and the H4 K16 acetyltransferase MOF. *Cell* 121(6): 873-885.
- Down TA, Hubbard TJ (2002) Computational detection and location of transcription start sites in mammalian genomic DNA. *Genome Res* 12(3): 458-461.
- Down TA, Hubbard TJ (2005) NestedMICA: sensitive inference of over-represented motifs in nucleic acid sequence. *Nucleic Acids Res* 33(5): 1445-1453.
- Draper JS, Pigott C, Thomson JA, Andrews PW (2002) Surface antigens of human embryonic stem cells: changes upon differentiation in culture. *J Anat* 200(Pt 3): 249-258.
- Duggan DJ, Bittner M, Chen Y, Meltzer P, Trent JM (1999) Expression profiling using cDNA microarrays. *Nat Genet* 21(1 Suppl): 10-14.
- Dunn KL, Zhao H, Davie JR (2003) The insulator binding protein CTCF associates with the nuclear matrix. *Exp Cell Res* 288(1): 218-223.
- Eichler EE, Sankoff D (2003) Structural dynamics of eukaryotic chromosome evolution. *Science* 301(5634): 793-797.
- Eisen MB, Spellman PT, Brown PO, Botstein D (1998) Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci U S A* 95(25): 14863-14868.
- Elnitski L, Giardine B, Shah P, Zhang Y, Riemer C et al. (2005) Improvements to GALA and dbERGE II: databases featuring genomic sequence alignment, annotation and experimental results. *Nucleic Acids Res* 33(Database issue): D466-470.
- Emanuelsson O, Nagalakshmi U, Zheng D, Rozowsky JS, Urban AE et al. (2007) Assessing the performance of different high-density tiling microarray strategies for mapping transcribed regions of the human genome. *Genome Res* 17(6): 886-897.
- Engel N, West AG, Felsenfeld G, Bartolomei MS (2004) Antagonism between DNA hypermethylation and enhancer-blocking activity at the H19 DMD is uncovered by CpG mutations. *Nat Genet* 36(8): 883-888.
- Enver T, Soneji S, Joshi C, Brown J, Iborra F et al. (2005) Cellular differentiation hierarchies in normal and culture-adapted human embryonic stem cells. *Hum Mol Genet* 14(21): 3129-3140.
- Euskirchen G, Royce TE, Bertone P, Martone R, Rinn JL et al. (2004) CREB binds to multiple loci on human chromosome 22. *Mol Cell Biol* 24(9): 3804-3814.
- Evans MJ, Kaufman MH (1981) Establishment in culture of pluripotential cells from mouse embryos. *Nature* 292(5819): 154-156.
- Farrell CM, West AG, Felsenfeld G (2002) Conserved CTCF insulator elements flank the mouse and human beta-globin loci. *Mol Cell Biol* 22(11): 3820-3831.
- Feinberg AP, Ohlsson R, Henikoff S (2006) The epigenetic progenitor origin of human cancer. *Nat Rev Genet* 7(1): 21-33.

- Feng Q, Wang H, Ng HH, Erdjument-Bromage H, Tempst P et al. (2002) Methylation of H3-lysine 79 is mediated by a new family of HMTases without a SET domain. *Curr Biol* 12(12): 1052-1058.
- Fickett JW, Hatzigeorgiou AG (1997) Eukaryotic promoter recognition. *Genome Res* 7(9): 861-878.
- Fiegler H, Carr P, Douglas EJ, Burford DC, Hunt S et al. (2003) DNA microarrays for comparative genomic hybridization based on DOP-PCR amplification of BAC and PAC clones. *Genes Chromosomes Cancer* 36(4): 361-374.
- Filippova GN, Fagerlie S, Klenova EM, Myers C, Dehner Y et al. (1996) An exceptionally conserved transcriptional repressor, CTCF, employs different combinations of zinc fingers to bind diverged promoter sequences of avian and mammalian c-myc oncogenes. *Mol Cell Biol* 16(6): 2802-2813.
- Filippova GN, Thienes CP, Penn BH, Cho DH, Hu YJ et al. (2001) CTCF-binding sites flank CTG/CAG repeats and form a methylation-sensitive insulator at the DM1 locus. *Nat Genet* 28(4): 335-343.
- Filippova GN, Cheng MK, Moore JM, Truong JP, Hu YJ et al. (2005) Boundaries between chromosomal domains of X inactivation and escape bind CTCF and lack CpG methylation during early development. *Dev Cell* 8(1): 31-42.
- Filippova GN (2008) Genetics and epigenetics of the multifunctional protein CTCF. *Curr Top Dev Biol* 80: 337-360.
- Fischle W, Wang Y, Allis CD (2003) Binary switches and modification cassettes in histone biology and beyond. *Nature* 425(6957): 475-479.
- Fischle W, Tseng BS, Dormann HL, Ueberheide BM, Garcia BA et al. (2005) Regulation of HP1-chromatin binding by histone H3 methylation and phosphorylation. *Nature* 438(7071): 1116-1122.
- Flanagan JF, Mi LZ, Chruszcz M, Cymborowski M, Clines KL et al. (2005) Double chromodomains cooperate to recognize the methylated histone H3 tail. *Nature* 438(7071): 1181-1185.
- Flicek P, Keibler E, Hu P, Korf I, Brent MR (2003) Leveraging the mouse genome for gene prediction in human: from whole-genome shotgun reads to a global synteny map. *Genome Res* 13(1): 46-54.
- Fogel GB, Weekes DG, Varga G, Dow ER, Craven AM et al. (2005) A statistical analysis of the TRANSFAC database. *Biosystems* 81(2): 137-154.
- Follows GA, Tagoh H, Lefevre P, Morgan GJ, Bonifer C (2003) Differential transcription factor occupancy but evolutionarily conserved chromatin features at the human and mouse M-CSF (CSF-1) receptor loci. *Nucleic Acids Res* 31(20): 5805-5816.
- Follows GA, Dhami P, Gottgens B, Bruce AW, Campbell PJ et al. (2006) Identifying gene regulatory elements by genomic microarray mapping of DNaseI hypersensitive sites. *Genome Res* 16(10): 1310-1319.
- Forrester WC, Fernandez LA, Grosschedl R (1999) Nuclear matrix attachment regions antagonize methylation-dependent repression of long-range enhancer-promoter interactions. *Genes Dev* 13(22): 3003-3014.
- Francis NJ, Kingston RE, Woodcock CL (2004) Chromatin compaction by a polycomb group protein complex. *Science* 306(5701): 1574-1577.

- Friedman AD (2007) Transcriptional control of granulocyte and monocyte development. *Oncogene* 26(47): 6816-6828.
- Frith MC, Pheasant M, Mattick JS (2005) The amazing complexity of the human transcriptome. *Eur J Hum Genet* 13(8): 894-897.
- Galas DJ, Schmitz A (1978) DNase footprinting: a simple method for the detection of protein-DNA binding specificity. *Nucleic Acids Res* 5(9): 3157-3170.
- Garcia BA, Shabanowitz J, Hunt DF (2007) Characterization of histones and their post-translational modifications by mass spectrometry. *Curr Opin Chem Biol* 11(1): 66-73.
- Gardiner-Garden M, Frommer M (1987) CpG islands in vertebrate genomes. *J Mol Biol* 196(2): 261-282.
- Garner MM, Revzin A (1981) A gel electrophoresis method for quantifying the binding of proteins to specific DNA regions: application to components of the Escherichia coli lactose operon regulatory system. *Nucleic Acids Res* 9(13): 3047-3060.
- Gaszner M, Felsenfeld G (2006) Insulators: exploiting transcriptional and epigenetic mechanisms. *Nat Rev Genet* 7(9): 703-713.
- Gautier T, Abbott DW, Molla A, Verdel A, Ausio J et al. (2004) Histone variant H2ABbd confers lower stability to the nucleosome. *EMBO Rep* 5(7): 715-720.
- Gavin I, Horn PJ, Peterson CL (2001) SWI/SNF chromatin remodeling requires changes in DNA topology. *Mol Cell* 7(1): 97-104.
- Geiger JH, Hahn S, Lee S, Sigler PB (1996) Crystal structure of the yeast TFIIA/TBP/DNA complex. *Science* 272(5263): 830-836.
- Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M et al. (2004) Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol* 5(10): R80.
- Gershenson NI, Ioshikhes IP (2005) Synergy of human Pol II core promoter elements revealed by statistical sequence analysis. *Bioinformatics* 21(8): 1295-1300.
- Gerstein MB, Bruce C, Rozowsky JS, Zheng D, Du J et al. (2007) What is a gene, post-ENCODE? History and updated definition. *Genome Res* 17(6): 669-681.
- Geyer PK, Spana C, Corces VG (1986) On the molecular mechanism of gypsy-induced mutations at the yellow locus of *Drosophila melanogaster*. *EMBO J* 5(10): 2657-2662.
- Ghioni P, D'Alessandra Y, Mansueto G, Jaffray E, Hay RT et al. (2005) The protein stability and transcriptional activity of p63alpha are regulated by SUMO-1 conjugation. *Cell Cycle* 4(1): 183-190.
- Gibbons FD, Proft M, Struhl K, Roth FP (2005) Chipper: discovering transcription-factor targets from chromatin immunoprecipitation microarrays using variance stabilization. *Genome Biol* 6(11): R96.
- Gilbert N, Boyle S, Fiegler H, Woodfine K, Carter NP et al. (2004) Chromatin architecture of the human genome: gene-rich domains are enriched in open chromatin fibers. *Cell* 118(5): 555-566.

- Gilbert DM (2007) Replication origin plasticity, Taylor-made: inhibition vs recruitment of origins under conditions of replication stress. *Chromosoma* 116(4): 341-347.
- Giresi PG, Kim J, McDaniel RM, Iyer VR, Lieb JD (2007) FAIRE (Formaldehyde-Assisted Isolation of Regulatory Elements) isolates active regulatory elements from human chromatin. *Genome Res* 17(6): 877-885.
- Golderer G, Grobner P (1991) ADP-ribosylation of core histones and their acetylated subspecies. *Biochem J* 277 (Pt 3): 607-610.
- Goldknopf IL, Taylor CW, Baum RM, Yeoman LC, Olson MO et al. (1975) Isolation and characterization of protein A24, a "histone-like" non-histone chromosomal protein. *J Biol Chem* 250(18): 7182-7187.
- Goodstadt L, Ponting CP (2006) Phylogenetic reconstruction of orthology, paralogy, and conserved synteny for dog and human. *PLoS Comput Biol* 2(9): e133.
- Gottgens B, McLaughlin F, Bockamp EO, Fordham JL, Begley CG et al. (1997) Transcription of the SCL gene in erythroid and CD34 positive primitive myeloid cells is controlled by a complex network of lineage-restricted chromatin-dependent and chromatin-independent regulatory elements. *Oncogene* 15(20): 2419-2428.
- Gottgens B, Barton LM, Gilbert JG, Bench AJ, Sanchez MJ et al. (2000) Analysis of vertebrate SCL loci identifies conserved enhancers. *Nat Biotechnol* 18(2): 181-186.
- Gottgens B, Gilbert JG, Barton LM, Grafham D, Rogers J et al. (2001) Long-range comparison of human and mouse SCL loci: localized regions of sensitivity to restriction endonucleases correspond precisely with peaks of conserved noncoding sequences. *Genome Res* 11(1): 87-97.
- Gottgens B, Nastos A, Kinston S, Piltz S, Delabesse EC et al. (2002) Establishing the transcriptional programme for blood: the SCL stem cell enhancer is regulated by a multiprotein complex containing Ets and GATA factors. *EMBO J* 21(12): 3039-3050.
- Greenbaum JA, Parker SC, Tullius TD (2007) Detection of DNA structural motifs in functional genomic elements. *Genome Res* 17(6): 940-946.
- Grewal SI, Moazed D (2003) Heterochromatin and epigenetic control of gene expression. *Science* 301(5634): 798-802.
- Grewal SI, Jia S (2007) Heterochromatin revisited. *Nat Rev Genet* 8(1): 35-46.
- Gross DS, Garrard WT (1988) Nuclease hypersensitive sites in chromatin. *Annu Rev Biochem* 57: 159-197.
- Grosveld F, van Assendelft GB, Greaves DR, Kollias G (1987) Position-independent, high-level expression of the human beta-globin gene in transgenic mice. *Cell* 51(6): 975-985.
- Gruzdeva N, Kyrchanova O, Parshikov A, Kullyev A, Georgiev P (2005) The Mcp element from the bithorax complex contains an insulator that is capable of pairwise interactions and can facilitate enhancer-promoter communication. *Mol Cell Biol* 25(9): 3682-3689.

- Guenther MG, Levine SS, Boyer LA, Jaenisch R, Young RA (2007) A chromatin landmark and transcription initiation at most promoters in human cells. *Cell* 130(1): 77-88.
- Hallikas O, Palin K, Sinjushina N, Rautiainen R, Partanen J et al. (2006) Genome-wide prediction of mammalian enhancers based on analysis of transcription-factor binding affinity. *Cell* 124(1): 47-59.
- Hampsey M (1998) Molecular genetics of the RNA polymerase II general transcriptional machinery. *Microbiol Mol Biol Rev* 62(2): 465-503.
- Hannich JT, Lewis A, Kroetz MB, Li SJ, Heide H et al. (2005) Defining the SUMO-modified proteome by multiple approaches in *Saccharomyces cerevisiae*. *J Biol Chem* 280(6): 4102-4110.
- Hark AT, Schoenherr CJ, Katz DJ, Ingram RS, Levorse JM et al. (2000) CTCF mediates methylation-sensitive enhancer-blocking activity at the H19/Igf2 locus. *Nature* 405(6785): 486-489.
- Harris MB, Mostecky J, Rothman PB (2005) Repression of an interleukin-4-responsive promoter requires cooperative BCL-6 function. *J Biol Chem* 280(13): 13114-13121.
- Harrow J, Denoeud F, Frankish A, Reymond A, Chen CK et al. (2006) GENCODE: producing a reference annotation for ENCODE. *Genome Biol* 7 Suppl 1: S4 1-9.
- Hassa PO, Haenni SS, Elser M, Hottiger MO (2006) Nuclear ADP-ribosylation reactions in mammalian cells: where are we today and where are we going? *Microbiol Mol Biol Rev* 70(3): 789-829.
- Hassan AH, Prochasson P, Neely KE, Galasinski SC, Chandy M et al. (2002) Function and selectivity of bromodomains in anchoring chromatin-modifying complexes to promoter nucleosomes. *Cell* 111(3): 369-379.
- Heintzman ND, Stuart RK, Hon G, Fu Y, Ching CW et al. (2007) Distinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome. *Nat Genet* 39(3): 311-318.
- Heinzel T, Lavinsky RM, Mullen TM, Soderstrom M, Laherty CD et al. (1997) A complex containing N-CoR, mSin3 and histone deacetylase mediates transcriptional repression. *Nature* 387(6628): 43-48.
- Henderson JK, Draper JS, Baillie HS, Fishel S, Thomson JA et al. (2002) Preimplantation human embryos and embryonic stem cells show comparable expression of stage-specific embryonic antigens. *Stem Cells* 20(4): 329-337.
- Hendrich B, Bird A (1998) Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Mol Cell Biol* 18(11): 6538-6547.
- Heng HH, Goetze S, Ye CJ, Liu G, Stevens JB et al. (2004) Chromatin loops are selectively anchored using scaffold/matrix-attachment regions. *J Cell Sci* 117(Pt 7): 999-1008.
- Higgs DR, Wood WG, Jarman AP, Sharpe J, Lida J et al. (1990) A major positive regulatory region located far upstream of the human alpha-globin gene locus. *Genes Dev* 4(9): 1588-1601.
- Hoheisel JD (2006) Microarray technology: beyond transcript profiling and genotype analysis. *Nat Rev Genet* 7(3): 200-210.

- Holohan EE, Kwong C, Adryan B, Bartkuhn M, Herold M et al. (2007) CTCF Genomic Binding Sites in *Drosophila* and the Organisation of the Bithorax Complex. *PLoS Genet* 3(7): e112.
- Honda BM, Dixon GH, Candido EP (1975) Sites of in vivo histone methylation in developing trout testis. *J Biol Chem* 250(22): 8681-8685.
- Hong S, Cho YW, Yu LR, Yu H, Veenstra TD et al. (2007) Identification of JmjC domain-containing UTX and JMJD3 as histone H3 lysine 27 demethylases. *Proc Natl Acad Sci U S A* 104(47): 18439-18444.
- Horak CE, Mahajan MC, Luscombe NM, Gerstein M, Weissman SM et al. (2002) GATA-1 binding sites mapped in the beta-globin locus by using mammalian ChIP-chip analysis. *Proc Natl Acad Sci U S A* 99(5): 2924-2929.
- Huang Y, Myers SJ, Dingledine R (1999) Transcriptional repression by REST: recruitment of Sin3A and histone deacetylase to neuronal genes. *Nat Neurosci* 2(10): 867-872.
- Huang S, Litt M, Felsenfeld G (2005) Methylation of histone H4 by arginine methyltransferase PRMT1 is essential in vivo for many subsequent histone modifications. *Genes Dev* 19(16): 1885-1893.
- Huang Y, Fang J, Bedford MT, Zhang Y, Xu RM (2006) Recognition of histone H3 lysine-4 methylation by the double tudor domain of JMJD2A. *Science* 312(5774): 748-751.
- Huang S, Li X, Yusufzai TM, Qiu Y, Felsenfeld G (2007) USF1 recruits histone modification complexes and is critical for maintenance of a chromatin barrier. *Mol Cell Biol* 27(22): 7991-8002.
- Hughes JD, Estep PW, Tavazoie S, Church GM (2000) Computational identification of cis-regulatory elements associated with groups of functionally related genes in *Saccharomyces cerevisiae*. *J Mol Biol* 296(5): 1205-1214.
- Hughes TR, Mao M, Jones AR, Burchard J, Marton MJ et al. (2001) Expression profiling using microarrays fabricated by an ink-jet oligonucleotide synthesizer. *Nat Biotechnol* 19(4): 342-347.
- Im H, Park C, Feng Q, Johnson KD, Kiekhäfer CM et al. (2003) Dynamic regulation of histone H3 methylated at lysine 79 within a tissue-specific chromatin domain. *J Biol Chem* 278(20): 18346-18352.
- Impey S, McCorkle SR, Cha-Molstad H, Dwyer JM, Yochum GS et al. (2004) Defining the CREB regulon: a genome-wide analysis of transcription factor regulatory regions. *Cell* 119(7): 1041-1054.
- Ioudinkova E, Petrov A, Razin SV, Vassetzky YS (2005) Mapping long-range chromatin organization within the chicken alpha-globin gene domain using oligonucleotide DNA arrays. *Genomics* 85(1): 143-151.
- Ishihara K, Oshimura M, Nakao M (2006) CTCF-dependent chromatin insulator is linked to epigenetic remodeling. *Mol Cell* 23(5): 733-742.
- Ishii K, Arib G, Lin C, Van Houwe G, Laemmli UK (2002) Chromatin boundaries in budding yeast: the nuclear pore connection. *Cell* 109(5): 551-562.
- Ivanova NB, Dimos JT, Schaniel C, Hackney JA, Moore KA et al. (2002) A stem cell molecular signature. *Science* 298(5593): 601-604.

- Iyer VR, Horak CE, Scafe CS, Botstein D, Snyder M et al. (2001) Genomic binding sites of the yeast cell-cycle transcription factors SBF and MBF. *Nature* 409(6819): 533-538.
- Jarman AP, Wood WG, Sharpe JA, Gourdon G, Ayyub H et al. (1991) Characterization of the major regulatory element upstream of the human alpha-globin gene cluster. *Mol Cell Biol* 11(9): 4679-4689.
- Jeffreys AJ, Kauppi L, Neumann R (2001) Intensely punctate meiotic recombination in the class II region of the major histocompatibility complex. *Nat Genet* 29(2): 217-222.
- Jenke AC, Stehle IM, Herrmann F, Eisenberger T, Baiker A et al. (2004) Nuclear scaffold/matrix attached region modules linked to a transcription unit are sufficient for replication and maintenance of a mammalian episome. *Proc Natl Acad Sci U S A* 101(31): 11322-11327.
- Jenuwein T, Forrester WC, Fernandez-Herrero LA, Laible G, Dull M et al. (1997) Extension of chromatin accessibility by nuclear matrix attachment regions. *Nature* 385(6613): 269-272.
- Jenuwein T, Allis CD (2001) Translating the histone code. *Science* 293(5532): 1074-1080.
- Jeon Y, Bekiranov S, Karnani N, Kapranov P, Ghosh S et al. (2005) Temporal profile of replication of human chromosomes. *Proc Natl Acad Sci U S A* 102(18): 6419-6424.
- Johansen KM, Johansen J (2006) Regulation of chromatin structure by histone H3S10 phosphorylation. *Chromosome Res* 14(4): 393-404.
- Johnson KD, Christensen HM, Zhao B, Bresnick EH (2001) Distinct mechanisms control RNA polymerase II recruitment to a tissue-specific locus control region and a downstream promoter. *Mol Cell* 8(2): 465-471.
- Johnson KD, Grass JA, Boyer ME, Kiekhäfer CM, Blobel GA et al. (2002) Cooperative activities of hematopoietic regulators recruit RNA polymerase II to a tissue-specific chromatin domain. *Proc Natl Acad Sci U S A* 99(18): 11760-11765.
- Johnson DS, Mortazavi A, Myers RM, Wold B (2007) Genome-wide mapping of in vivo protein-DNA interactions. *Science* 316(5830): 1497-1502.
- Jones N (1990) Transcriptional regulation by dimerization: two sides to an incestuous relationship. *Cell* 61(1): 9-11.
- Jones PL, Veenstra GJ, Wade PA, Vermaak D, Kass SU et al. (1998) Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat Genet* 19(2): 187-191.
- Jones PA, Baylin SB (2007) The epigenomics of cancer. *Cell* 128(4): 683-692.
- Kadonaga JT (2002) The DPE, a core promoter element for transcription by RNA polymerase II. *Exp Mol Med* 34(4): 259-264.
- Kadota M, Yang HH, Hu N, Wang C, Hu Y et al. (2007) Allele-specific chromatin immunoprecipitation studies show genetic influence on chromatin state in human genome. *PLoS Genet* 3(5): e81.
- Kamakaka RT, Biggins S (2005) Histone variants: deviants? *Genes Dev* 19(3): 295-310.

- Kanduri C, Pant V, Loukinov D, Pugacheva E, Qi CF et al. (2000) Functional association of CTCF with the insulator upstream of the H19 gene is parent of origin-specific and methylation-sensitive. *Curr Biol* 10(14): 853-856.
- Kanduri M, Kanduri C, Mariano P, Vostrov AA, Quitschke W et al. (2002) Multiple nucleosome positioning sites regulate the CTCF-mediated insulator function of the H19 imprinting control region. *Mol Cell Biol* 22(10): 3339-3344.
- Kapranov P, Cheng J, Dike S, Nix DA, Duttagupta R et al. (2007) RNA maps reveal new RNA classes and a possible function for pervasive transcription. *Science* 316(5830): 1484-1488.
- Karnani N, Taylor C, Malhotra A, Dutta A (2007) Pan-S replication patterns and chromosomal domains defined by genome-tiling arrays of ENCODE genomic areas. *Genome Res* 17(6): 865-876.
- Katayama S, Tomaru Y, Kasukawa T, Waki K, Nakanishi M et al. (2005) Antisense transcription in the mammalian transcriptome. *Science* 309(5740): 1564-1566.
- Kel AE, Gossling E, Reuter I, Cheremushkin E, Kel-Margoulis OV et al. (2003) MATCH: A tool for searching transcription factor binding sites in DNA sequences. *Nucleic Acids Res* 31(13): 3576-3579.
- Keogh MC, Kurdistani SK, Morris SA, Ahn SH, Podolny V et al. (2005) Cotranscriptional set2 methylation of histone H3 lysine 36 recruits a repressive Rpd3 complex. *Cell* 123(4): 593-605.
- Khorasanizadeh S (2004) The nucleosome: from genomic organization to genomic regulation. *Cell* 116(2): 259-272.
- Kim M, Park CH, Lee MS, Carlson BA, Hatfield DL et al. (2003) A novel TBP-interacting zinc finger protein represses transcription by inhibiting the recruitment of TFIIA and TFIIB. *Biochem Biophys Res Commun* 306(1): 231-238.
- Kim A, Dean A (2004) Developmental stage differences in chromatin subdomains of the beta-globin locus. *Proc Natl Acad Sci U S A* 101(18): 7028-7033.
- Kim TH, Barrera LO, Zheng M, Qu C, Singer MA et al. (2005) A high-resolution map of active promoters in the human genome. *Nature* 436(7052): 876-880.
- Kim TH, Abdullaev ZK, Smith AD, Ching KA, Loukinov DI et al. (2007) Analysis of the vertebrate insulator protein CTCF-binding sites in the human genome. *Cell* 128(6): 1231-1245.
- King DC, Taylor J, Zhang Y, Cheng Y, Lawson HA et al. (2007) Finding cis-regulatory elements using comparative genomics: some lessons from ENCODE data. *Genome Res* 17(6): 775-786.
- Kirmizis A, Bartley SM, Kuzmichev A, Margueron R, Reinberg D et al. (2004) Silencing of human polycomb target genes is associated with methylation of histone H3 Lys 27. *Genes Dev* 18(13): 1592-1605.
- Kirmizis A, Santos-Rosa H, Penkett CJ, Singer MA, Vermeulen M et al. (2007) Arginine methylation at histone H3R2 controls deposition of H3K4 trimethylation. *Nature* 449(7164): 928-932.
- Kleff S, Andrulis ED, Anderson CW, Sternglanz R (1995) Identification of a gene encoding a yeast histone H4 acetyltransferase. *J Biol Chem* 270(42): 24674-24677.

- Klenova EM, Nicolas RH, Paterson HF, Carne AF, Heath CM et al. (1993) CTCF, a conserved nuclear factor required for optimal transcriptional activity of the chicken c-myc gene, is an 11-Zn-finger protein differentially expressed in multiple forms. *Mol Cell Biol* 13(12): 7612-7624.
- Klenova EM, Nicolas RH, U S, Carne AF, Lee RE et al. (1997) Molecular weight abnormalities of the CTCF transcription factor: CTCF migrates aberrantly in SDS-PAGE and the size of the expressed protein is affected by the UTRs and sequences within the coding region of the CTCF gene. *Nucleic Acids Res* 25(3): 466-474.
- Klochkov D, Rincon-Arango H, Ioudinkova ES, Valadez-Graham V, Gavrilov A et al. (2006) A CTCF-dependent silencer located in the differentially methylated area may regulate expression of a housekeeping gene overlapping a tissue-specific gene domain. *Mol Cell Biol* 26(5): 1589-1597.
- Klose RJ, Bird AP (2006) Genomic DNA methylation: the mark and its mediators. *Trends Biochem Sci* 31(2): 89-97.
- Klose RJ, Gardner KE, Liang G, Erdjument-Bromage H, Tempst P et al. (2007) Demethylation of histone H3K36 and H3K9 by Rph1: a vestige of an H3K9 methylation system in *Saccharomyces cerevisiae*? *Mol Cell Biol* 27(11): 3951-3961.
- Klug J (1997) Ku autoantigen is a potential major cause of nonspecific bands in electrophoretic mobility shift assays. *Biotechniques* 22(2): 212-214, 216.
- Koch CM, Andrews RM, Flicek P, Dillon SC, Karaoz U et al. (2007) The landscape of histone modifications across 1% of the human genome in five human cell lines. *Genome Res* 17(6): 691-707.
- Komarnitsky P, Cho EJ, Buratowski S (2000) Different phosphorylated forms of RNA polymerase II and associated mRNA processing factors during transcription. *Genes Dev* 14(19): 2452-2460.
- Kondrashov AS, Shabalina SA (2002) Classification of common conserved sequences in mammalian intergenic regions. *Hum Mol Genet* 11(6): 669-674.
- Kornberg RD (1974) Chromatin structure: a repeating unit of histones and DNA. *Science* 184(139): 868-871.
- Kornberg RD, Thomas JO (1974) Chromatin structure; oligomers of the histones. *Science* 184(139): 865-868.
- Kornberg RD, Lorch Y (1999) Twenty-five years of the nucleosome, fundamental particle of the eukaryote chromosome. *Cell* 98(3): 285-294.
- Kouzarides T (2007) Chromatin modifications and their function. *Cell* 128(4): 693-705.
- Krogan NJ, Kim M, Tong A, Golshani A, Cagney G et al. (2003) Methylation of histone H3 by Set2 in *Saccharomyces cerevisiae* is linked to transcriptional elongation by RNA polymerase II. *Mol Cell Biol* 23(12): 4207-4218.
- Kuhn RM, Karolchik D, Zweig AS, Trumbower H, Thomas DJ et al. (2007) The UCSC genome browser database: update 2007. *Nucleic Acids Res* 35(Database issue): D668-673.

- Kumar PP, Bischof O, Purbey PK, Notani D, Urlaub H et al. (2007) Functional interaction between PML and SATB1 regulates chromatin-loop architecture and transcription of the MHC class I locus. *Nat Cell Biol* 9(1): 45-56.
- Kummerfeld SK, Teichmann SA (2006) DBD: a transcription factor prediction database. *Nucleic Acids Res* 34(Database issue): D74-81.
- Kurdistani SK, Robyr D, Tavazoie S, Grunstein M (2002) Genome-wide binding map of the histone deacetylase Rpd3 in yeast. *Nat Genet* 31(3): 248-254.
- Kurdistani SK, Grunstein M (2003) Histone acetylation and deacetylation in yeast. *Nat Rev Mol Cell Biol* 4(4): 276-284.
- Kurdistani SK, Tavazoie S, Grunstein M (2004) Mapping global histone acetylation patterns to gene expression. *Cell* 117(6): 721-733.
- Kurukuti S, Tiwari VK, Tavosidana G, Pugacheva E, Murrell A et al. (2006) CTCF binding at the H19 imprinting control region mediates maternally inherited higher-order chromatin conformation to restrict enhancer access to Igf2. *Proc Natl Acad Sci U S A* 103(28): 10684-10689.
- Kuzmichev A, Nishioka K, Erdjument-Bromage H, Tempst P, Reinberg D (2002) Histone methyltransferase activity associated with a human multiprotein complex containing the Enhancer of Zeste protein. *Genes Dev* 16(22): 2893-2905.
- Kuzmichev A, Margueron R, Vaquero A, Preissner TS, Scher M et al. (2005) Composition and histone substrates of polycomb repressive group complexes change during cellular differentiation. *Proc Natl Acad Sci U S A* 102(6): 1859-1864.
- Lachner M, O'Carroll D, Rea S, Mechtler K, Jenuwein T (2001) Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature* 410(6824): 116-120.
- Lachner M, O'Sullivan RJ, Jenuwein T (2003) An epigenetic road map for histone lysine methylation. *J Cell Sci* 116(Pt 11): 2117-2124.
- Lan F, Bayliss PE, Rinn JL, Whetstone JR, Wang JK et al. (2007) A histone H3 lysine 27 demethylase regulates animal posterior development. *Nature* 449(7163): 689-694.
- Lan F, Zaratiegui M, Villen J, Vaughn MW, Verdel A et al. (2007 b) *S. pombe* LSD1 homologs regulate heterochromatin propagation and euchromatic gene transcription. *Mol Cell* 26(1): 89-101.
- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC et al. (2001) Initial sequencing and analysis of the human genome. *Nature* 409(6822): 860-921.
- Lang G, Mamalaki C, Greenberg D, Yannoutsos N, Kioussis D (1991) Deletion analysis of the human CD2 gene locus control region in transgenic mice. *Nucleic Acids Res* 19(21): 5851-5856.
- Lee TI, Young RA (2000) Transcription of eukaryotic protein-coding genes. *Annu Rev Genet* 34: 77-137.
- Lee TI, Rinaldi NJ, Robert F, Odom DT, Bar-Joseph Z et al. (2002) Transcriptional regulatory networks in *Saccharomyces cerevisiae*. *Science* 298(5594): 799-804.

- Lee YH, Koh SS, Zhang X, Cheng X, Stallcup MR (2002 b) Synergy among nuclear receptor coactivators: selective requirement for protein methyltransferase and acetyltransferase activities. *Mol Cell Biol* 22(11): 3621-3632.
- Lee CK, Shibata Y, Rao B, Strahl BD, Lieb JD (2004) Evidence for nucleosome depletion at active regulatory regions genome-wide. *Nat Genet* 36(8): 900-905.
- Lee DY, Teyssier C, Strahl BD, Stallcup MR (2005) Role of protein methylation in regulation of transcription. *Endocr Rev* 26(2): 147-170.
- Lee MG, Wynder C, Cooch N, Shiekhhattar R (2005) An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature* 437(7057): 432-435.
- Lee JH, Skalnik DG (2005) CpG-binding protein (CXXC finger protein 1) is a component of the mammalian Set1 histone H3-Lys4 methyltransferase complex, the analogue of the yeast Set1/COMPASS complex. *J Biol Chem* 280(50): 41725-41731.
- Lee TI, Jenner RG, Boyer LA, Guenther MG, Levine SS et al. (2006) Control of developmental regulators by Polycomb in human embryonic stem cells. *Cell* 125(2): 301-313.
- Lee N, Zhang J, Klose RJ, Erdjument-Bromage H, Tempst P et al. (2007) The trithorax-group protein Lid is a histone H3 trimethyl-Lys4 demethylase. *Nat Struct Mol Biol* 14(4): 341-343.
- Lee JS, Shilatifard A (2007) A site to remember: H3K36 methylation a mark for histone deacetylation. *Mutat Res* 618(1-2): 130-134.
- Leroy-Viard K, Vinit MA, Lecoite N, Mathieu-Mahul D, Romeo PH (1994) Distinct DNase-I hypersensitive sites are associated with TAL-1 transcription in erythroid and T-cell lines. *Blood* 84(11): 3819-3827.
- Lestrade L, Weber MJ (2006) snoRNA-LBME-db, a comprehensive database of human H/ACA and C/D box snoRNAs. *Nucleic Acids Res* 34(Database issue): D158-162.
- Levine SS, King IF, Kingston RE (2004) Division of labor in polycomb group repression. *Trends Biochem Sci* 29(9): 478-485.
- Lewis EB (1978) A gene complex controlling segmentation in *Drosophila*. *Nature* 276(5688): 565-570.
- Li Q, Peterson KR, Fang X, Stamatoyannopoulos G (2002) Locus control regions. *Blood* 100(9): 3077-3086.
- Li Z, Van Calcar S, Qu C, Cavenee WK, Zhang MQ et al. (2003) A global transcriptional regulatory role for c-Myc in Burkitt's lymphoma cells. *Proc Natl Acad Sci U S A* 100(14): 8164-8169.
- Li L, He S, Sun JM, Davie JR (2004) Gene regulation by Sp1 and Sp3. *Biochem Cell Biol* 82(4): 460-471.
- Li W, Meyer CA, Liu XS (2005) A hidden Markov model for analyzing ChIP-chip experiments on genome tiling arrays and its application to p53 binding sequences. *Bioinformatics* 21 Suppl 1: i274-282.
- Li H, Ilin S, Wang W, Duncan EM, Wysocka J et al. (2006) Molecular basis for site-specific read-out of histone H3K4me3 by the BPTF PHD finger of NURF. *Nature* 442(7098): 91-95.
- Li B, Carey M, Workman JL (2007) The role of chromatin during transcription. *Cell* 128(4): 707-719.

- Li H, Fischle W, Wang W, Duncan EM, Liang L et al. (2007 b) Structural basis for lower lysine methylation state-specific readout by MBT repeats of L3MBTL1 and an engineered PHD finger. *Mol Cell* 28(4): 677-691.
- Li B, Gogol M, Carey M, Lee D, Seidel C et al. (2007 c) Combined action of PHD and chromo domains directs the Rpd3S HDAC to transcribed chromatin. *Science* 316(5827): 1050-1054.
- Liang G, Lin JC, Wei V, Yoo C, Cheng JC et al. (2004) Distinct localization of histone H3 acetylation and H3-K4 methylation to the transcription start sites in the human genome. *Proc Natl Acad Sci U S A* 101(19): 7357-7362.
- Liang G, Klose RJ, Gardner KE, Zhang Y (2007) Yeast Jhd2p is a histone H3 Lys4 trimethyl demethylase. *Nat Struct Mol Biol* 14(3): 243-245.
- Lieb M, Rehmat S, Bhagwat AS (2001) Interaction of MutS and Vsr: some dominant-negative mutS mutations that disable methyladenine-directed mismatch repair are active in very-short-patch repair. *J Bacteriol* 183(21): 6487-6490.
- Lieu PT, Jozsi P, Gilles P, Peterson T (2005) Development of a DNA-labeling system for array-based comparative genomic hybridization. *J Biomol Tech* 16(2): 104-111.
- Ling J, Ainol L, Zhang L, Yu X, Pi W et al. (2004) HS2 enhancer function is blocked by a transcriptional terminator inserted between the enhancer and the promoter. *J Biol Chem* 279(49): 51704-51713.
- Ling JQ, Li T, Hu JF, Vu TH, Chen HL et al. (2006) CTCF mediates interchromosomal colocalization between Igf2/H19 and Wsb1/Nf1. *Science* 312(5771): 269-272.
- Lipshutz RJ, Fodor SP, Gingeras TR, Lockhart DJ (1999) High density synthetic oligonucleotide arrays. *Nat Genet* 21(1 Suppl): 20-24.
- Litt MD, Simpson M, Recillas-Targa F, Prioleau MN, Felsenfeld G (2001) Transitions in histone acetylation reveal boundaries of three separately regulated neighboring loci. *EMBO J* 20(9): 2224-2235.
- Liu CL, Schreiber SL, Bernstein BE (2003) Development and validation of a T7 based linear amplification for genomic DNA. *BMC Genomics* 4(1): 19.
- Liu CL, Kaplan T, Kim M, Buratowski S, Schreiber SL et al. (2005) Single-nucleosome mapping of histone modifications in *S. cerevisiae*. *PLoS Biol* 3(10): e328.
- Liu L, Zhang J, Bates S, Li JJ, Peehl DM et al. (2005 b) A methylation profile of in vitro immortalized human cell lines. *Int J Oncol* 26(1): 275-285.
- Lobanenkov VV, Nicolas RH, Adler VV, Paterson H, Klenova EM et al. (1990) A novel sequence-specific DNA binding protein which interacts with three regularly spaced direct repeats of the CCCTC-motif in the 5'-flanking sequence of the chicken c-myc gene. *Oncogene* 5(12): 1743-1753.
- Loh YH, Wu Q, Chew JL, Vega VB, Zhang W et al. (2006) The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nat Genet* 38(4): 431-440.
- Lohr D, Lopez J (1995) GAL4/GAL80-dependent nucleosome disruption/deposition on the upstream regions of the yeast GAL1-10 and GAL80 genes. *J Biol Chem* 270(46): 27671-27678.

- Lorincz MC, Dickerson DR, Schmitt M, Groudine M (2004) Intragenic DNA methylation alters chromatin structure and elongation efficiency in mammalian cells. *Nat Struct Mol Biol* 11(11): 1068-1075.
- Lozzio BB, Lozzio CB (1979) Properties and usefulness of the original K-562 human myelogenous leukemia cell line. *Leuk Res* 3(6): 363-370.
- Luger K, Mader AW, Richmond RK, Sargent DF, Richmond TJ (1997) Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature* 389(6648): 251-260.
- Lunyak VV, Burgess R, Prefontaine GG, Nelson C, Sze SH et al. (2002) Corepressor-dependent silencing of chromosomal regions encoding neuronal genes. *Science* 298(5599): 1747-1752.
- Lusser A, Kadonaga JT (2003) Chromatin remodeling by ATP-dependent molecular machines. *Bioessays* 25(12): 1192-1200.
- Lutz M, Burke LJ, Barreto G, Goeman F, Greb H et al. (2000) Transcriptional repression by the insulator protein CTCF involves histone deacetylases. *Nucleic Acids Res* 28(8): 1707-1713.
- Maeda N, Kasukawa T, Oyama R, Gough J, Frith M et al. (2006) Transcript annotation in FANTOM3: mouse gene catalog based on physical cDNAs. *PLoS Genet* 2(4): e62.
- Mahadevan LC, Willis AC, Barratt MJ (1991) Rapid histone H3 phosphorylation in response to growth factors, phorbol esters, okadaic acid, and protein synthesis inhibitors. *Cell* 65(5): 775-783.
- Mak W, Baxter J, Silva J, Newall AE, Otte AP et al. (2002) Mitotically stable association of polycomb group proteins *eed* and *enx1* with the inactive X chromosome in trophoblast stem cells. *Curr Biol* 12(12): 1016-1020.
- Maniatis T, Goodbourn S, Fischer JA (1987) Regulation of inducible and tissue-specific gene expression. *Science* 236(4806): 1237-1245.
- Manza LL, Codreanu SG, Stamer SL, Smith DL, Wells KS et al. (2004) Global shifts in protein sumoylation in response to electrophile and oxidative stress. *Chem Res Toxicol* 17(12): 1706-1715.
- Mao DY, Watson JD, Yan PS, Barsyte-Lovejoy D, Khosravi F et al. (2003) Analysis of Myc bound loci identified by CpG island arrays shows that Max is essential for Myc-dependent repression. *Curr Biol* 13(10): 882-886.
- Margulies EH, Cooper GM, Asimenos G, Thomas DJ, Dewey CN et al. (2007) Analyses of deep mammalian sequence alignments and constraint predictions for 1% of the human genome. *Genome Res* 17(6): 760-774.
- Marmorstein R (2003) Structure of SET domain proteins: a new twist on histone methylation. *Trends Biochem Sci* 28(2): 59-62.
- Martin GR (1981) Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc Natl Acad Sci U S A* 78(12): 7634-7638.
- Martin N, Patel S, Segre JA (2004) Long-range comparison of human and mouse *Spr* loci to identify conserved noncoding sequences involved in coordinate regulation. *Genome Res* 14(12): 2430-2438.

- Martin D, Brun C, Remy E, Mouren P, Thieffry D et al. (2004) GOToolBox: functional analysis of gene datasets based on Gene Ontology. *Genome Biol* 5(12): R101.
- Martin C, Zhang Y (2005) The diverse functions of histone lysine methylation. *Nat Rev Mol Cell Biol* 6(11): 838-849.
- Martone R, Euskirchen G, Bertone P, Hartman S, Royce TE et al. (2003) Distribution of NF-kappaB-binding sites across human chromosome 22. *Proc Natl Acad Sci U S A* 100(21): 12247-12252.
- Maston GA, Evans SK, Green MR (2006) Transcriptional regulatory elements in the human genome. *Annu Rev Genomics Hum Genet* 7: 29-59.
- Mattick JS, Makunin IV (2006) Non-coding RNA. *Hum Mol Genet* 15 Spec No 1: R17-29.
- Mattick JS (2007) A new paradigm for developmental biology. *J Exp Biol* 210(Pt 9): 1526-1547.
- Matys V, Kel-Margoulis OV, Fricke E, Liebich I, Land S et al. (2006) TRANSFAC and its module TRANSCompel: transcriptional gene regulation in eukaryotes. *Nucleic Acids Res* 34(Database issue): D108-110.
- Mautner J, Joos S, Werner T, Eick D, Bornkamm GW et al. (1995) Identification of two enhancer elements downstream of the human c-myc gene. *Nucleic Acids Res* 23(1): 72-80.
- McNairn AJ, Gilbert DM (2003) Epigenomic replication: linking epigenetics to DNA replication. *Bioessays* 25(7): 647-656.
- Meneghini MD, Wu M, Madhani HD (2003) Conserved histone variant H2A.Z protects euchromatin from the ectopic spread of silent heterochromatin. *Cell* 112(5): 725-736.
- Meshorer E, Yellajoshula D, George E, Scambler PJ, Brown DT et al. (2006) Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic stem cells. *Dev Cell* 10(1): 105-116.
- Metzger E, Wissmann M, Yin N, Muller JM, Schneider R et al. (2005) LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription. *Nature* 437(7057): 436-439.
- Miao F, Natarajan R (2005) Mapping global histone methylation patterns in the coding regions of human genes. *Mol Cell Biol* 25(11): 4650-4661.
- Mikkelsen TS, Ku M, Jaffe DB, Issac B, Lieberman E et al. (2007) Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* 448(7153): 553-560.
- Millar CB, Xu F, Zhang K, Grunstein M (2006) Acetylation of H2AZ Lys 14 is associated with genome-wide gene activity in yeast. *Genes Dev* 20(6): 711-722.
- Min J, Zhang Y, Xu RM (2003) Structural basis for specific binding of Polycomb chromodomain to histone H3 methylated at Lys 27. *Genes Dev* 17(15): 1823-1828.
- Mirkovitch J, Mirault ME, Laemmli UK (1984) Organization of the higher-order chromatin loop: specific DNA attachment sites on nuclear scaffold. *Cell* 39(1): 223-232.
- Mito Y, Henikoff JG, Henikoff S (2005) Genome-scale profiling of histone H3.3 replacement patterns. *Nat Genet* 37(10): 1090-1097.
- Mito Y, Henikoff JG, Henikoff S (2007) Histone replacement marks the boundaries of cis-regulatory domains. *Science* 315(5817): 1408-1411.

- Mizuguchi G, Shen X, Landry J, Wu WH, Sen S et al. (2004) ATP-driven exchange of histone H2AZ variant catalyzed by SWR1 chromatin remodeling complex. *Science* 303(5656): 343-348.
- Mizzen CA, Yang XJ, Kokubo T, Brownell JE, Bannister AJ et al. (1996) The TAF(II)250 subunit of TFIID has histone acetyltransferase activity. *Cell* 87(7): 1261-1270.
- Monier K, Mouradian S, Sullivan KF (2007) DNA methylation promotes Aurora-B-driven phosphorylation of histone H3 in chromosomal subdomains. *J Cell Sci* 120(Pt 1): 101-114.
- Moon H, Filippova G, Loukinov D, Pugacheva E, Chen Q et al. (2005) CTCF is conserved from *Drosophila* to humans and confers enhancer blocking of the Fab-8 insulator. *EMBO Rep* 6(2): 165-170.
- Morris SA, Rao B, Garcia BA, Hake SB, Diaz RL et al. (2007) Identification of histone H3 lysine 36 acetylation as a highly conserved histone modification. *J Biol Chem* 282(10): 7632-7640.
- Mukhopadhyay R, Yu W, Whitehead J, Xu J, Lezcano M et al. (2004) The binding sites for the chromatin insulator protein CTCF map to DNA methylation-free domains genome-wide. *Genome Res* 14(8): 1594-1602.
- Murray K (1964) The Occurrence of Epsilon-N-Methyl Lysine in Histones. *Biochemistry* 3: 10-15.
- Nagy PL, Cleary ML, Brown PO, Lieb JD (2003) Genomewide demarcation of RNA polymerase II transcription units revealed by physical fractionation of chromatin. *Proc Natl Acad Sci U S A* 100(11): 6364-6369.
- Nakayama J, Rice JC, Strahl BD, Allis CD, Grewal SI (2001) Role of histone H3 lysine 9 methylation in epigenetic control of heterochromatin assembly. *Science* 292(5514): 110-113.
- Nallur GN, Prakash K, Weissman SM (1996) Multiplex selection technique (MuST): an approach to clone transcription factor binding sites. *Proc Natl Acad Sci U S A* 93(3): 1184-1189.
- Nan X, Ng HH, Johnson CA, Laherty CD, Turner BM et al. (1998) Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393(6683): 386-389.
- Narlikar GJ, Fan HY, Kingston RE (2002) Cooperation between complexes that regulate chromatin structure and transcription. *Cell* 108(4): 475-487.
- Nathan D, Ingvarsdottir K, Sterner DE, Bylebyl GR, Dokmanovic M et al. (2006) Histone sumoylation is a negative regulator in *Saccharomyces cerevisiae* and shows dynamic interplay with positive-acting histone modifications. *Genes Dev* 20(8): 966-976.
- Negre N, Hennetin J, Sun LV, Lavrov S, Bellis M et al. (2006) Chromosomal distribution of PcG proteins during *Drosophila* development. *PLoS Biol* 4(6): e170.
- Ng HH, Feng Q, Wang H, Erdjument-Bromage H, Tempst P et al. (2002) Lysine methylation within the globular domain of histone H3 by Dot1 is important for telomeric silencing and Sir protein association. *Genes Dev* 16(12): 1518-1527.

- Ng HH, Robert F, Young RA, Struhl K (2003) Targeted recruitment of Set1 histone methylase by elongating Pol II provides a localized mark and memory of recent transcriptional activity. *Mol Cell* 11(3): 709-719.
- Ng HH, Ciccone DN, Morshead KB, Oettinger MA, Struhl K (2003 b) Lysine-79 of histone H3 is hypomethylated at silenced loci in yeast and mammalian cells: a potential mechanism for position-effect variegation. *Proc Natl Acad Sci U S A* 100(4): 1820-1825.
- Nishida H, Suzuki T, Kondo S, Miura H, Fujimura Y et al. (2006) Histone H3 acetylated at lysine 9 in promoter is associated with low nucleosome density in the vicinity of transcription start site in human cell. *Chromosome Res* 14(2): 203-211.
- Nobrega MA, Ovcharenko I, Afzal V, Rubin EM (2003) Scanning human gene deserts for long-range enhancers. *Science* 302(5644): 413.
- Nowak SJ, Corces VG (2000) Phosphorylation of histone H3 correlates with transcriptionally active loci. *Genes Dev* 14(23): 3003-3013.
- Nowak SJ, Corces VG (2004) Phosphorylation of histone H3: a balancing act between chromosome condensation and transcriptional activation. *Trends Genet* 20(4): 214-220.
- Odom DT, Zizlsperger N, Gordon DB, Bell GW, Rinaldi NJ et al. (2004) Control of pancreas and liver gene expression by HNF transcription factors. *Science* 303(5662): 1378-1381.
- Odom DT, Dowell RD, Jacobsen ES, Gordon W, Danford TW et al. (2007) Tissue-specific transcriptional regulation has diverged significantly between human and mouse. *Nat Genet* 39(6): 730-732.
- Ogbourne S, Antalis TM (1998) Transcriptional control and the role of silencers in transcriptional regulation in eukaryotes. *Biochem J* 331 (Pt 1): 1-14.
- Ohlsson R, Renkawitz R, Lobanenko V (2001) CTCF is a uniquely versatile transcription regulator linked to epigenetics and disease. *Trends Genet* 17(9): 520-527.
- O'Neill LP, Turner BM (1995) Histone H4 acetylation distinguishes coding regions of the human genome from heterochromatin in a differentiation-dependent but transcription-independent manner. *EMBO J* 14(16): 3946-3957.
- O'Neill LP, Turner BM (2003) Immunoprecipitation of native chromatin: NChIP. *Methods* 31(1): 76-82.
- O'Neill LP, VerMilyea MD, Turner BM (2006) Epigenetic characterization of the early embryo with a chromatin immunoprecipitation protocol applicable to small cell populations. *Nat Genet* 38(7): 835-841.
- Orlando V, Paro R (1993) Mapping Polycomb-repressed domains in the bithorax complex using in vivo formaldehyde cross-linked chromatin. *Cell* 75(6): 1187-1198.
- Orlando V, Strutt H, Paro R (1997) Analysis of chromatin structure by in vivo formaldehyde cross-linking. *Methods* 11(2): 205-214.
- Orlando V (2000) Mapping chromosomal proteins in vivo by formaldehyde-crosslinked-chromatin immunoprecipitation. *Trends Biochem Sci* 25(3): 99-104.

- Orphanides G, Lagrange T, Reinberg D (1996) The general transcription factors of RNA polymerase II. *Genes Dev* 10(21): 2657-2683.
- Osborne CS, Chakalova L, Brown KE, Carter D, Horton A et al. (2004) Active genes dynamically colocalize to shared sites of ongoing transcription. *Nat Genet* 36(10): 1065-1071.
- Pan G, S. Tian, et al. (2007) Whole-Genome Analysis of Histone H3 Lysine 4 and Lysine 27 Methylation in Human Embryonic Stem Cells. *Cell Stem Cell* 1: 299-312.
- Pant V, Kurukuti S, Pugacheva E, Shamsuddin S, Mariano P et al. (2004) Mutation of a single CTCF target site within the H19 imprinting control region leads to loss of Igf2 imprinting and complex patterns of de novo methylation upon maternal inheritance. *Mol Cell Biol* 24(8): 3497-3504.
- Park YJ, Chodaparambil JV, Bao Y, McBryant SJ, Luger K (2005) Nucleosome assembly protein 1 exchanges histone H2A-H2B dimers and assists nucleosome sliding. *J Biol Chem* 280(3): 1817-1825.
- Patrinos GP, de Krom M, de Boer E, Langeveld A, Imam AM et al. (2004) Multiple interactions between regulatory regions are required to stabilize an active chromatin hub. *Genes Dev* 18(12): 1495-1509.
- Patterson BD, Davies DD (1969) Specificity of the enzymatic methylation of pea histone. *Biochem Biophys Res Commun* 34(6): 791-794.
- Pearson JC, Lemons D, McGinnis W (2005) Modulating Hox gene functions during animal body patterning. *Nat Rev Genet* 6(12): 893-904.
- Pena PV, Davrazou F, Shi X, Walter KL, Verkhusha VV et al. (2006) Molecular mechanism of histone H3K4me3 recognition by plant homeodomain of ING2. *Nature* 442(7098): 100-103.
- Pennacchio LA, Ahituv N, Moses AM, Prabhakar S, Nobrega MA et al. (2006) In vivo enhancer analysis of human conserved non-coding sequences. *Nature* 444(7118): 499-502.
- Pennacchio LA, Loots GG, Nobrega MA, Ovcharenko I (2007) Predicting tissue-specific enhancers in the human genome. *Genome Res* 17(2): 201-211.
- Pezzolesi MG, Zbuk KM, Waite KA, Eng C (2007) Comparative genomic and functional analyses reveal a novel cis-acting PTEN regulatory element as a highly conserved functional E-box motif deleted in Cowden syndrome. *Hum Mol Genet* 16(9): 1058-1071.
- Pikaart MJ, Recillas-Targa F, Felsenfeld G (1998) Loss of transcriptional activity of a transgene is accompanied by DNA methylation and histone deacetylation and is prevented by insulators. *Genes Dev* 12(18): 2852-2862.
- Pimanda JE, Silberstein L, Dominici M, Dekel B, Bowen M et al. (2006) Transcriptional link between blood and bone: the stem cell leukemia gene and its +19 stem cell enhancer are active in bone cells. *Mol Cell Biol* 26(7): 2615-2625.
- Pokholok DK, Harbison CT, Levine S, Cole M, Hannett NM et al. (2005) Genome-wide map of nucleosome acetylation and methylation in yeast. *Cell* 122(4): 517-527.

- Prabhakar S, Poulin F, Shoukry M, Afzal V, Rubin EM et al. (2006) Close sequence comparisons are sufficient to identify human cis-regulatory elements. *Genome Res* 16(7): 855-863.
- Prakash A, Tompa M (2005) Discovery of regulatory elements in vertebrates through comparative genomics. *Nat Biotechnol* 23(10): 1249-1256.
- Pray-Grant MG, Daniel JA, Schieltz D, Yates JR, 3rd, Grant PA (2005) Chd1 chromodomain links histone H3 methylation with SAGA- and SLIK-dependent acetylation. *Nature* 433(7024): 434-438.
- Prelich G (2002) RNA polymerase II carboxy-terminal domain kinases: emerging clues to their function. *Eukaryot Cell* 1(2): 153-162.
- Prendergast GC, Ziff EB (1989) DNA-binding motif. *Nature* 341(6241): 392.
- Pruitt KD, Tatusova T, Maglott DR (2007) NCBI reference sequences (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* 35(Database issue): D61-65.
- Puig O, Casparly F, Rigaut G, Rutz B, Bouveret E et al. (2001) The tandem affinity purification (TAP) method: a general procedure of protein complex purification. *Methods* 24(3): 218-229.
- Quackenbush J (2002) Microarray data normalization and transformation. *Nat Genet* 32 Suppl: 496-501.
- Rada-Iglesias A, Enroth S, Ameer A, Koch CM, Clelland GK et al. (2007) Butyrate mediates decrease of histone acetylation centered on transcription start sites and down-regulation of associated genes. *Genome Res* 17(6): 708-719.
- Raghuraman MK, Winzeler EA, Collingwood D, Hunt S, Wodicka L et al. (2001) Replication dynamics of the yeast genome. *Science* 294(5540): 115-121.
- Ramalho-Santos M, Yoon S, Matsuzaki Y, Mulligan RC, Melton DA (2002) "Stemness": transcriptional profiling of embryonic and adult stem cells. *Science* 298(5593): 597-600.
- Rao B, Shibata Y, Strahl BD, Lieb JD (2005) Dimethylation of histone H3 at lysine 36 demarcates regulatory and nonregulatory chromatin genome-wide. *Mol Cell Biol* 25(21): 9447-9459.
- Rea S, Eisenhaber F, O'Carroll D, Strahl BD, Sun ZW et al. (2000) Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature* 406(6796): 593-599.
- Recillas-Targa F, Pikaart MJ, Burgess-Beusse B, Bell AC, Litt MD et al. (2002) Position-effect protection and enhancer blocking by the chicken beta-globin insulator are separable activities. *Proc Natl Acad Sci U S A* 99(10): 6883-6888.
- Reid JL, Iyer VR, Brown PO, Struhl K (2000) Coordinate regulation of yeast ribosomal protein genes is associated with targeted recruitment of Esa1 histone acetylase. *Mol Cell* 6(6): 1297-1307.
- Reik W, Dean W, Walter J (2001) Epigenetic reprogramming in mammalian development. *Science* 293(5532): 1089-1093.
- Reik W, Walter J (2001) Genomic imprinting: parental influence on the genome. *Nat Rev Genet* 2(1): 21-32.
- Reik W (2007) Stability and flexibility of epigenetic gene regulation in mammalian development. *Nature* 447(7143): 425-432.

- Reinke H, Horz W (2003) Histones are first hyperacetylated and then lose contact with the activated PHO5 promoter. *Mol Cell* 11(6): 1599-1607.
- Ren B, Maniatis T (1998) Regulation of Drosophila Adh promoter switching by an initiator-targeted repression mechanism. *EMBO J* 17(4): 1076-1086.
- Ren B, Robert F, Wyrick JJ, Aparicio O, Jennings EG et al. (2000) Genome-wide location and function of DNA binding proteins. *Science* 290(5500): 2306-2309.
- Ren B, Cam H, Takahashi Y, Volkert T, Terragni J et al. (2002) E2F integrates cell cycle progression with DNA repair, replication, and G(2)/M checkpoints. *Genes Dev* 16(2): 245-256.
- Reubinoff BE, Pera MF, Fong CY, Trounson A, Bongso A (2000) Embryonic stem cell lines from human blastocysts: somatic differentiation in vitro. *Nat Biotechnol* 18(4): 399-404.
- Reya T, Morrison SJ, Clarke MF, Weissman IL (2001) Stem cells, cancer, and cancer stem cells. *Nature* 414(6859): 105-111.
- Richards EJ, Elgin SC (2002) Epigenetic codes for heterochromatin formation and silencing: rounding up the usual suspects. *Cell* 108(4): 489-500.
- Ringrose L, Ehret H, Paro R (2004) Distinct contributions of histone H3 lysine 9 and 27 methylation to locus-specific stability of polycomb complexes. *Mol Cell* 16(4): 641-653.
- Ringrose L, Paro R (2004) Epigenetic regulation of cellular memory by the Polycomb and Trithorax group proteins. *Annu Rev Genet* 38: 413-443.
- Rinn JL, Euskirchen G, Bertone P, Martone R, Luscombe NM et al. (2003) The transcriptional activity of human Chromosome 22. *Genes Dev* 17(4): 529-540.
- Robb L, Begley CG (1997) The SCL/TAL1 gene: roles in normal and malignant haematopoiesis. *Bioessays* 19(7): 607-613.
- Robert F, Pokholok DK, Hannett NM, Rinaldi NJ, Chandy M et al. (2004) Global position and recruitment of HATs and HDACs in the yeast genome. *Mol Cell* 16(2): 199-209.
- Robertson G, Hirst M, Bainbridge M, Bilenky M, Zhao Y et al. (2007) Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nat Methods* 4(8): 651-657.
- Roby D, Suka Y, Xenarios I, Kurdistani SK, Wang A et al. (2002) Microarray deacetylation maps determine genome-wide functions for yeast histone deacetylases. *Cell* 109(4): 437-446.
- Roby D, Grunstein M (2003) Genomewide histone acetylation microarrays. *Methods* 31(1): 83-89.
- Roguev A, Schaft D, Shevchenko A, Pijnappel WW, Wilm M et al. (2001) The *Saccharomyces cerevisiae* Set1 complex includes an Ash2 homologue and methylates histone 3 lysine 4. *EMBO J* 20(24): 7137-7148.
- Roh TY, Ngau WC, Cui K, Landsman D, Zhao K (2004) High-resolution genome-wide mapping of histone modifications. *Nat Biotechnol* 22(8): 1013-1016.
- Roh TY, Cuddapah S, Zhao K (2005) Active chromatin domains are defined by acetylation islands revealed by genome-wide mapping. *Genes Dev* 19(5): 542-552.

- Roh TY, Cuddapah S, Cui K, Zhao K (2006) The genomic landscape of histone modifications in human T cells. *Proc Natl Acad Sci U S A* 103(43): 15782-15787.
- Roh TY, Wei G, Farrell CM, Zhao K (2007) Genome-wide prediction of conserved and nonconserved enhancers by histone acetylation patterns. *Genome Res* 17(1): 74-81.
- Roth SY, Denu JM, Allis CD (2001) Histone acetyltransferases. *Annu Rev Biochem* 70: 81-120.
- Rudolph T, Yonezawa M, Lein S, Heidrich K, Kubicek S et al. (2007) Heterochromatin formation in *Drosophila* is initiated through active removal of H3K4 methylation by the LSD1 homolog SU(VAR)3-3. *Mol Cell* 26(1): 103-115.
- Ruthenburg AJ, Li H, Patel DJ, Allis CD (2007) Multivalent engagement of chromatin modifications by linked binding modules. *Nat Rev Mol Cell Biol* 8(12): 983-994.
- Sabo PJ, Humbert R, Hawrylycz M, Wallace JC, Dorschner MO et al. (2004) Genome-wide identification of DNaseI hypersensitive sites using active chromatin sequence libraries. *Proc Natl Acad Sci U S A* 101(13): 4537-4542.
- Sabo PJ, Kuehn MS, Thurman R, Johnson BE, Johnson EM et al. (2006) Genome-scale mapping of DNase I sensitivity in vivo using tiling DNA microarrays. *Nat Methods* 3(7): 511-518.
- Sanchez M, Gottgens B, Sinclair AM, Stanley M, Begley CG et al. (1999) An SCL 3' enhancer targets developing endothelium together with embryonic and adult haematopoietic progenitors. *Development* 126(17): 3891-3904.
- Sanchez MJ, Bockamp EO, Miller J, Gambardella L, Green AR (2001) Selective rescue of early haematopoietic progenitors in *Scl(-/-)* mice by expressing *Scl* under the control of a stem cell enhancer. *Development* 128(23): 4815-4827.
- Sandelin A, Alkema W, Engstrom P, Wasserman WW, Lenhard B (2004) JASPAR: an open-access database for eukaryotic transcription factor binding profiles. *Nucleic Acids Res* 32(Database issue): D91-94.
- Sandelin A, Carninci P, Lenhard B, Ponjavic J, Hayashizaki Y et al. (2007) Mammalian RNA polymerase II core promoters: insights from genome-wide studies. *Nat Rev Genet* 8(6): 424-436.
- Sanders SL, Portoso M, Mata J, Bahler J, Allshire RC et al. (2004) Methylation of histone H4 lysine 20 controls recruitment of Crb2 to sites of DNA damage. *Cell* 119(5): 603-614.
- Santos-Rosa H, Schneider R, Bannister AJ, Sherriff J, Bernstein BE et al. (2002) Active genes are trimethylated at K4 of histone H3. *Nature* 419(6905): 407-411.
- Sarraf SA, Stancheva I (2004) Methyl-CpG binding protein MBD1 couples histone H3 methylation at lysine 9 by SETDB1 to DNA replication and chromatin assembly. *Mol Cell* 15(4): 595-605.
- Sassone-Corsi P, Mizzen CA, Cheung P, Crosio C, Monaco L et al. (1999) Requirement of Rsk-2 for epidermal growth factor-activated phosphorylation of histone H3. *Science* 285(5429): 886-891.
- Savas U, Hsu MH, Johnson EF (2003) Differential regulation of human CYP4A genes by peroxisome proliferators and dexamethasone. *Arch Biochem Biophys* 409(1): 212-220.

- Saxonov S, Berg P, Brutlag DL (2006) A genome-wide analysis of CpG dinucleotides in the human genome distinguishes two distinct classes of promoters. *Proc Natl Acad Sci U S A* 103(5): 1412-1417.
- Schaft D, Roguev A, Kotovic KM, Shevchenko A, Sarov M et al. (2003) The histone 3 lysine 36 methyltransferase, SET2, is involved in transcriptional elongation. *Nucleic Acids Res* 31(10): 2475-2482.
- Schena M, Shalon D, Davis RW, Brown PO (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 270(5235): 467-470.
- Scherf M, Klingenhoff A, Werner T (2000) Highly specific localization of promoter regions in large genomic sequences by PromoterInspector: a novel context analysis approach. *J Mol Biol* 297(3): 599-606.
- Schneider R, Bannister AJ, Myers FA, Thorne AW, Crane-Robinson C et al. (2004) Histone H3 lysine 4 methylation patterns in higher eukaryotic genes. *Nat Cell Biol* 6(1): 73-77.
- Schotta G, Lachner M, Sarma K, Ebert A, Sengupta R et al. (2004) A silencing pathway to induce H3-K9 and H4-K20 trimethylation at constitutive heterochromatin. *Genes Dev* 18(11): 1251-1262.
- Schubeler D, Scalzo D, Kooperberg C, van Steensel B, Delrow J et al. (2002) Genome-wide DNA replication profile for *Drosophila melanogaster*: a link between transcription and replication timing. *Nat Genet* 32(3): 438-442.
- Schubeler D, MacAlpine DM, Scalzo D, Wirbelauer C, Kooperberg C et al. (2004) The histone modification pattern of active genes revealed through genome-wide chromatin analysis of a higher eukaryote. *Genes Dev* 18(11): 1263-1271.
- Schuettengruber B, Chourrout D, Vervoort M, Leblanc B, Cavalli G (2007) Genome regulation by polycomb and trithorax proteins. *Cell* 128(4): 735-745.
- Schurter BT, Koh SS, Chen D, Bunick GJ, Harp JM et al. (2001) Methylation of histone H3 by coactivator-associated arginine methyltransferase 1. *Biochemistry* 40(19): 5747-5756.
- Schwartz YB, Kahn TG, Nix DA, Li XY, Bourgon R et al. (2006) Genome-wide analysis of Polycomb targets in *Drosophila melanogaster*. *Nat Genet* 38(6): 700-705.
- Secombe J, Li L, Carlos L, Eisenman RN (2007) The Trithorax group protein Lid is a trimethyl histone H3K4 demethylase required for dMyc-induced cell growth. *Genes Dev* 21(5): 537-551.
- Sertil O, Vemula A, Salmon SL, Morse RH, Lowry CV (2007) Direct role for the Rpd3 complex in transcriptional induction of the anaerobic DAN/TIR genes in yeast. *Mol Cell Biol* 27(6): 2037-2047.
- Seta N, Kuwana M (2007) Human circulating monocytes as multipotential progenitors. *Keio J Med* 56(2): 41-47.
- Shahbazian MD, Grunstein M (2007) Functions of site-specific histone acetylation and deacetylation. *Annu Rev Biochem* 76: 75-100.

- Sharma VM, Tomar RS, Dempsey AE, Reese JC (2007) Histone deacetylases RPD3 and HOS2 regulate the transcriptional activation of DNA damage-inducible genes. *Mol Cell Biol* 27(8): 3199-3210.
- Sharpe JA, Summerhill RJ, Vyas P, Gourdon G, Higgs DR et al. (1993) Role of upstream DNase I hypersensitive sites in the regulation of human alpha globin gene expression. *Blood* 82(5): 1666-1671.
- Shi Y, Lan F, Matson C, Mulligan P, Whetstine JR et al. (2004) Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119(7): 941-953.
- Shi X, Hong T, Walter KL, Ewalt M, Michishita E et al. (2006) ING2 PHD domain links histone H3 lysine 4 methylation to active gene repression. *Nature* 442(7098): 96-99.
- Shiio Y, Eisenman RN (2003) Histone sumoylation is associated with transcriptional repression. *Proc Natl Acad Sci U S A* 100(23): 13225-13230.
- Shiio Y, Eisenman RN (2003) Histone sumoylation is associated with transcriptional repression. *Proc Natl Acad Sci U S A* 100(23): 13225-13230.
- Shilatifard A (2006) Chromatin modifications by methylation and ubiquitination: implications in the regulation of gene expression. *Annu Rev Biochem* 75: 243-269.
- Shipp MA, Ross KN, Tamayo P, Weng AP, Kutok JL et al. (2002) Diffuse large B-cell lymphoma outcome prediction by gene-expression profiling and supervised machine learning. *Nat Med* 8(1): 68-74.
- Shogren-Knaak M, Ishii H, Sun JM, Pazin MJ, Davie JR et al. (2006) Histone H4-K16 acetylation controls chromatin structure and protein interactions. *Science* 311(5762): 844-847.
- Siepel A, Bejerano G, Pedersen JS, Hinrichs AS, Hou M et al. (2005) Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome Res* 15(8): 1034-1050.
- Silverstein RA, Ekwall K (2005) Sin3: a flexible regulator of global gene expression and genome stability. *Curr Genet* 47(1): 1-17.
- Simonis M, Klous P, Splinter E, Moshkin Y, Willemsen R et al. (2006) Nuclear organization of active and inactive chromatin domains uncovered by chromosome conformation capture-on-chip (4C). *Nat Genet* 38(11): 1348-1354.
- Sims RJ, 3rd, Belotserkovskaya R, Reinberg D (2004) Elongation by RNA polymerase II: the short and long of it. *Genes Dev* 18(20): 2437-2468.
- Sims RJ, 3rd, Chen CF, Santos-Rosa H, Kouzarides T, Patel SS et al. (2005) Human but not yeast CHD1 binds directly and selectively to histone H3 methylated at lysine 4 via its tandem chromodomains. *J Biol Chem* 280(51): 41789-41792.
- Sinclair AM, Gottgens B, Barton LM, Stanley ML, Pardanaud L et al. (1999) Distinct 5' SCL enhancers direct transcription to developing brain, spinal cord, and endothelium: neural expression is mediated by GATA factor binding sites. *Dev Biol* 209(1): 128-142.
- Singh-Gasson S, Green RD, Yue Y, Nelson C, Blattner F et al. (1999) Maskless fabrication of light-directed oligonucleotide microarrays using a digital micromirror array. *Nat Biotechnol* 17(10): 974-978.

- Sinha I, Wiren M, Ekwall K (2006) Genome-wide patterns of histone modifications in fission yeast. *Chromosome Res* 14(1): 95-105.
- Smith AD, Sumazin P, Das D, Zhang MQ (2005) Mining ChIP-chip data for transcription factor and cofactor binding sites. *Bioinformatics* 21 Suppl 1: i403-412.
- Solomon MJ, Varshavsky A (1985) Formaldehyde-mediated DNA-protein crosslinking: a probe for in vivo chromatin structures. *Proc Natl Acad Sci U S A* 82(19): 6470-6474.
- Solomon MJ, Larsen PL, Varshavsky A (1988) Mapping protein-DNA interactions in vivo with formaldehyde: evidence that histone H4 is retained on a highly transcribed gene. *Cell* 53(6): 937-947.
- Spilianakis CG, Lalioti MD, Town T, Lee GR, Flavell RA (2005) Interchromosomal associations between alternatively expressed loci. *Nature* 435(7042): 637-645.
- Spivakov M, Fisher AG (2007) Epigenetic signatures of stem-cell identity. *Nat Rev Genet* 8(4): 263-271.
- Splinter E, Heath H, Kooren J, Palstra RJ, Klous P et al. (2006) CTCF mediates long-range chromatin looping and local histone modification in the beta-globin locus. *Genes Dev* 20(17): 2349-2354.
- Sproul D, Gilbert N, Bickmore WA (2005) The role of chromatin structure in regulating the expression of clustered genes. *Nat Rev Genet* 6(10): 775-781.
- Stein LD, Bao Z, Blasiar D, Blumenthal T, Brent MR et al. (2003) The genome sequence of *Caenorhabditis briggsae*: a platform for comparative genomics. *PLoS Biol* 1(2): E45.
- Stephanou A, Scarabelli TM, Townsend PA, Bell R, Yellon D et al. (2002) The carboxyl-terminal activation domain of the STAT-1 transcription factor enhances ischemia/reperfusion-induced apoptosis in cardiac myocytes. *FASEB J* 16(13): 1841-1843.
- Sterner DE, Berger SL (2000) Acetylation of histones and transcription-related factors. *Microbiol Mol Biol Rev* 64(2): 435-459.
- Strahl BD, Allis CD (2000) The language of covalent histone modifications. *Nature* 403(6765): 41-45.
- Strahl BD, Briggs SD, Brame CJ, Caldwell JA, Koh SS et al. (2001) Methylation of histone H4 at arginine 3 occurs in vivo and is mediated by the nuclear receptor coactivator PRMT1. *Curr Biol* 11(12): 996-1000.
- Straka C, Horz W (1991) A functional role for nucleosomes in the repression of a yeast promoter. *EMBO J* 10(2): 361-368.
- Suka N, Suka Y, Carmen AA, Wu J, Grunstein M (2001) Highly specific antibodies determine histone acetylation site usage in yeast heterochromatin and euchromatin. *Mol Cell* 8(2): 473-479.
- Sumer H, Craig JM, Sibson M, Choo KH (2003) A rapid method of genomic array analysis of scaffold/matrix attachment regions (S/MARs) identifies a 2.5-Mb region of enhanced scaffold/matrix attachment at a human neocentromere. *Genome Res* 13(7): 1737-1743.
- Sun FL, Elgin SC (1999) Putting boundaries on silence. *Cell* 99(5): 459-462.
- Sundstrom C, Nilsson K (1976) Establishment and characterization of a human histiocytic lymphoma cell line (U-937). *Int J Cancer* 17(5): 565-577.

- Surani MA, Hayashi K, Hajkova P (2007) Genetic and epigenetic regulators of pluripotency. *Cell* 128(4): 747-762.
- Szabo P, Tang SH, Rentsendorj A, Pfeifer GP, Mann JR (2000) Maternal-specific footprints at putative CTCF sites in the H19 imprinting control region give evidence for insulator function. *Curr Biol* 10(10): 607-610.
- Szutorisz H, Dillon N, Tora L (2005) The role of enhancers as centres for general transcription factor recruitment. *Trends Biochem Sci* 30(11): 593-599.
- Szutorisz H, Canzonetta C, Georgiou A, Chow CM, Tora L et al. (2005 b) Formation of an active tissue-specific chromatin domain initiated by epigenetic marking at the embryonic stem cell stage. *Mol Cell Biol* 25(5): 1804-1820.
- Taft RJ, Pheasant M, Mattick JS (2007) The relationship between non-protein-coding DNA and eukaryotic complexity. *Bioessays* 29(3): 288-299.
- Talasz H, Lindner HH, Sarg B, Helliger W (2005) Histone H4-lysine 20 monomethylation is increased in promoter and coding regions of active genes and correlates with hyperacetylation. *J Biol Chem* 280(46): 38814-38822.
- Tanimoto K, Liu Q, Bungert J, Engel JD (1999) Effects of altered gene order or orientation of the locus control region on human beta-globin gene expression in mice. *Nature* 398(6725): 344-348.
- Taverna SD, Li H, Ruthenburg AJ, Allis CD, Patel DJ (2007) How chromatin-binding modules interpret histone modifications: lessons from professional pocket pickers. *Nat Struct Mol Biol* 14(11): 1025-1040.
- Thomas MC, Chiang CM (2006) The general transcription machinery and general cofactors. *Crit Rev Biochem Mol Biol* 41(3): 105-178.
- Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ et al. (1998) Embryonic stem cell lines derived from human blastocysts. *Science* 282(5391): 1145-1147.
- Thomson S, Clayton AL, Hazzalin CA, Rose S, Barratt MJ et al. (1999) The nucleosomal response associated with immediate-early gene induction is mediated via alternative MAP kinase cascades: MSK1 as a potential histone H3/HMG-14 kinase. *EMBO J* 18(17): 4779-4793.
- Tian B, Nowak DE, Jamaluddin M, Wang S, Brasier AR (2005) Identification of direct genomic targets downstream of the nuclear factor-kappaB transcription factor mediating tumor necrosis factor signaling. *J Biol Chem* 280(17): 17435-17448.
- Tolhuis B, Palstra RJ, Splinter E, Grosveld F, de Laat W (2002) Looping and interaction between hypersensitive sites in the active beta-globin locus. *Mol Cell* 10(6): 1453-1465.
- Tolhuis B, de Wit E, Muijers I, Teunissen H, Talhout W et al. (2006) Genome-wide profiling of PRC1 and PRC2 Polycomb chromatin binding in *Drosophila melanogaster*. *Nat Genet* 38(6): 694-699.
- Torres-Padilla ME, Parfitt DE, Kouzarides T, Zernicka-Goetz M (2007) Histone arginine methylation regulates pluripotency in the early mouse embryo. *Nature* 445(7124): 214-218.

- Tse C, Sera T, Wolffe AP, Hansen JC (1998) Disruption of higher-order folding by core histone acetylation dramatically enhances transcription of nucleosomal arrays by RNA polymerase III. *Mol Cell Biol* 18(8): 4629-4638.
- Tsukada Y, Fang J, Erdjument-Bromage H, Warren ME, Borchers CH et al. (2006) Histone demethylation by a family of JmjC domain-containing proteins. *Nature* 439(7078): 811-816.
- Tuerk C, Gold L (1990) Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* 249(4968): 505-510.
- Turner BM (2000) Histone acetylation and an epigenetic code. *Bioessays* 22(9): 836-845.
- Uesugi M, Nyanguile O, Lu H, Levine AJ, Verdine GL (1997) Induced alpha helix in the VP16 activation domain upon binding to a human TAF. *Science* 277(5330): 1310-1313.
- Vakoc CR, Mandat SA, Olenchock BA, Blobel GA (2005) Histone H3 lysine 9 methylation and HP1gamma are associated with transcription elongation through mammalian chromatin. *Mol Cell Biol* 25(3): 381-391.
- Vakoc CR, Sachdeva MM, Wang H, Blobel GA (2006) Profile of histone lysine methylation across transcribed mammalian chromatin. *Mol Cell Biol* 26(24): 9185-9195.
- Valadez-Graham V, Razin SV, Recillas-Targa F (2004) CTCF-dependent enhancer blockers at the upstream region of the chicken alpha-globin gene domain. *Nucleic Acids Res* 32(4): 1354-1362.
- Vallier L, Alexander M, Pedersen R (2007) Conditional gene expression in human embryonic stem cells. *Stem Cells* 25(6): 1490-1497.
- van Leeuwen F, Gafken PR, Gottschling DE (2002) Dot1p modulates silencing in yeast by methylation of the nucleosome core. *Cell* 109(6): 745-756.
- van Steensel B, Henikoff S (2000) Identification of in vivo DNA targets of chromatin proteins using tethered dam methyltransferase. *Nat Biotechnol* 18(4): 424-428.
- Varshavsky AJ, Sundin O, Bohn M (1979) A stretch of "late" SV40 viral DNA about 400 bp long which includes the origin of replication is specifically exposed in SV40 minichromosomes. *Cell* 16(2): 453-466.
- Vavouri T, Elgar G (2005) Prediction of cis-regulatory elements using binding site matrices--the successes, the failures and the reasons for both. *Curr Opin Genet Dev* 15(4): 395-402.
- Venter JC, Adams MD, Myers EW, Li PW, Mural RJ et al. (2001) The sequence of the human genome. *Science* 291(5507): 1304-1351.
- Vermeulen M, Mulder KW, Denissov S, Pijnappel WW, van Schaik FM et al. (2007) Selective anchoring of TFIID to nucleosomes by trimethylation of histone H3 lysine 4. *Cell* 131(1): 58-69.
- Vettese-Dadey M, Grant PA, Hebbes TR, Crane-Robinson C, Allis CD et al. (1996) Acetylation of histone H4 plays a primary role in enhancing transcription factor binding to nucleosomal DNA in vitro. *EMBO J* 15(10): 2508-2518.
- Vire E, Brenner C, Deplus R, Blanchon L, Fraga M et al. (2006) The Polycomb group protein EZH2 directly controls DNA methylation. *Nature* 439(7078): 871-874.

- Visel A, Minovitsky S, Dubchak I, Pennacchio LA (2007) VISTA Enhancer Browser--a database of tissue-specific human enhancers. *Nucleic Acids Res* 35(Database issue): D88-92.
- Vogelauer M, Wu J, Suka N, Grunstein M (2000) Global histone acetylation and deacetylation in yeast. *Nature* 408(6811): 495-498.
- Vostrov AA, Quitschke WW (1997) The zinc finger protein CTCF binds to the APBbeta domain of the amyloid beta-protein precursor promoter. Evidence for a role in transcriptional activation. *J Biol Chem* 272(52): 33353-33359.
- Vyas P, Vickers MA, Picketts DJ, Higgs DR (1995) Conservation of position and sequence of a novel, widely expressed gene containing the major human alpha-globin regulatory element. *Genomics* 29(3): 679-689.
- Waddington C (1942) The epigenotype. *Endeavour* 1: 18-20.
- Walker GT, Fraiser MS, Schram JL, Little MC, Nadeau JG et al. (1992) Strand displacement amplification--an isothermal, in vitro DNA amplification technique. *Nucleic Acids Res* 20(7): 1691-1696.
- Wallace JA, Felsenfeld G (2007) We gather together: insulators and genome organization. *Curr Opin Genet Dev* 17(5): 400-407.
- Walter J, Dever CA, Biggin MD (1994) Two homeo domain proteins bind with similar specificity to a wide range of DNA sites in *Drosophila* embryos. *Genes Dev* 8(14): 1678-1692.
- Wang J, Mager J, Chen Y, Schneider E, Cross JC et al. (2001) Imprinted X inactivation maintained by a mouse Polycomb group gene. *Nat Genet* 28(4): 371-375.
- Wang H, Huang ZQ, Xia L, Feng Q, Erdjument-Bromage H et al. (2001 b) Methylation of histone H4 at arginine 3 facilitating transcriptional activation by nuclear hormone receptor. *Science* 293(5531): 853-857.
- Wang J, Mager J, Schnedier E, Magnuson T (2002) The mouse PcG gene *eed* is required for Hox gene repression and extraembryonic development. *Mamm Genome* 13(9): 493-503.
- Wang Y, Wysocka J, Sayegh J, Lee YH, Perlin JR et al. (2004) Human PAD4 regulates histone arginine methylation levels via demethylimination. *Science* 306(5694): 279-283.
- Wang H, Wang L, Erdjument-Bromage H, Vidal M, Tempst P et al. (2004) Role of histone H2A ubiquitination in Polycomb silencing. *Nature* 431(7010): 873-878.
- Watt F, Molloy PL (1988) Cytosine methylation prevents binding to DNA of a HeLa cell transcription factor required for optimal expression of the adenovirus major late promoter. *Genes Dev* 2(9): 1136-1143.
- Weber M, Hellmann I, Stadler MB, Ramos L, Paabo S et al. (2007) Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome. *Nat Genet* 39(4): 457-466.
- Wei Y, Mizzen CA, Cook RG, Gorovsky MA, Allis CD (1998) Phosphorylation of histone H3 at serine 10 is correlated with chromosome condensation during mitosis and meiosis in *Tetrahymena*. *Proc Natl Acad Sci U S A* 95(13): 7480-7484.

- Wei CL, Wu Q, Vega VB, Chiu KP, Ng P et al. (2006) A global map of p53 transcription-factor binding sites in the human genome. *Cell* 124(1): 207-219.
- Weinmann AS, Bartley SM, Zhang T, Zhang MQ, Farnham PJ (2001) Use of chromatin immunoprecipitation to clone novel E2F target promoters. *Mol Cell Biol* 21(20): 6820-6832.
- Weinmann AS, Yan PS, Oberley MJ, Huang TH, Farnham PJ (2002) Isolating human transcription factor targets by coupling chromatin immunoprecipitation and CpG island microarray analysis. *Genes Dev* 16(2): 235-244.
- Weintraub H, Groudine M (1976) Chromosomal subunits in active genes have an altered conformation. *Science* 193(4256): 848-856.
- Wells J, Graveel CR, Bartley SM, Madore SJ, Farnham PJ (2002) The identification of E2F1-specific target genes. *Proc Natl Acad Sci U S A* 99(6): 3890-3895.
- West AG, Huang S, Gaszner M, Litt MD, Felsenfeld G (2004) Recruitment of histone modifications by USF proteins at a vertebrate barrier element. *Mol Cell* 16(3): 453-463.
- Whetstine JR, Nottke A, Lan F, Huarte M, Smolikov S et al. (2006) Reversal of histone lysine trimethylation by the JMJD2 family of histone demethylases. *Cell* 125(3): 467-481.
- White KP, Rifkin SA, Hurban P, Hogness DS (1999) Microarray analysis of *Drosophila* development during metamorphosis. *Science* 286(5447): 2179-2184.
- White EJ, Emanuelsson O, Scalzo D, Royce T, Kosak S et al. (2004) DNA replication-timing analysis of human chromosome 22 at high resolution and different developmental states. *Proc Natl Acad Sci U S A* 101(51): 17771-17776.
- Whitehouse I, Flaus A, Cairns BR, White MF, Workman JL et al. (1999) Nucleosome mobilization catalysed by the yeast SWI/SNF complex. *Nature* 400(6746): 784-787.
- Wiblin AE, Cui W, Clark AJ, Bickmore WA (2005) Distinctive nuclear organisation of centromeres and regions involved in pluripotency in human embryonic stem cells. *J Cell Sci* 118(Pt 17): 3861-3868.
- Wingender E, Chen X, Hehl R, Karas H, Liebich I et al. (2000) TRANSFAC: an integrated system for gene expression regulation. *Nucleic Acids Res* 28(1): 316-319.
- Wiren M, Silverstein RA, Sinha I, Walfridsson J, Lee HM et al. (2005) Genomewide analysis of nucleosome density histone acetylation and HDAC function in fission yeast. *EMBO J* 24(16): 2906-2918.
- Woodfine K, Fiegler H, Beare DM, Collins JE, McCann OT et al. (2004) Replication timing of the human genome. *Hum Mol Genet* 13(2): 191-202.
- Woolfe A, Goodson M, Goode DK, Snell P, McEwen GK et al. (2005) Highly conserved non-coding sequences are associated with vertebrate development. *PLoS Biol* 3(1): e7.
- Wright WE, Binder M, Funk W (1991) Cyclic amplification and selection of targets (CASTing) for the myogenin consensus binding site. *Mol Cell Biol* 11(8): 4104-4110.

- Wu C (1980) The 5' ends of *Drosophila* heat shock genes in chromatin are hypersensitive to DNase I. *Nature* 286(5776): 854-860.
- Wyrick JJ, Aparicio JG, Chen T, Barnett JD, Jennings EG et al. (2001) Genome-wide distribution of ORC and MCM proteins in *S. cerevisiae*: high-resolution mapping of replication origins. *Science* 294(5550): 2357-2360.
- Wysocka J, Swigut T, Milne TA, Dou Y, Zhang X et al. (2005) WDR5 associates with histone H3 methylated at K4 and is essential for H3 K4 methylation and vertebrate development. *Cell* 121(6): 859-872.
- Wysocka J, Swigut T, Xiao H, Milne TA, Kwon SY et al. (2006) A PHD finger of NURF couples histone H3 lysine 4 trimethylation with chromatin remodelling. *Nature* 442(7098): 86-90.
- Xi H, Shulha HP, Lin JM, Vales TR, Fu Y et al. (2007) Identification and characterization of cell type-specific and ubiquitous chromatin regulatory structures in the human genome. *PLoS Genet* 3(8): e136.
- Xiao T, Hall H, Kizer KO, Shibata Y, Hall MC et al. (2003) Phosphorylation of RNA polymerase II CTD regulates H3 methylation in yeast. *Genes Dev* 17(5): 654-663.
- Xie X, Lu J, Kulbokas EJ, Golub TR, Mootha V et al. (2005) Systematic discovery of regulatory motifs in human promoters and 3' UTRs by comparison of several mammals. *Nature* 434(7031): 338-345.
- Xie X, Mikkelsen TS, Gnirke A, Lindblad-Toh K, Kellis M et al. (2007) Systematic discovery of regulatory motifs in conserved regions of the human genome, including thousands of CTCF insulator sites. *Proc Natl Acad Sci U S A* 104(17): 7145-7150.
- Xu W, Aparicio JG, Aparicio OM, Tavare S (2006) Genome-wide mapping of ORC and Mcm2p binding sites on tiling arrays and identification of essential ARS consensus sequences in *S. cerevisiae*. *BMC Genomics* 7: 276.
- Yamane K, Toumazou C, Tsukada Y, Erdjument-Bromage H, Tempst P et al. (2006) JHDM2A, a JmjC-containing H3K9 demethylase, facilitates transcription activation by androgen receptor. *Cell* 125(3): 483-495.
- Yamane K, Tateishi K, Klose RJ, Fang J, Fabrizio LA et al. (2007) PLU-1 is an H3K4 demethylase involved in transcriptional repression and breast cancer cell proliferation. *Mol Cell* 25(6): 801-812.
- Yan Q, Moreland RJ, Conaway JW, Conaway RC (1999) Dual roles for transcription factor IIF in promoter escape by RNA polymerase II. *J Biol Chem* 274(50): 35668-35675.
- Yang A, Zhu Z, Kapranov P, McKeon F, Church GM et al. (2006) Relationships between p63 binding, DNA sequence, transcription activity, and biological function in human cells. *Mol Cell* 24(4): 593-602.
- Yauk CL, Bois PR, Jeffreys AJ (2003) High-resolution sperm typing of meiotic recombination in the mouse MHC Ebeta gene. *EMBO J* 22(6): 1389-1397.

- Yeh RF, Lim LP, Burge CB (2001) Computational inference of homologous gene structures in the human genome. *Genome Res* 11(5): 803-816.
- Yokoyama A, Wang Z, Wysocka J, Sanyal M, Aufiero DJ et al. (2004) Leukemia proto-oncoprotein MLL forms a SET1-like histone methyltransferase complex with menin to regulate Hox gene expression. *Mol Cell Biol* 24(13): 5639-5649.
- Young RA (1991) RNA polymerase II. *Annu Rev Biochem* 60: 689-715.
- Young RA (2000) Biomedical discovery with DNA arrays. *Cell* 102(1): 9-15.
- Yu W, Ginjala V, Pant V, Chernukhin I, Whitehead J et al. (2004) Poly(ADP-ribosyl)ation regulates CTCF-dependent chromatin insulation. *Nat Genet* 36(10): 1105-1110.
- Yu MC, Lamming DW, Eskin JA, Sinclair DA, Silver PA (2006) The role of protein arginine methylation in the formation of silent chromatin. *Genes Dev* 20(23): 3249-3254.
- Yuan GC, Liu YJ, Dion MF, Slack MD, Wu LF et al. (2005) Genome-scale identification of nucleosome positions in *S. cerevisiae*. *Science* 309(5734): 626-630.
- Yusufzai TM, Felsenfeld G (2004) The 5'-HS4 chicken beta-globin insulator is a CTCF-dependent nuclear matrix-associated element. *Proc Natl Acad Sci U S A* 101(23): 8620-8624.
- Zardo G, Caiafa P (1998) The unmethylated state of CpG islands in mouse fibroblasts depends on the poly(ADP-ribosyl)ation process. *J Biol Chem* 273(26): 16517-16520.
- Zeng L, Zhou MM (2002) Bromodomain: an acetyl-lysine binding domain. *FEBS Lett* 513(1): 124-128.
- Zhang Y (2004) Molecular biology: no exception to reversibility. *Nature* 431(7009): 637-639.
- Zhao H, Dean A (2004) An insulator blocks spreading of histone acetylation and interferes with RNA polymerase II transfer between an enhancer and gene. *Nucleic Acids Res* 32(16): 4903-4919.
- Zhao H, Kim A, Song SH, Dean A (2006) Enhancer blocking by chicken beta-globin 5'-HS4: role of enhancer strength and insulator nucleosome depletion. *J Biol Chem* 281(41): 30573-30580.
- Zhao Z, Tavoosidana G, Sjolinder M, Gondor A, Mariano P et al. (2006 b) Circular chromosome conformation capture (4C) uncovers extensive networks of epigenetically regulated intra- and interchromosomal interactions. *Nat Genet* 38(11): 1341-1347.
- Zhao XD, X. Han, et al. (2007) Whole-Genome Mapping of Histone H3 Lys4 and 27 Trimethylations Reveals Distinct Genomic Compartments in Human Embryonic Stem Cells. *Cell Stem Cell* 1: 286-298.
- Zheng D, Frankish A, Baertsch R, Kapranov P, Reymond A et al. (2007) Pseudogenes in the ENCODE regions: consensus annotation, analysis of transcription, and evolution. *Genome Res* 17(6): 839-851.
- Zhou T, Chiang CM (2001) The intronless and TATA-less human TAF(II)55 gene contains a functional initiator and a downstream promoter element. *J Biol Chem* 276(27): 25503-25511.
- Zhu B, Zheng Y, Pham AD, Mandal SS, Erdjument-Bromage H et al. (2005) Monoubiquitination of human histone H2B: the factors involved and their roles in HOX gene regulation. *Mol Cell* 20(4): 601-611.