

Chapter 8

References

- Aaronson, J. S., Eckman, B., Blevins, R. A., Borkowski, J. A., Myerson, J., Imran, S., *et al.* (1996). Toward the development of a gene index to the human genome: an assessment of the nature of high-throughput EST sequence data. *Genome Res* **6**: 829-45.
- Abbs, S., Roberts, R. G., Mathew, C. G., Bentley, D. R. and Bobrow, M. (1990). Accurate assessment of intragenic recombination frequency within the Duchenne muscular dystrophy gene. *Genomics* **7**: 602-6.
- Adams, M. D., Kelley, J. M., Gocayne, J. D., Dubnick, M., Polymeropoulos, M. H., Xiao, H., *et al.* (1991). Complementary DNA sequencing: expressed sequence tags and human genome project. *Science* **252**: 1651-6.
- Adams, M. D., Kerlavage, A. R., Fleischmann, R. D., Fuldner, R. A., Bult, C. J., Lee, N. H., *et al.* (1995). Initial assessment of human gene diversity and expression patterns based upon 83 million nucleotides of cDNA sequence. *Nature* **377**: 3-174.
- Adams, M. D., Celtniker, S. E., Holt, R. A., Evans, C. A., Gocayne, J. D., Amanatides, P. G., *et al.* (2000). The genome sequence of *Drosophila melanogaster*. *Science* **287**: 2185-95.
- Ahmad, K. F., Engel, C. K. and Prive, G. G. (1998). Crystal structure of the BTB domain from PLZF. *Proc Natl Acad Sci U S A* **95**: 12123-8.
- Albertsen, H. M., Abderrahim, H., Cann, H. M., Dausset, J., Le Paslier, D. and Cohen, D. (1990). Construction and characterization of a yeast artificial chromosome library containing seven haploid human genome equivalents. *Proc Natl Acad Sci U S A* **87**: 4256-60.
- Aldridge, J., Kunkel, L., Bruns, G., Tantravahi, U., Lalande, M., Brewster, T., *et al.* (1984). A strategy to reveal high-frequency RFLPs along the human X chromosome. *Am J Hum Genet* **36**: 546-64.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* **215**: 403-10.

- Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., *et al.* (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389-402.
- Aman, M. J., Tayebi, N., Obiri, N. I., Puri, R. K., Modi, W. S. and Leonard, W. J. (1996). cDNA cloning and characterization of the human interleukin 13 receptor alpha chain. *J Biol Chem* **271**: 29265-70.
- Anand, R., Riley, J. H., Butler, R., Smith, J. C. and Markham, A. F. (1990). A 3.5 genome equivalent multi access YAC library: construction, characterisation, screening and storage. *Nucleic Acids Res* **18**: 1951-6.
- Anderson, S. (1981). Shotgun DNA sequencing using cloned DNase I-generated fragments. *Nucleic Acids Res* **9**: 3015-27.
- Ansari-Lari, M. A., Oeltjen, J. C., Schwartz, S., Zhang, Z., Muzny, D. M., Lu, J., *et al.* (1998). Comparative sequence analysis of a gene-rich cluster at human chromosome 12p13 and its syntenic region in mouse chromosome 6. *Genome Res* **8**: 29-40.
- Antequera, F. and Bird, A. (1993). Number of CpG islands and genes in human and mouse. *Proc Natl Acad Sci U S A* **90**: 11995-9.
- Arabidopsis Genome Initiative, The. (2000). Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* **408**: 796-815.
- Arpin, M., Friederich, E., Algrain, M., Vernel, F. and Louvard, D. (1994). Functional differences between L- and T-plastin isoforms. *J Cell Biol* **127**: 1995-2008.
- Au, H. C., Seo, B. B., Matsuno-Yagi, A., Yagi, T. and Scheffler, I. E. (1999). The NDUFA1 gene product (MWFE protein) is essential for activity of complex I in mammalian mitochondria. *Proc Natl Acad Sci U S A* **96**: 4354-9.
- Bailey, J. A., Carrel, L., Chakravarti, A. and Eichler, E. E. (2000). Molecular evidence for a relationship between LINE-1 elements and X chromosome inactivation: the Lyon repeat hypothesis. *Proc Natl Acad Sci U S A* **97**: 6634-9.
- Barbazuk, W. B., Korf, I., Kadavi, C., Heyen, J., Tate, S., Wun, E., *et al.* (2000). The syntenic relationship of the zebrafish and human genomes. *Genome Res* **10**: 1351-8.

- Bardwell, V. J. and Treisman, R. (1994). The POZ domain: a conserved protein-protein interaction motif. *Genes Dev* **8**: 1664-77.
- Barr, M. L. and Bertram, E. G. (1949). A morphological distinction between neurones of the male and female, and the behaviour of the nucleolar satellite during accelerated nucleoprotein synthesis. *Nature* **163**: 676-677.
- Barr, M. L. and Carr, D. H. (1961). Correlations between sex chromatin and sex chromosomes. *Acta Cytol.* **6**: 34-45.
- Bateman, A., Birney, E., Cerruti, L., Durbin, R., Etwiller, L., Eddy, S. R., et al. (2002). The Pfam protein families database. *Nucleic Acids Res* **30**: 276-80.
- Battini, R., Ferrari, S., Kaczmarek, L., Calabretta, B., Chen, S. T. and Baserga, R. (1987). Molecular cloning of a cDNA for a human ADP/ATP carrier which is growth-regulated. *J Biol Chem* **262**: 4355-9.
- Batzoglou, S., Pachter, L., Mesirov, J. P., Berger, B. and Lander, E. S. (2000). Human and mouse gene structure: comparative analysis and application to exon prediction. *Genome Res* **10**: 950-8.
- Beaudoin, E., Freier, S., Wyatt, J. R., Claverie, J. M. and Gautheret, D. (2000). Patterns of variant polyadenylation signal usage in human genes. *Genome Res* **10**: 1001-10.
- Bell, C. J., Budarf, M. L., Nieuwenhuijsen, B. W., Barnoski, B. L., Buetow, K. H., Campbell, K., et al. (1995). Integration of physical, breakpoint and genetic maps of chromosome 22. Localization of 587 yeast artificial chromosomes with 238 mapped markers. *Hum Mol Genet* **4**: 59-69.
- Bellanne-Chantelot, C., Lacroix, B., Ougen, P., Billault, A., Beaufils, S., Bertrand, S., et al. (1992). Mapping the whole human genome by fingerprinting yeast artificial chromosomes. *Cell* **70**: 1059-68.
- Belyaev, N., Keohane, A. M. and Turner, B. M. (1996). Differential underacetylation of histones H2A, H3 and H4 on the inactive X chromosome in human female cells. *Hum Genet* **97**: 573-8.

- Bentley, D. R., Deloukas, P., Dunham, A., French, L., Gregory, S. G., Humphray, S. J., *et al.* (2001). The physical maps for sequencing human chromosomes 1, 6, 9, 10, 13, 20 and X. *Nature* **409**: 942-3.
- Bernstein, H. S., Bishop, D. F., Astrin, K. H., Kornreich, R., Eng, C. M., Sakuraba, H., *et al.* (1989). Fabry disease: six gene rearrangements and an exonic point mutation in the alpha-galactosidase gene. *J Clin Invest* **83**: 1390-9.
- Bickmore, W. A. and Sumner, A. T. (1989). Mammalian chromosome banding--an expression of genome organization. *Trends Genet* **5**: 144-8.
- Blattner, F. R., Plunkett, G., 3rd, Bloch, C. A., Perna, N. T., Burland, V., Riley, M., *et al.* (1997). The complete genome sequence of Escherichia coli K-12. *Science* **277**: 1453-74.
- Boguski, M. S. (1995). Hunting for genes in computer data bases. *N Engl J Med* **333**: 645-7.
- Borsani, G., Tonlorenzi, R., Simmler, M. C., Dandolo, L., Arnaud, D., Capra, V., *et al.* (1991). Characterization of a murine gene expressed from the inactive X chromosome. *Nature* **351**: 325-9.
- Bouffard, G. G., Idol, J. R., Braden, V. V., Iyer, L. M., Cunningham, A. F., Weintraub, L. A., *et al.* (1997). A physical map of human chromosome 7: an integrated YAC contig map with average STS spacing of 79 kb. *Genome Res* **7**: 673-92.
- Boyd, Y., Blair, H. J., Cunliffe, P., Denny, P., Gormally, E. and Herman, G. E. (1998). Encyclopedia of the mouse genome VII. Mouse chromosome X. *Mamm Genome* **8**: S361-77.
- Brenner, S. (1990). The human genome: the nature of the enterprise. *Ciba Found Symp* **149**: 6-12; discussion 12-7.
- Brickner, A. G., Koop, B. F., Aronow, B. J. and Wiginton, D. A. (1999). Genomic sequence comparison of the human and mouse adenosine deaminase gene regions. *Mamm Genome* **10**: 95-101.

- Brockdorff, N., Ashworth, A., Kay, G. F., Cooper, P., Smith, S., McCabe, V. M., *et al.* (1991). Conservation of position and exclusive expression of mouse Xist from the inactive X chromosome. *Nature* **351**: 329-31.
- Brockdorff, N. (1998). The role of Xist in X-inactivation. *Curr Opin Genet Dev* **8**: 328-33.
- Bruls, T., Gyapay, G., Petit, J. L., Artiguenave, F., Vico, V., Qin, S., *et al.* (2001). A physical map of human chromosome 14. *Nature* **409**: 947-8.
- Buetow, K. H., Ludwigsen, S., Scherbier-Heddemeyer, T., Quillen, J., Murray, J. C., Sheffield, V. C., *et al.* (1994). Human genetic map. Genome maps V. Wall chart. *Science* **265**: 2055-70.
- Burge, C. and Karlin, S. (1997). Prediction of complete gene structures in human genomic DNA. *J Mol Biol* **268**: 78-94.
- Burglin, T. R. and Barnes, T. M. (1992). Introns in sequence tags. *Nature* **357**: 367-8.
- Burke, D. T., Carle, G. F. and Olson, M. V. (1987). Cloning of large segments of exogenous DNA into yeast by means of artificial chromosome vectors. *Science* **236**: 806-12.
- Cabezas, D. A., Arena, J. F., Stevenson, R. E., Schwartz, C., Goldberg, S., Morales, A., *et al.* (1999). XLMR database. *Am J Med Genet* **85**: 202-5.
- Caput, D., Laurent, P., Kaghad, M., Lelias, J. M., Lefort, S., Vita, N., *et al.* (1996). Cloning and characterization of a specific interleukin (IL)-13 binding protein structurally related to the IL-5 receptor alpha chain. *J Biol Chem* **271**: 16921-6.
- Carninci, P., Shibata, Y., Hayatsu, N., Itoh, M., Shiraki, T., Hirozane, T., *et al.* (2001). Balanced-size and long-size cloning of full-length, cap-trapped cDNAs into vectors of the novel lambda-FLC family allows enhanced gene discovery rate and functional analysis. *Genomics* **77**: 79-90.
- C. elegans Sequencing Consortium, The. (1998). Genome sequence of the nematode *C. elegans*: a platform for investigating biology. The *C. elegans* Sequencing Consortium. *Science* **282**: 2012-8.
- Chapman, V. M. (1986). X chromosome regulation in oogenesis and early mammalian development. Rossant, J. and Pedersen, E. D. Cambridge University Press.

- Chen, J. D., Mackey, D., Fuller, H., Serravalle, S., Olsson, J. and Denton, M. J. (1989). X-linked megalocornea: close linkage to DXS87 and DXS94. *Hum Genet* **83**: 292-4.
- Chumakov, I., Rigault, P., Guillou, S., Ougen, P., Billaut, A., Guasconi, G., *et al.* (1992a). Continuum of overlapping clones spanning the entire human chromosome 21q. *Nature* **359**: 380-7.
- Chumakov, I. M., Le Gall, I., Billault, A., Ougen, P., Soularue, P., Guillou, S., *et al.* (1992b). Isolation of chromosome 21-specific yeast artificial chromosomes from a total human genome library. *Nat Genet* **1**: 222-5.
- Chumakov, I. M., Rigault, P., Le Gall, I., Bellanne-Chantelot, C., Billault, A., Guillou, S., *et al.* (1995). A YAC contig map of the human genome. *Nature* **377**: 175-297.
- Clerc, P. and Avner, P. (1998). Role of the region 3' to Xist exon 6 in the counting process of X-chromosome inactivation. *Nat Genet* **19**: 249-53.
- Coffey, A. J., Roberts, R. G., Green, E. D., Cole, C. G., Butler, R., Anand, R., *et al.* (1992). Construction of a 2.6-Mb contig in yeast artificial chromosomes spanning the human dystrophin gene using an STS-based approach. *Genomics* **12**: 474-84.
- Coffey, A. J., Brooksbank, R. A., Brandau, O., Oohashi, T., Howell, G. R., Bye, J. M., *et al.* (1998). Host response to EBV infection in X-linked lymphoproliferative disease results from mutations in an SH2-domain encoding gene. *Nat Genet* **20**: 129-35.
- Cohen, D., Chumakov, I. and Weissenbach, J. (1993). A first-generation physical map of the human genome. *Nature* **366**: 698-701.
- Collins, J. and Bruning, H. J. (1978). Plasmids useable as gene-cloning vectors in an in vitro packaging by coliphage lambda: "cosmids". *Gene* **4**: 85-107.
- Collins, J. E., Cole, C. G., Smink, L. J., Garrett, C. L., Leversha, M. A., Soderlund, C. A., *et al.* (1995). A high-density YAC contig map of human chromosome 22. *Nature* **377**: 367-79.
- Coulson, A., Sulston, J., Brenner, S. and Karn, J. (1986). Towards a physical map of the genome of the nematode *C. elegans*. *Proc. Natl. Acad. Sci., USA* **83**: 7821-7825.

- Coulson, A., Waterston, R., Kiff, J., Sulston, J. and Kohara, Y. (1988). Genome linking with yeast artificial chromosomes. *Nature* **335**: 184-6.
- Crow, Y. J. and Tolmie, J. L. (1998). Recurrence risks in mental retardation. *J Med Genet* **35**: 177-82.
- Daniel, J. M., Ireton, R. C. and Reynolds, A. B. (2001). Monoclonal antibodies to Kaiso: a novel transcription factor and p120ctn-binding protein. *Hybridoma* **20**: 159-66.
- Dehal, P., Predki, P., Olsen, A. S., Kobayashi, A., Folta, P., Lucas, S., *et al.* (2001). Human chromosome 19 and related regions in mouse: conservative and lineage-specific evolution. *Science* **293**: 104-11.
- Delbruck, S., Sonneborn, A., Gerads, M., Grablowitz, A. H. and Ernst, J. F. (1997). Characterization and regulation of the genes encoding ribosomal proteins L39 and S7 of the human pathogen *Candida albicans*. *Yeast* **13**: 1199-210.
- Deloukas, P., Schuler, G. D., Gyapay, G., Beasley, E. M., Soderlund, C., Rodriguez-Tome, P., *et al.* (1998). A physical map of 30,000 human genes. *Science* **282**: 744-6.
- Deloukas, P., Matthews, L. H., Ashurst, J., Burton, J., Gilbert, J. G., Jones, M., *et al.* (2001). The DNA sequence and comparative analysis of human chromosome 20. *Nature* **414**: 865-71.
- Dib, C., Faure, S., Fizames, C., Samson, D., Drouot, N., Vignal, A., *et al.* (1996). A comprehensive genetic map of the human genome based on 5,264 microsatellites. *Nature* **380**: 152-4.
- Dietz-Band, J. N., Turco, A. E., Willard, H. F., Vincent, A., Skolnick, M. H. and Barker, D. F. (1990). Isolation, characterization, and physical localization of 33 human X-chromosome RFLP markers. *Cytogenet Cell Genet* **54**: 137-41.
- Doggett, N. A., Goodwin, L. A., Tesmer, J. G., Meincke, L. J., Bruce, D. C., Clark, L. M., *et al.* (1995). An integrated physical map of human chromosome 16. *Nature* **377**: 335-65.
- Donis-Keller, H., Green, P., Helms, C., Cartinhour, S., Weiffenbach, B., Stephens, K., *et al.* (1987). A genetic linkage map of the human genome. *Cell* **51**: 319-37.

- Drayna, D. and White, R. (1985). The genetic linkage map of the human X chromosome. *Science* **230**: 753-8.
- Driever, W., Solnica-Krezel, L., Schier, A. F., Neuhauss, S. C., Malicki, J., Stemple, D. L., *et al.* (1996). A genetic screen for mutations affecting embryogenesis in zebrafish. *Development* **123**: 37-46.
- Dubchak, I., Brudno, M., Loots, G. G., Pachter, L., Mayor, C., Rubin, E. M., *et al.* (2000). Active conservation of noncoding sequences revealed by three-way species comparisons. *Genome Res* **10**: 1304-6.
- Dunham, I., Shimizu, N., Roe, B. A., Chissoe, S., Hunt, A. R., Collins, J. E., *et al.* (1999). The DNA sequence of human chromosome 22. *Nature* **402**: 489-95.
- Duret, L. and Bucher, P. (1997). Searching for regulatory elements in human noncoding sequences. *Curr Opin Struct Biol* **7**: 399-406.
- Ekker, M., Fritz, A. and Westerfield, M. (1992). Identification of two families of satellite-like repetitive DNA sequences from the zebrafish (*Brachydanio rerio*). *Genomics* **13**: 1169-73.
- Elgar, G., Clark, M. S., Meek, S., Smith, S., Warner, S., Edwards, Y. J., *et al.* (1999). Generation and analysis of 25 Mb of genomic DNA from the pufferfish *Fugu rubripes* by sequence scanning. *Genome Res* **9**: 960-71.
- Eliceiri, G. L. (1999). Small nucleolar RNAs. *Cell Mol Life Sci* **56**: 22-31.
- Evans, M. J., Carlton, M. B. and Russ, A. P. (1997). Gene trapping and functional genomics. *Trends Genet* **13**: 370-4.
- Ewing, B. and Green, P. (2000). Analysis of expressed sequence tags indicates 35,000 human genes. *Nat Genet* **25**: 232-4.
- Feinberg, A. P. and Vogelstein, B. (1983). A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* **132**: 6-13.
- Fields, C., Adams, M. D., White, O. and Venter, J. C. (1994). How many genes in the human genome? *Nat Genet* **7**: 345-6.

- Fleischmann, R. D., Adams, M. D., White, O., Clayton, R. A., Kirkness, E. F., Kerlavage, A. R., *et al.* (1995). Whole-genome random sequencing and assembly of Haemophilus influenzae Rd. *Science* **269**: 496-512.
- Foote, S., Vollrath, D., Hilton, A. and Page, D. C. (1992). The human Y chromosome: overlapping DNA clones spanning the euchromatic region. *Science* **258**: 60-6.
- Frattini, A., Faranda, S., Bagnasco, L., Patrosso, C., Nulli, P., Zucchi, I., *et al.* (1997). Identification of a new member (ZNF183) of the Ring finger gene family in Xq24-25. *Gene* **192**: 291-8.
- Gates, M. A., Kim, L., Egan, E. S., Cardozo, T., Sirotkin, H. I., Dougan, S. T., *et al.* (1999). A genetic linkage map for zebrafish: comparative analysis and localization of genes and expressed sequences. *Genome Res* **9**: 334-47.
- Gecz, J., Barnett, S., Liu, J., Hollway, G., Donnelly, A., Eyre, H., *et al.* (1999). Characterization of the human glutamate receptor subunit 3 gene (GRIA3), a candidate for bipolar disorder and nonspecific X-linked mental retardation. *Genomics* **62**: 356-68.
- Gemmill, R. M., Chumakov, I., Scott, P., Waggoner, B., Rigault, P., Cypser, J., *et al.* (1995). A second-generation YAC contig map of human chromosome 3. *Nature* **377**: 299-319.
- Gerdes, D., Wehling, M., Leube, B. and Falkenstein, E. (1998). Cloning and tissue expression of two putative steroid membrane receptors. *Biol Chem* **379**: 907-11.
- Gilbert, W. (1992). The Codes of Codes. Kelves, D. J. and Hood, L. Harvard University Press.
- Giraud, S., Bonod-Bidaud, C., Wesolowski-Louvel, M. and Stepien, G. (1998). Expression of human ANT2 gene in highly proliferative cells: GRBOX, a new transcriptional element, is involved in the regulation of glycolytic ATP import into mitochondria. *J Mol Biol* **281**: 409-18.
- Gish, W. and States, D. J. (1993). Identification of protein coding regions by database similarity search. *Nature Genetics* **3**: 266-72.

- Goffeau, A., Barrell, B. G., Bussey, H., Davis, R. W., Dujon, B., Feldmann, H., *et al.* (1996). Life with 6000 genes. *Science* **274**: 546, 563-7.
- Goss, S. J. and Harris, H. (1975). New method for mapping genes in human chromosomes. *Nature* **255**: 680-4.
- Gottgens, B., Barton, L. M., Gilbert, J. G., Bench, A. J., Sanchez, M. J., Bahn, S., *et al.* (2000). Analysis of vertebrate SCL loci identifies conserved enhancers. *Nat Biotechnol* **18**: 181-6.
- Gottgens, B., Gilbert, J. G., Barton, L. M., Grafham, D., Rogers, J., Bentley, D. R., *et al.* (2001). Long-range comparison of human and mouse SCL loci: localized regions of sensitivity to restriction endonucleases correspond precisely with peaks of conserved noncoding sequences. *Genome Res* **11**: 87-97.
- Graves, J. A. (1996). Mammals that break the rules: genetics of marsupials and monotremes. *Annu Rev Genet* **30**: 233-60.
- Green, E. D. and Olson, M. V. (1990). Chromosomal region of the cystic fibrosis gene in yeast artificial chromosomes: a model for human genome mapping. *Science* **250**: 94-8.
- Green, P. (1997). Against a whole-genome shotgun. *Genome Res* **7**: 410-7.
- Gregg, R. G., Palmer, C., Kirkpatrick, S. and Simantel, A. (1996). Localization of a non-specific X-linked mental retardation gene, MRX23, to Xq23-q24. *Hum Mol Genet* **5**: 411-4.
- Gregory, S. G., Howell, G. R. and Bentley, D. R. (1997). Genome mapping by fluorescent fingerprinting. *Genome Res* **7**: 1162-8.
- Gubler, U. and Hoffman, B. J. (1983). A simple and very efficient method for generating cDNA libraries. *Gene* **25**: 263-9.
- Guigo, R., Agarwal, P., Abril, J. F., Burset, M. and Fickett, J. W. (2000). An assessment of gene prediction accuracy in large DNA sequences. *Genome Res* **10**: 1631-42.
- Gyapay, G., Morissette, J., Vignal, A., Dib, C., Fizames, C., Millasseau, P., *et al.* (1994). The 1993-94 Genethon human genetic linkage map. *Nat Genet* **7**: 246-339.

- Gyapay, G., Schmitt, K., Fizames, C., Jones, H., Vega-Czarny, N., Spillett, D., *et al.* (1996). A radiation hybrid map of the human genome. *Hum Mol Genet* **5**: 339-46.
- Haffter, P., Granato, M., Brand, M., Mullins, M. C., Hammerschmidt, M., Kane, D. A., *et al.* (1996). The identification of genes with unique and essential functions in the development of the zebrafish, *Danio rerio*. *Development* **123**: 1-36.
- Hannenhalli, S. and Levy, S. (2001). Promoter prediction in the human genome. *Bioinformatics* **17**: S90-6.
- Hardison, R., Xu, J., Jackson, J., Mansberger, J., Selifonova, O., Grotch, B., *et al.* (1993). Comparative analysis of the locus control region of the rabbit beta-like gene cluster: HS3 increases transient expression of an embryonic epsilon-globin gene. *Nucleic Acids Res* **21**: 1265-72.
- Hardison, R., Slightom, J. L., Gumucio, D. L., Goodman, M., Stojanovic, N. and Miller, W. (1997). Locus control regions of mammalian beta-globin gene clusters: combining phylogenetic analyses and experimental results to gain functional insights. *Gene* **205**: 73-94.
- Hardison, R. C. (2000). Conserved noncoding sequences are reliable guides to regulatory elements. *Trends Genet* **16**: 369-72.
- Hattori, M., Fujiyama, A., Taylor, T. D., Watanabe, H., Yada, T., Park, H. S., *et al.* (2000). The DNA sequence of human chromosome 21. *Nature* **405**: 311-9.
- Hillier, L. D., Lennon, G., Becker, M., Bonaldo, M. F., Chiapelli, B., Chissoe, S., *et al.* (1996). Generation and analysis of 280,000 human expressed sequence tags. *Genome Res* **6**: 807-28.
- Holland, J., Coffey, A. J., Giannelli, F. and Bentley, D. R. (1993). Vertical integration of cosmid and YAC resources for interval mapping on the X-chromosome. *Genomics* **15**: 297-304.
- Huang, S.-H., Yang, A. Y. and Holcnenberg, J. (1993). Amplification of gene ends from gene libraries by polymerase chain reaction with single-sided specificity. White, B. A. Humana Press.

- Hudson, L. D., Friedrich, V. L., Jr., Behar, T., Dubois-Dalcq, M. and Lazzarini, R. A. (1989). The initial events in myelin synthesis: orientation of proteolipid protein in the plasma membrane of cultured oligodendrocytes. *J Cell Biol* **109**: 717-27.
- Hudson, T. J., Stein, L. D., Gerety, S. S., Ma, J., Castle, A. B., Silva, J., et al. (1995). An STS-based map of the human genome. *Science* **270**: 1945-54.
- Hudson, T. J., Church, D. M., Greenaway, S., Nguyen, H., Cook, A., Steen, R. G., et al. (2001). A radiation hybrid map of mouse genes. *Nat Genet* **29**: 201-5.
- IHGSC (International Human Genome Sequencing Consortium). (2001). Initial sequencing and analysis of the human genome. *Nature* **409**: 860-921.
- Imai, T. and Olson, M. V. (1990). Second-generation approach to the construction of yeast artificial-chromosome libraries. *Genomics* **8**: 297-303.
- Inoue, K., Osaka, H., Imaizumi, K., Nezu, A., Takanashi, J., Arii, J., et al. (1999). Proteolipid protein gene duplications causing Pelizaeus-Merzbacher disease: molecular mechanism and phenotypic manifestations. *Ann Neurol* **45**: 624-32.
- Ioannou, P. A., Amemiya, C. T., Garnes, J., Kroisel, P. M., Shizuya, H., Chen, C., et al. (1994). A new bacteriophage P1-derived vector for the propagation of large human DNA fragments. *Nat Genet* **6**: 84-9.
- Izsvák, Z., Ivics, Z. and Hackett, P. B. (1995). Characterization of a Tc1-like transposable element in zebrafish (*Danio rerio*). *Mol Gen Genet* **247**: 312-22.
- Izsvák, Z., Ivics, Z., Garcia-Estefanía, D., Fahrenkrug, S. C. and Hackett, P. B. (1996). DANA elements: a family of composite, tRNA-derived short interspersed DNA elements associated with mutational activities in zebrafish (*Danio rerio*). *Proc Natl Acad Sci U S A* **93**: 1077-81.
- Izsvák, Z., Ivics, Z., Shimoda, N., Mohn, D., Okamoto, H. and Hackett, P. B. (1999). Short inverted-repeat transposable elements in teleost fish and implications for a mechanism of their amplification. *J Mol Evol* **48**: 13-21.
- Jackson, M. S. and Strachan, T. (1996). Human Genome Evolution. Oxford Bios.

- Jang, W., Hua, A., Spilson, S. V., Miller, W., Roe, B. A. and Meisler, M. H. (1999). Comparative sequence of human and mouse BAC clones from the mnd2 region of chromosome 2p13. *Genome Res* **9**: 53-61.
- Jeppesen, P. and Turner, B. M. (1993). The inactive X chromosome in female mammals is distinguished by a lack of histone H4 acetylation, a cytogenetic marker for gene expression. *Cell* **74**: 281-9.
- Justice, M. J. (2000). Capitalizing on large-scale mouse mutagenesis screens. *Nat Rev Genet* **1**: 109-15.
- Kannan, K., Stewart, R. M., Bounds, W., Carlsson, S. R., Fukuda, M., Betzing, K. W., *et al.* (1996). Lysosome-associated membrane proteins h-LAMP1 (CD107a) and h-LAMP2 (CD107b) are activation-dependent cell surface glycoproteins in human peripheral blood mononuclear cells which mediate cell adhesion to vascular endothelium. *Cell Immunol* **171**: 10-9.
- Kaplan, J. C., Kahn, A. and Chelly, J. (1992). Illegitimate transcription: its use in the study of inherited disease. *Hum Mutat* **1**: 357-60.
- Kendall, E., Evans, W., Jin, H., Holland, J. and Vetrie, D. (1997). A complete YAC contig and cosmid interval map covering the entirety of human Xq21.33 to Xq22.3 from DXS3 to DXS287. *Genomics* **43**: 171-82.
- Khan, A. S., Wilcox, A. S., Hopkins, J. A. and Sikela, J. M. (1991). Efficient double stranded sequencing of cDNA clones containing long poly(A) tails using anchored poly(dT) primers. *Nucleic Acids Res* **19**: 1715.
- Khan, A. S., Wilcox, A. S., Polymeropoulos, M. H., Hopkins, J. A., Stevens, T. J., Robinson, M., *et al.* (1992). Single pass sequencing and physical and genetic mapping of human brain cDNAs. *Nat Genet* **2**: 180-5.
- Kim, U. J., Shizuya, H., de Jong, P. J., Birren, B. and Simon, M. I. (1992). Stable propagation of cosmid sized human DNA inserts in an F factor based vector. *Nucleic Acids Res* **20**: 1083-5.

- Kinoshita, M., Kumar, S., Mizoguchi, A., Ide, C., Kinoshita, A., Haraguchi, T., *et al.* (1997). Nedd5, a mammalian septin, is a novel cytoskeletal component interacting with actin-based structures. *Genes Dev* **11**: 1535-47.
- Kohara, Y., Akiyama, K. and Isono, K. (1987). The physical map of the whole *E. coli* chromosome: application of a new strategy for rapid analysis and sorting of a large genomic library. *Cell* **50**: 495-508.
- Koken, M. H., Hoogerbrugge, J. W., Jasper-Dekker, I., de Wit, J., Willemse, R., Roest, H. P., *et al.* (1996). Expression of the ubiquitin-conjugating DNA repair enzymes HHR6A and B suggests a role in spermatogenesis and chromatin modification. *Dev Biol* **173**: 119-32.
- Koller, B. H. and Smithies, O. (1992). Altering genes in animals by gene targeting. *Annu Rev Immunol* **10**: 705-30.
- Koop, B. F. and Hood, L. (1994). Striking sequence similarity over almost 100 kilobases of human and mouse T-cell receptor DNA. *Nat Genet* **7**: 48-53.
- Kozak, M. (1991). Structural features in eukaryotic mRNAs that modulate the initiation of translation. *J Biol Chem* **266**: 19867-70.
- Krauter, K., Montgomery, K., Yoon, S. J., LeBlanc-Straceski, J., Renault, B., Marondel, I., *et al.* (1995). A second-generation YAC contig map of human chromosome 12. *Nature* **377**: 321-33.
- Lapenta, V., Chiurazzi, P., van der Spek, P., Pizzuti, A., Hanaoka, F. and Brahe, C. (1997). SMT3A, a human homologue of the *S. cerevisiae* SMT3 gene, maps to chromosome 21qter and defines a novel gene family. *Genomics* **40**: 362-6.
- Larin, Z., Monaco, A. P. and Lehrach, H. (1991). Yeast artificial chromosome libraries containing large inserts from mouse and human DNA. *Proc Natl Acad Sci U S A* **88**: 4123-7.
- Lehrke, R. G. (1974). X-linked mental retardation and verbal disability. *Birth Defects Orig Artic Ser* **10**: 1-100.

- Levine, A. and Durbin, R. (2001). A computational scan for U12-dependent introns in the human genome sequence. *Nucleic Acids Res* **29**: 4006-13.
- Liang, F., Holt, I., Pertea, G., Karamycheva, S., Salzberg, S. L. and Quackenbush, J. (2000). Gene index analysis of the human genome estimates approximately 120,000 genes. *Nat Genet* **25**: 239-40.
- Lin, C. S., Park, T., Chen, Z. P. and Leavitt, J. (1993). Human plastin genes. Comparative gene structure, chromosome location, and differential expression in normal and neoplastic cells. *J Biol Chem* **268**: 2781-92.
- Little, P. F., Flavell, R. A., Kooter, J. M., Annison, G. and Williamson, R. (1979). Structure of the human fetal globin gene locus. *Nature* **278**: 227-31.
- Luciakova, K., Hodny, Z., Barath, P. and Nelson, B. D. (2000). In vivo mapping of the human adenine nucleotide translocator-2 (ANT2) promoter provides support for regulation by a pair of proximal Sp1-activating sites and an upstream silencer element. *Biochem J* **352**: 519-23.
- Lyon, M. F. (1961). Gene action in the X chromosome of the mouse. *Nature* **190**: 372-373.
- Lyon, M. F. (1998). X-chromosome inactivation: a repeat hypothesis. *Cytogenet Cell Genet* **80**: 133-7.
- Mackey, D. A., Buttery, R. G., Wise, G. M. and Denton, M. J. (1991). Description of X-linked megalocornea with identification of the gene locus. *Arch Ophthalmol* **109**: 829-33.
- Maestrini, E., Rivella, S., Tribioli, C., Purtilo, D., Rocchi, M., Archidiacono, N., et al. (1990). Probes for CpG islands on the distal long arm of the human X chromosome are clustered in Xq24 and Xq28. *Genomics* **8**: 664-70.
- Makalowski, W., Zhang, J. and Boguski, M. S. (1996). Comparative analysis of 1196 orthologous mouse and human full-length mRNA and protein sequences. *Genome Res* **6**: 846-57.
- Manoni, M., Tribioli, C., Lazzari, B., DeBellis, G., Patrosso, C., Pergolizzi, R., et al. (1991). The nucleotide sequence of a CpG island demonstrates the presence of the first exon

- of the gene encoding the human lysosomal membrane protein lamp2 and assigns the gene to Xq24. *Genomics* **9**: 551-4.
- Mathias, N., Johnson, S. L., Winey, M., Adams, A. E., Goetsch, L., Pringle, J. R., *et al.* (1996). Cdc53p acts in concert with Cdc4p and Cdc34p to control the G1-to-S-phase transition and identifies a conserved family of proteins. *Mol Cell Biol* **16**: 6634-43.
- McKusick, V. A. (1998). Mendelian Inheritance in Man. Catalogs of Human Genes and Genetic Disorders. John Hopkins University Press.
- Mei, X., Singh, I. S., Erlichman, J. and Orr, G. A. (1997). Cloning and characterization of a testis-specific, developmentally regulated A-kinase-anchoring protein (TAKAP-80) present on the fibrous sheath of rat sperm. *Eur J Biochem* **246**: 425-32.
- Metscher, B. D. and Ahlberg, P. E. (1999). Zebrafish in context: uses of a laboratory model in comparative studies. *Dev Biol* **210**: 1-14.
- Miller, W. (2001). Comparison of genomic DNA sequences: solved and unsolved problems. *Bioinformatics* **17**: 391-7.
- Mills, A. A., Zheng, B., Wang, X. J., Vogel, H., Roop, D. R. and Bradley, A. (1999). p63 is a p53 homologue required for limb and epidermal morphogenesis. *Nature* **398**: 708-13.
- Monaco, A. P. and Larin, Z. (1994). YACs, BACs, PACs and MACs: artificial chromosomes as research tools. *Trends Biotechnol* **12**: 280-6.
- Murray, J. C., Buetow, K. H., Weber, J. L., Ludwigsen, S., Scherbier-Heddema, T., Manion, F., *et al.* (1994). A comprehensive human linkage map with centimorgan density. Cooperative Human Linkage Center (CHLC). *Science* **265**: 2049-54.
- Nagaraja, R., MacMillan, S., Kere, J., Jones, C., Griffin, S., Schmatz, M., *et al.* (1997). X chromosome map at 75-kb STS resolution, revealing extremes of recombination and GC content. *Genome Res* **7**: 210-22.

- Nagaraja, R., MacMillan, S., Jones, C., Masisi, M., Pengue, G., Porta, G., *et al.* (1998). Integrated YAC/STS physical and genetic map of 22.5 Mb of human Xq24-q26 at 56-kb inter-STS resolution. *Genomics* **52**: 247-66.
- Nakamura, Y., Julier, C., Wolff, R., Holm, T., O'Connell, P., Leppert, M., *et al.* (1987). Characterization of a human 'midisatellite' sequence. *Nucleic Acids Res* **15**: 2537-47.
- Naylor, J. A., Buck, D., Green, P., Williamson, H., Bentley, D. and Giannelli, F. (1995). Investigation of the factor VIII intron 22 repeated region (int22h) and the associated inversion junctions. *Hum Mol Genet* **4**: 1217-24.
- Neri, G. and Chiurazzi, P. (1999). X-linked mental retardation. *Adv Genet* **41**: 55-94.
- Nizetic, D., Gellen, L., Hamvas, R. M., Mott, R., Grigoriev, A., Vatcheva, R., *et al.* (1994). An integrated YAC-overlap and 'cosmid-pocket' map of the human chromosome 21. *Hum Mol Genet* **3**: 759-70.
- Nourbakhsh, M., Oumard, A., Schwarzer, M. and Hauser, H. (2000). NRF, a nuclear inhibitor of NF-kappaB proteins silencing interferon-beta promoter. *Eur Cytokine Netw* **11**: 500-1.
- O'Brien, S. J., Eisenberg, J. F., Miyamoto, M., Hedges, S. B., Kumar, S., Wilson, D. E., *et al.* (1999). Genome maps 10. Comparative genomics. Mammalian radiations. Wall chart. *Science* **286**: 463-78.
- Ohno, S. (1967). Sex chromosomes and sex linked genes. Springer.
- Olson, M. V., Dutchik, J. E., Graham, M. Y., Brodeur, G. M., Helms, C., Frank, M., *et al.* (1986). Random-clone strategy for genomic restriction mapping in yeast. *Proc Natl Acad Sci U S A* **83**: 7826-30.
- Osaka, F., Kawasaki, H., Aida, N., Saeki, M., Chiba, T., Kawashima, S., *et al.* (1998). A new NEDD8-ligating system for cullin-4A. *Genes Dev* **12**: 2263-8.
- Panning, B. and Jaenisch, R. (1998). RNA and the epigenetic regulation of X chromosome inactivation. *Cell* **93**: 305-8.
- Papadakis, M. N. and Patrinos, G. P. (1999). Contribution of gene conversion in the evolution of the human beta-like globin gene family. *Hum Genet* **104**: 117-25.

- Pestov, N. B., Adams, G., Shakhparyonov, M. I. and Modyanov, N. N. (1999). Identification of a novel gene of the X,K-ATPase beta-subunit family that is predominantly expressed in skeletal and heart muscles. *FEBS Lett* **456**: 243-8.
- Postlethwait, J. H., Yan, Y. L., Gates, M. A., Horne, S., Amores, A., Brownlie, A., *et al.* (1998). Vertebrate genome evolution and the zebrafish gene map. *Nat Genet* **18**: 345-9.
- Rabbitts, T. H. (1976). Bacterial cloning of plasmids carrying copies of rabbit globin messenger RNA. *Nature* **260**: 221-5.
- Rastan, S. (1983). Non-random X-chromosome inactivation in mouse X-autosome translocation embryos--location of the inactivation centre. *J Embryol Exp Morphol* **78**: 1-22.
- Rastan, S. (1994). X chromosome inactivation and the Xist gene. *Curr Opin Genet Dev* **4**: 292-7.
- Richards, R. I., Holman, K., Kozman, H., Kremer, E., Lynch, M., Pritchard, M., *et al.* (1991). Fragile X syndrome: genetic localisation by linkage mapping of two microsatellite repeats FRAXAC1 and FRAXAC2 which immediately flank the fragile site. *J Med Genet* **28**: 818-23.
- Riley, J., Butler, R., Ogilvie, D., Finniear, R., Jenner, D., Powell, S., *et al.* (1990). A novel, rapid method for the isolation of terminal sequences from yeast artificial chromosome (YAC) clones. *Nucleic Acids Res* **18**: 2887-90.
- Roberts, R. G., Barby, T. F., Manners, E., Bobrow, M. and Bentley, D. R. (1991). Direct detection of dystrophin gene rearrangements by analysis of dystrophin mRNA in peripheral blood lymphocytes. *Am J Hum Genet* **49**: 298-310.
- Roest Crollius, H., Jaillon, O., Bernot, A., Dasilva, C., Bouneau, L., Fischer, C., *et al.* (2000). Estimate of human gene number provided by genome-wide analysis using Tetraodon nigroviridis DNA sequence. *Nat Genet* **25**: 235-8.

- Ross, M. and Langford, C. F. (1997). The use of flow-sorted chromosomes in genome mapping. In *Genome Mapping: a practical approach*. Dear, P. H. IRL Press at Oxford University Press. 165-184.
- Rubin, E. M. (2001). Comparing Species. *Nature* **409**: 820-821.
- Rubin, G. M., Yandell, M. D., Wortman, J. R., Gabor Miklos, G. L., Nelson, C. R., Hariharan, I. K., et al. (2000). Comparative genomics of the eukaryotes. *Science* **287**: 2204-15.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989). Molecular cloning; a laboratory manual. Cold Spring Harbor Laboratory Press.
- Sanchez, R. and Sali, A. (1997). Evaluation of comparative protein structure modeling by MODELLER-3. *Proteins Suppl:* 50-8.
- Sanger, F., Air, G. M., Barrell, B. G., Brown, N. L., Coulson, A. R., Fiddes, C. A., et al. (1977a). Nucleotide sequence of bacteriophage phi X174 DNA. *Nature* **265**: 687-95.
- Sanger, F., Nicklen, S. and Coulson, A. R. (1977b). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**: 5463-7.
- Sanger, F., Coulson, A. R., Friedmann, T., Air, G. M., Barrell, B. G., Brown, N. L., et al. (1978). The nucleotide sequence of bacteriophage phiX174. *J Mol Biol* **125**: 225-46.
- Sanger, F., Coulson, A. R., Hong, G. F., Hill, D. F. and Petersen, G. B. (1982). Nucleotide sequence of bacteriophage lambda DNA. *J Mol Biol* **162**: 729-73.
- Scherf, M., Klingenhoff, A. and Werner, T. (2000). Highly specific localization of promoter regions in large genomic sequences by PromoterInspector: a novel context analysis approach. *J Mol Biol* **297**: 599-606.
- Schiebel, K., Mertz, A., Winkelmann, M., Nagaraja, R. and Rappold, G. (1994). Localization of the adenine nucleotide translocase gene ANT2 to chromosome Xq24-q25 with tight linkage to DXS425. *Genomics* **24**: 605-6.
- Schnedl, W., Mikelsaar, A. V., Breitenbach, M. and Dann, O. (1977). DIPI and DAPI: fluorescence banding with only negligible fading. *Hum Genet* **36**: 167-72.

- Schuler, G. D., Boguski, M. S., Stewart, E. A., Stein, L. D., Gyapay, G., Rice, K., *et al.* (1996). A gene map of the human genome. *Science* **274**: 540-6.
- Schwartz, S., Zhang, Z., Frazer, K. A., Smit, A., Riemer, C., Bouck, J., *et al.* (2000). PipMaker--a web server for aligning two genomic DNA sequences. *Genome Res* **10**: 577-86.
- Shashidharan, P., Michaelidis, T. M., Robakis, N. K., Kresovali, A., Papamatheakis, J. and Plaitakis, A. (1994). Novel human glutamate dehydrogenase expressed in neural and testicular tissues and encoded by an X-linked intronless gene. *J Biol Chem* **269**: 16971-6.
- Shizuya, H., Birren, B., Kim, U. J., Mancino, V., Slepak, T., Tachiiri, Y., *et al.* (1992). Cloning and stable maintenance of 300-kilobase-pair fragments of human DNA in *Escherichia coli* using an F-factor-based vector. *Proc Natl Acad Sci U S A* **89**: 8794-7.
- Simmons, D. L. 1993. Cloning cell surface molecules by transient expression in mammalian cells. IRL Press at Oxford University PressOxford. 93-127
- Sloan, J. L. and Mager, S. (1999). Cloning and functional expression of a human Na(+) and Cl(-)-dependent neutral and cationic amino acid transporter B(0+). *J Biol Chem* **274**: 23740-5.
- Solovyev, V. V., Salamov, A. A. and Lawrence, C. B. (1995). Identification of human gene structure using linear discriminant functions and dynamic programming. *Proc Int Conf Intell Syst Mol Biol* **3**: 367-75.
- Sonnhammer, E. L. and Durbin, R. (1995). A dot-matrix program with dynamic threshold control suited for genomic DNA and protein sequence analysis. *Gene* **167**: GC1-10.
- Srivastava, A. K., McMillan, S., Jermak, C., Shomaker, M., Copeland-Yates, S. A., Sossey-Alaoui, K., *et al.* (1999). Integrated STS/YAC physical, genetic, and transcript map of human Xq21.3 to q23/q24 (DXS1203-DXS1059). *Genomics* **58**: 188-201.
- Staden, R. (1984). Graphic methods to determine the function of nucleic acid sequences. *Nucleic Acids Res* **12**: 521-38.

- Steingruber, H. E., Dunham, A., Coffey, A. J., Clegg, S. M., Howell, G. R., Maslen, G. L., *et al.* (1999). High-resolution landmark framework for the sequence-ready mapping of Xq23-q26.1. *Genome Res* **9**: 751-62.
- Stewart, E. A., McKusick, K. B., Aggarwal, A., Bajorek, E., Brady, S., Chu, A., *et al.* (1997). An STS-based radiation hybrid map of the human genome. *Genome Res* **7**: 422-33.
- Takagi, N. (1974). Differentiation of X chromosomes in early female mouse embryos. *Exp Cell Res* **86**: 127-35.
- Thanaraj, T. A. and Clark, F. (2001). Human GC-AG alternative intron isoforms with weak donor sites show enhanced consensus at acceptor exon positions. *Nucleic Acids Res* **29**: 2581-93.
- Toniolo, D. and D'Adamo, P. (2000). X-linked non-specific mental retardation. *Curr Opin Genet Dev* **10**: 280-5.
- Touchman, J. W., Dehejia, A., Chiba-Falek, O., Cabin, D. E., Schwartz, J. R., Orrison, B. M., *et al.* (2001). Human and mouse alpha-synuclein genes: comparative genomic sequence analysis and identification of a novel gene regulatory element. *Genome Res* **11**: 78-86.
- Trofatter, J. A., Dlouhy, S. R., DeMyer, W., Conneally, P. M. and Hodes, M. E. (1989). Pelizaeus-Merzbacher disease: tight linkage to proteolipid protein gene exon variant. *Proc Natl Acad Sci U S A* **86**: 9427-30.
- Trofatter, J. A., Long, K. R., Murrell, J. R., Stotler, C. J., Gusella, J. F. and Buckler, A. J. (1995). An expression-independent catalog of genes from human chromosome 22. *Genome Res* **5**: 214-24.
- Tsukada, S., Saffran, D. C., Rawlings, D. J., Parolini, O., Allen, R. C., Klisak, I., *et al.* (1993). Deficient expression of a B cell cytoplasmic tyrosine kinase in human X-linked agammaglobulinemia. *Cell* **72**: 279-90.
- Tyson, J., Bellman, S., Newton, V., Simpson, P., Malcolm, S., Pembrey, M. E., *et al.* (1996). Mapping of DFN2 to Xq22. *Hum Mol Genet* **5**: 2055-60.

- Uberbacher, E. C. and Mural, R. J. (1991). Locating protein-coding regions in human DNA sequences by a multiple sensor-neural network approach. *Proc Natl Acad Sci U S A* **88:** 11261-5.
- Venter, J. C., Adams, M. D., Sutton, G. G., Kerlavage, A. R., Smith, H. O. and Hunkapiller, M. (1998). Shotgun sequencing of the human genome. *Science* **280:** 1540-2.
- Venter, J. C., Adams, M. D., Myers, E. W., Li, P. W., Mural, R. J., Sutton, G. G., et al. (2001). The sequence of the human genome. *Science* **291:** 1304-51.
- Vetrie, D. (1993). Isolation of the defective gene in X linked agammaglobulinaemia. *J Med Genet* **30:** 452-3.
- Vetrie, D., Kendall, E., Coffey, A., Hassock, S., Collins, J., Todd, C., et al. (1994). A 6.5-Mb yeast artificial chromosome contig incorporating 33 DNA markers on the human X chromosome at Xq22. *Genomics* **19:** 42-7.
- Vogel, A. M. and Gerster, T. (1999). Promoter activity of the zebrafish bhikhari retroelement requires an intact activin signaling pathway. *Mech Dev* **85:** 133-46.
- Walhout, A. J., Boulton, S. J. and Vidal, M. (2000). Yeast two-hybrid systems and protein interaction mapping projects for yeast and worm. *Yeast* **17:** 88-94.
- Way, M., Sanders, M., Chafel, M., Tu, Y. H., Knight, A. and Matsudaira, P. (1995). beta-Scruin, a homologue of the actin crosslinking protein scruin, is localized to the acrosomal vesicle of Limulus sperm. *J Cell Sci* **108:** 3155-62.
- Weber, J. L. and Myers, E. W. (1997). Human whole-genome shotgun sequencing. *Genome Res* **7:** 401-9.
- Weissenbach, J., Gyapay, G., Dib, C., Vignal, A., Morissette, J., Millasseau, P., et al. (1992). A second-generation linkage map of the human genome. *Nature* **359:** 794-801.
- Willard, H. F. (1996). X chromosome inactivation, XIST, and pursuit of the X-inactivation center. *Cell* **86:** 5-7.
- Wingender, E., Chen, X., Hehl, R., Karas, H., Liebich, I., Matys, V., et al. (2000). TRANSFAC: an integrated system for gene expression regulation. *Nucleic Acids Res* **28:** 316-9.

- Wyman, A. R. and White, R. (1980). A highly polymorphic locus in human DNA. *Proc Natl Acad Sci U S A* **77**: 6754-8.
- Zhang, M. Q. (1997). Identification of protein coding regions in the human genome by quadratic discriminant analysis. *Proc Natl Acad Sci U S A* **94**: 565-8.
- Zhang, P., Li, M. Z. and Elledge, S. J. (2002). Towards genetic genome projects: genomic library screening and gene-targeting vector construction in a single step. *Nat Genet* **30**: 31-9.
- Zollman, S., Godt, D., Prive, G. G., Couderc, J. L. and Laski, F. A. (1994). The BTB domain, found primarily in zinc finger proteins, defines an evolutionarily conserved family that includes several developmentally regulated genes in *Drosophila*. *Proc Natl Acad Sci U S A* **91**: 10717-21.