

References

- (2004). The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science* (New York, NY 306, 636-640.
- (2004). Finishing the euchromatic sequence of the human genome. *Nature* 431, 931-945.
- (2007). Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 447, 799-816.
- Acevedo, L.G., Iniguez, A.L., Holster, H.L., Zhang, X., Green, R., and Farnham, P.J. (2007). Genome-scale ChIP-chip analysis using 10,000 human cells. *BioTechniques* 43, 791-797.
- Agulnick, A.D., Taira, M., Breen, J.J., Tanaka, T., Dawid, I.B., and Westphal, H. (1996). Interactions of the LIM-domain-binding factor *Ldb1* with LIM homeodomain proteins. *Nature* 384, 270-272.
- Akashi, K., Traver, D., Miyamoto, T., and Weissman, I.L. (2000). A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature* 404, 193-197.
- Akiyama, Y. (1998). TFSEARCH: Searching Transcription Factor Binding Sites. <http://www.rwcp.or.jp/papia/>.
- Allfrey, V.G., Faulkner, R., and Mirsky, A.E. (1964). Acetylation And Methylation Of Histones And Their Possible Role In The Regulation Of Rna Synthesis. *Proceedings of the National Academy of Sciences of the United States of America* 51, 786-794.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990). Basic local alignment search tool. *Journal of molecular biology* 215, 403-410.
- Alwine, J.C., Kemp, D.J., and Stark, G.R. (1977). Method for detection of specific RNAs in agarose gels by transfer to diazobenzyloxymethyl-paper and hybridization with DNA probes. *Proceedings of the National Academy of Sciences of the United States of America* 74, 5350-5354.
- Anderson, K.P., Crable, S.C., and Lingrel, J.B. (1998). Multiple proteins binding to a GATA-E box-GATA motif regulate the erythroid Kruppel-like factor (EKLF) gene. *The Journal of biological chemistry* 273, 14347-14354.
- Anderson, K.P., Crable, S.C., and Lingrel, J.B. (2000). The GATA-E box-GATA motif in the EKLF promoter is required for in vivo expression. *Blood* 95, 1652-1655.
- Anguita, E., Hughes, J., Heyworth, C., Blobel, G.A., Wood, W.G., and Higgs, D.R. (2004). Globin gene activation during haemopoiesis is driven by protein complexes nucleated by GATA-1 and GATA-2. *Embo J* 23, 2841-2852.
- Antes, T.J., Namciu, S.J., Fournier, R.E., and Levy-Wilson, B. (2001). The 5' boundary of the human apolipoprotein B chromatin domain in intestinal cells. *Biochemistry* 40, 6731-6742.
- Aplan, P.D., Begley, C.G., Bertness, V., Nussmeier, M., Ezquerra, A., Coligan, J., and Kirsch, I.R. (1990). The SCL gene is formed from a transcriptionally complex locus. *Molecular and cellular biology* 10, 6426-6435.
- Aplan, P.D., Lombardi, D.P., Ginsberg, A.M., Cossman, J., Bertness, V.L., and Kirsch, I.R. (1990). Disruption of the human SCL locus by "illegitimate" V-(D)-J recombinase activity. *Science* (New York, NY 250, 1426-1429.
- Attema, J.L., Papathanasiou, P., Forsberg, E.C., Xu, J., Smale, S.T., and Weissman, I.L. (2007). Epigenetic characterization of hematopoietic stem cell differentiation using miniChIP and bisulfite sequencing analysis. *Proceedings of the National Academy of Sciences of the United States of America* 104, 12371-12376.
- Aza-Blanc, P., Cooper, C.L., Wagner, K., Batalov, S., Deveraux, Q.L., and Cooke, M.P. (2003). Identification of modulators of TRAIL-induced apoptosis via RNAi-based phenotypic screening. *Mol Cell* 12, 627-637.
- Baeuerle, P.A., and Baltimore, D. (1988). I kappa B: a specific inhibitor of the NF-kappa B transcription factor. *Science* (New York, NY 242, 540-546.
- Bailey, T.L., and Elkan, C. (1995). The value of prior knowledge in discovering motifs with MEME. *Proceedings / International Conference on Intelligent Systems for Molecular Biology ; ISMB* 3, 21-29.
- Bain, G., Maandag, E.C., Izon, D.J., Amsen, D., Kruisbeek, A.M., Weintraub, B.C., Krop, I., Schlissel, M.S., Feeney, A.J., van Roon, M., et al. (1994). E2A proteins are required for proper B cell development and initiation of immunoglobulin gene rearrangements. *Cell* 79, 885-892.

- Bajic, V.B., Tan, S.L., Suzuki, Y., and Sugano, S. (2004). Promoter prediction analysis on the whole human genome. *Nature biotechnology* 22, 1467-1473.
- Balduini, C.L., Pecci, A., Loffredo, G., Izzo, P., Noris, P., Grosso, M., Bergamaschi, G., Rosti, V., Magrini, U., Ceresa, I.F., et al. (2004). Effects of the R216Q mutation of GATA-1 on erythropoiesis and megakaryocytopoiesis. *Thrombosis and haemostasis* 91, 129-140.
- Balhoff, J.P., and Wray, G.A. (2005). Evolutionary analysis of the well characterized endo16 promoter reveals substantial variation within functional sites. *Proceedings of the National Academy of Sciences of the United States of America* 102, 8591-8596.
- Banerji, J., Olson, L., and Schaffner, W. (1983). A lymphocyte-specific cellular enhancer is located downstream of the joining region in immunoglobulin heavy chain genes. *Cell* 33, 729-740.
- Banerji, J., Rusconi, S., and Schaffner, W. (1981). Expression of a beta-globin gene is enhanced by remote SV40 DNA sequences. *Cell* 27, 299-308.
- Barndt, R.J., Dai, M., and Zhuang, Y. (2000). Functions of E2A-HEB heterodimers in T-cell development revealed by a dominant negative mutation of HEB. *Molecular and cellular biology* 20, 6677-6685.
- Barski, A., Cuddapah, S., Cui, K., Roh, T.Y., Schones, D.E., Wang, Z., Wei, G., Chepelev, I., and Zhao, K. (2007). High-resolution profiling of histone methylations in the human genome. *Cell* 129, 823-837.
- Barski, A., and Frenkel, B. (2004). ChIP Display: novel method for identification of genomic targets of transcription factors. *Nucleic acids research* 32, e104.
- Barton, G.M., and Medzhitov, R. (2002). Retroviral delivery of small interfering RNA into primary cells. *Proceedings of the National Academy of Sciences of the United States of America* 99, 14943-14945.
- Basu, P., Morris, P.E., Haar, J.L., Wani, M.A., Lingrel, J.B., Gaensler, K.M., and Lloyd, J.A. (2005). KLF2 is essential for primitive erythropoiesis and regulates the human and murine embryonic beta-like globin genes in vivo. *Blood* 106, 2566-2571.
- Becker, D.M., Fikes, J.D., and Guarente, L. (1991). A cDNA encoding a human CCAAT-binding protein cloned by functional complementation in yeast. *Proceedings of the National Academy of Sciences of the United States of America* 88, 1968-1972.
- Begley, C.G., Aplan, P.D., Davey, M.P., Nakahara, K., Tchorz, K., Kurtzberg, J., Hershfield, M.S., Haynes, B.F., Cohen, D.I., Waldmann, T.A., et al. (1989). Chromosomal translocation in a human leukemic stem-cell line disrupts the T-cell antigen receptor delta-chain diversity region and results in a previously unreported fusion transcript. *Proceedings of the National Academy of Sciences of the United States of America* 86, 2031-2035.
- Begley, C.G., Aplan, P.D., Denning, S.M., Haynes, B.F., Waldmann, T.A., and Kirsch, I.R. (1989). The gene SCL is expressed during early hematopoiesis and encodes a differentiation-related DNA-binding motif. *Proceedings of the National Academy of Sciences of the United States of America* 86, 10128-10132.
- Begley, C.G., Visvader, J., Green, A.R., Aplan, P.D., Metcalf, D., Kirsch, I.R., and Gough, N.M. (1991). Molecular cloning and chromosomal localization of the murine homolog of the human helix-loop-helix gene SCL. *Proceedings of the National Academy of Sciences of the United States of America* 88, 869-873.
- Bell, A.C., West, A.G., and Felsenfeld, G. (1999). The protein CTCF is required for the enhancer blocking activity of vertebrate insulators. *Cell* 98, 387-396.
- Bennett, S. (2004). Solexa Ltd. *Pharmacogenomics* 5, 433-438.
- Bernard, O., Azogui, O., Lecointe, N., Mugneret, F., Berger, R., Larsen, C.J., and Mathieu-Mahul, D. (1992). A third tal-1 promoter is specifically used in human T cell leukemias. *The Journal of experimental medicine* 176, 919-925.
- Bernard, O., Lecointe, N., Jonveaux, P., Souyri, M., Mauchauffe, M., Berger, R., Larsen, C.J., and Mathieu-Mahul, D. (1991). Two site-specific deletions and t(1;14) translocation restricted to human T-cell acute leukemias disrupt the 5' part of the tal-1 gene. *Oncogene* 6, 1477-1488.
- Berns, K., Hijmans, E.M., Mullenders, J., Brummelkamp, T.R., Velds, A., Heimerikx, M., Kerkhoven, R.M., Madiredjo, M., Nijkamp, W., Weigelt, B., et al. (2004). A large-scale RNAi screen in human cells identifies new components of the p53 pathway. *Nature* 428, 431-437.
- Bernstein, B.E., Humphrey, E.L., Erlich, R.L., Schneider, R., Bouman, P., Liu, J.S., Kouzarides, T., and Schreiber, S.L. (2002). Methylation of histone H3 Lys 4 in coding regions of active genes. *Proceedings of the National Academy of Sciences of the United States of America* 99, 8695-8700.

- Bernstein, B.E., Kamal, M., Lindblad-Toh, K., Bekiranov, S., Bailey, D.K., Huebert, D.J., McMahon, S., Karlsson, E.K., Kulbokas, E.J., 3rd, Gingeras, T.R., et al. (2005). Genomic maps and comparative analysis of histone modifications in human and mouse. *Cell* 120, 169-181.
- Birmingham, A., Anderson, E.M., Reynolds, A., Ilsley-Tyree, D., Leake, D., Fedorov, Y., Baskerville, S., Maksimova, E., Robinson, K., Karpilow, J., et al. (2006). 3' UTR seed matches, but not overall identity, are associated with RNAi off-targets. *Nature methods* 3, 199-204.
- Blais, A., Tsikitis, M., Acosta-Alvear, D., Sharan, R., Kluger, Y., and Dynlacht, B.D. (2005). An initial blueprint for myogenic differentiation. *Genes & development* 19, 553-569.
- Blanchette, M., and Tompa, M. (2003). FootPrinter: A program designed for phylogenetic footprinting. *Nucleic acids research* 31, 3840-3842.
- Bochar, D.A., Savard, J., Wang, W., Lafleur, D.W., Moore, P., Cote, J., and Shiekhatar, R. (2000). A family of chromatin remodeling factors related to Williams syndrome transcription factor. *Proceedings of the National Academy of Sciences of the United States of America* 97, 1038-1043.
- Bockamp, E.O., Fordham, J.L., Gottgens, B., Murrell, A.M., Sanchez, M.J., and Green, A.R. (1998). Transcriptional regulation of the stem cell leukemia gene by PU.1 and Elf-1. *The Journal of biological chemistry* 273, 29032-29042.
- Bockamp, E.O., McLaughlin, F., Gottgens, B., Murrell, A.M., Elefanty, A.G., and Green, A.R. (1997). Distinct mechanisms direct SCL/tal-1 expression in erythroid cells and CD34 positive primitive myeloid cells. *The Journal of biological chemistry* 272, 8781-8790.
- Bode, J., Benham, C., Knopp, A., and Mielke, C. (2000). Transcriptional augmentation: modulation of gene expression by scaffold/matrix-attached regions (S/MAR elements). *Critical reviews in eukaryotic gene expression* 10, 73-90.
- Bode, J., Goetze, S., Heng, H., Krawetz, S.A., and Benham, C. (2003). From DNA structure to gene expression: mediators of nuclear compartmentalization and dynamics. *Chromosome Res* 11, 435-445.
- Boden, D., Pusch, O., Lee, F., Tucker, L., and Ramratnam, B. (2004). Efficient gene transfer of HIV-1-specific short hairpin RNA into human lymphocytic cells using recombinant adeno-associated virus vectors. *Mol Ther* 9, 396-402.
- Boden, D., Pusch, O., Silbermann, R., Lee, F., Tucker, L., and Ramratnam, B. (2004). Enhanced gene silencing of HIV-1 specific siRNA using microRNA designed hairpins. *Nucleic acids research* 32, 1154-1158.
- Boehm, T., Foroni, L., Kaneko, Y., Perutz, M.F., and Rabbitts, T.H. (1991). The rhombotin family of cysteine-rich LIM-domain oncogenes: distinct members are involved in T-cell translocations to human chromosomes 11p15 and 11p13. *Proceedings of the National Academy of Sciences of the United States of America* 88, 4367-4371.
- Bose, F., Fugazza, C., Casalgrandi, M., Capelli, A., Cunningham, J.M., Zhao, Q., Jane, S.M., Ottolenghi, S., and Ronchi, A. (2006). Functional interaction of CP2 with GATA-1 in the regulation of erythroid promoters. *Molecular and cellular biology* 26, 3942-3954.
- Boutros, M., Kiger, A.A., Armknecht, S., Kerr, K., Hild, M., Koch, B., Haas, S.A., Paro, R., and Perrimon, N. (2004). Genome-wide RNAi analysis of growth and viability in *Drosophila* cells. *Science (New York, NY)* 303, 832-835.
- Boyer, L.A., Lee, T.I., Cole, M.F., Johnstone, S.E., Levine, S.S., Zucker, J.P., Guenther, M.G., Kumar, R.M., Murray, H.L., Jenner, R.G., et al. (2005). Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 122, 947-956.
- Boyer, L.A., Plath, K., Zeitlinger, J., Brambrink, T., Medeiros, L.A., Lee, T.I., Levine, S.S., Wernig, M., Tajonar, A., Ray, M.K., et al. (2006). Polycomb complexes repress developmental regulators in murine embryonic stem cells. *Nature* 441, 349-353.
- Boyes, J., Byfield, P., Nakatani, Y., and Ogryzko, V. (1998). Regulation of activity of the transcription factor GATA-1 by acetylation. *Nature* 396, 594-598.
- Boyle, E.I., Weng, S., Gollub, J., Jin, H., Botstein, D., Cherry, J.M., and Sherlock, G. (2004). GO::TermFinder--open source software for accessing Gene Ontology information and finding significantly enriched Gene Ontology terms associated with a list of genes. *Bioinformatics (Oxford, England)* 20, 3710-3715.

- Brandenberger, R., Khrebtukova, I., Thies, R.S., Miura, T., Jingli, C., Puri, R., Vasicek, T., Lebkowski, J., and Rao, M. (2004). MPSS profiling of human embryonic stem cells. *BMC developmental biology* 4, 10.
- Brass, A.L., Dykxhoorn, D.M., Benita, Y., Yan, N., Engelman, A., Xavier, R.J., Lieberman, J., and Elledge, S.J. (2008). Identification of host proteins required for HIV infection through a functional genomic screen. *Science (New York, NY)* 319, 921-926.
- Brehm, A., Miska, E.A., McCance, D.J., Reid, J.L., Bannister, A.J., and Kouzarides, T. (1998). Retinoblastoma protein recruits histone deacetylase to repress transcription. *Nature* 391, 597-601.
- Brenner, S., Johnson, M., Bridgham, J., Golda, G., Lloyd, D.H., Johnson, D., Luo, S., McCurdy, S., Foy, M., Ewan, M., et al. (2000). Gene expression analysis by massively parallel signature sequencing (MPSS) on microbead arrays. *Nature biotechnology* 18, 630-634.
- Brenner, S., Williams, S.R., Vermaas, E.H., Storck, T., Moon, K., McCollum, C., Mao, J.I., Luo, S., Kirchner, J.J., Eletr, S., et al. (2000). In vitro cloning of complex mixtures of DNA on microbeads: physical separation of differentially expressed cDNAs. *Proceedings of the National Academy of Sciences of the United States of America* 97, 1665-1670.
- Brent, R., and Ptashne, M. (1985). A eukaryotic transcriptional activator bearing the DNA specificity of a prokaryotic repressor. *Cell* 43, 729-736.
- Bridge, A.J., Pebernard, S., Ducraux, A., Nicoulaz, A.L., and Iggo, R. (2003). Induction of an interferon response by RNAi vectors in mammalian cells. *Nature genetics* 34, 263-264.
- Brudno, M., Do, C.B., Cooper, G.M., Kim, M.F., Davydov, E., Green, E.D., Sidow, A., and Batzoglou, S. (2003). LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome research* 13, 721-731.
- Brummelkamp, T.R., Bernards, R., and Agami, R. (2002). A system for stable expression of short interfering RNAs in mammalian cells. *Science (New York, NY)* 296, 550-553.
- Brummelkamp, T.R., Nijman, S.M., Dirac, A.M., and Bernards, R. (2003). Loss of the cylindromatosis tumour suppressor inhibits apoptosis by activating NF-kappaB. *Nature* 424, 797-801.
- Cai, J., Shin, S., Wright, L., Liu, Y., Zhou, D., Xue, H., Khrebtukova, I., Mattson, M.P., Svendsen, C.N., and Rao, M.S. (2006). Massively parallel signature sequencing profiling of fetal human neural precursor cells. *Stem cells and development* 15, 232-244.
- Cao, H.B., Wang, A., Martin, B., Koehler, D.R., Zeitlin, P.L., Tanawell, A.K., and Hu, J. (2005). Down-regulation of IL-8 expression in human airway epithelial cells through helper-dependent adenoviral-mediated RNA interference. *Cell research* 15, 111-119.
- Cao, R., Wang, L., Wang, H., Xia, L., Erdjument-Bromage, H., Tempst, P., Jones, R.S., and Zhang, Y. (2002). Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science (New York, NY)* 298, 1039-1043.
- Caplen, N.J., Parrish, S., Imani, F., Fire, A., and Morgan, R.A. (2001). Specific inhibition of gene expression by small double-stranded RNAs in invertebrate and vertebrate systems. *Proceedings of the National Academy of Sciences of the United States of America* 98, 9742-9747.
- Capron C, Lécluse Y, Kaushik AL, Foudi A, Lacout C, Sekkai D, Godin I, Albagli O, Poullion I, Svinartchouk F, Schanze E, Vainchenker W, Sablitzky F, Bennaceur-Griscelli A, Duménil D. (2006). The SCL relative LYL-1 is required for fetal and adult hematopoietic stem cell function and B-cell differentiation. *Blood* 107, 4678-4686.
- Cawley, S., Bekiranov, S., Ng, H.H., Kapranov, P., Sekinger, E.A., Kampa, D., Piccolboni, A., Sementchenko, V., Cheng, J., Williams, A.J., et al. (2004). Unbiased mapping of transcription factor binding sites along human chromosomes 21 and 22 points to widespread regulation of noncoding RNAs. *Cell* 116, 499-509.
- Chambeyron, S., and Bickmore, W.A. (2004). Chromatin decondensation and nuclear reorganization of the HoxB locus upon induction of transcription. *Genes & development* 18, 1119-1130.
- Chan, W.C., and Nie, S. (1998). Quantum dot bioconjugates for ultrasensitive nonisotopic detection. *Science (New York, NY)* 281, 2016-2018.
- Chan, W.Y., Follows, G.A., Lacaud, G., Pimanda, J.E., Landry, J.R., Kinston, S., Knezevic, K., Piltz, S., Donaldson, I.J., Gambardella, L., et al. (2007). The paralogous hematopoietic regulators Ly11 and Scl are

- coregulated by Ets and GATA factors, but Lyl1 cannot rescue the early Scl^{-/-} phenotype. *Blood* 109, 1908-1916.
- Chen, A.A., Derfus, A.M., Khetani, S.R., and Bhatia, S.N. (2005). Quantum dots to monitor RNAi delivery and improve gene silencing. *Nucleic acids research* 33, e190.
- Chen, C.C., Ko, T.M., Ma, H.I., Wu, H.L., Xiao, X., Li, J., Chang, C.M., Wu, P.Y., Chen, C.H., Han, J.M., et al. (2007). Long-term inhibition of hepatitis B virus in transgenic mice by double-stranded adeno-associated virus 8-delivered short hairpin RNA. *Gene therapy* 14, 11-19.
- Chen, J.D., and Evans, R.M. (1995). A transcriptional co-repressor that interacts with nuclear hormone receptors. *Nature* 377, 454-457.
- Chen, L., and Widom, J. (2005). Mechanism of transcriptional silencing in yeast. *Cell* 120, 37-48.
- Chen, Y.T., Scanlan, M.J., Venditti, C.A., Chua, R., Theiler, G., Stevenson, B.J., Iseli, C., Gure, A.O., Vasicek, T., Strausberg, R.L., et al. (2005). Identification of cancer/testis-antigen genes by massively parallel signature sequencing. *Proceedings of the National Academy of Sciences of the United States of America* 102, 7940-7945.
- Choi, I., Cho, B.R., Kim, D., Miyagawa, S., Kubo, T., Kim, J.Y., Park, C.G., Hwang, W.S., Lee, J.S., and Ahn, C. (2005). Choice of the adequate detection time for the accurate evaluation of the efficiency of siRNA-induced gene silencing. *Journal of biotechnology* 120, 251-261.
- Choi, K., Kennedy, M., Kazarov, A., Papadimitriou, J.C., and Keller, G. (1998). A common precursor for hematopoietic and endothelial cells. *Development (Cambridge, England)* 125, 725-732.
- Chung, Y.S., Zhang, W.J., Arentson, E., Kingsley, P.D., Palis, J., and Choi, K. (2002). Lineage analysis of the hemangioblast as defined by FLK1 and SCL expression. *Development (Cambridge, England)* 129, 5511-5520.
- Clemens, J.C., Worby, C.A., Simonson-Leff, N., Muda, M., Maehama, T., Hemmings, B.A., and Dixon, J.E. (2000). Use of double-stranded RNA interference in *Drosophila* cell lines to dissect signal transduction pathways. *Proceedings of the National Academy of Sciences of the United States of America* 97, 6499-6503.
- Cohen, B.A., Mitra, R.D., Hughes, J.D., and Church, G.M. (2000). A computational analysis of whole-genome expression data reveals chromosomal domains of gene expression. *Nature genetics* 26, 183-186.
- Coller, H.A., Grandori, C., Tamayo, P., Colbert, T., Lander, E.S., Eisenman, R.N., and Golub, T.R. (2000). Expression analysis with oligonucleotide microarrays reveals that MYC regulates genes involved in growth, cell cycle, signaling, and adhesion. *Proceedings of the National Academy of Sciences of the United States of America* 97, 3260-3265.
- Corey, D.R. (2007). Chemical modification: the key to clinical application of RNA interference? *The Journal of clinical investigation* 117, 3615-3622.
- Courey, A.J., and Tjian, R. (1988). Analysis of Sp1 in vivo reveals multiple transcriptional domains, including a novel glutamine-rich activation motif. *Cell* 55, 887-898.
- Crawford, G.E., Davis, S., Scacheri, P.C., Renaud, G., Halawi, M.J., Erdos, M.R., Green, R., Meltzer, P.S., Wolfsberg, T.G., and Collins, F.S. (2006). DNase-chip: a high-resolution method to identify DNase I hypersensitive sites using tiled microarrays. *Nature methods* 3, 503-509.
- Crawford, G.E., Holt, I.E., Mullikin, J.C., Tai, D., Blakesley, R., Bouffard, G., Young, A., Masiello, C., Green, E.D., Wolfsberg, T.G., et al. (2004). Identifying gene regulatory elements by genome-wide recovery of DNase hypersensitive sites. *Proceedings of the National Academy of Sciences of the United States of America* 101, 992-997.
- Crawford, G.E., Holt, I.E., Whittle, J., Webb, B.D., Tai, D., Davis, S., Margulies, E.H., Chen, Y., Bernat, J.A., Ginsburg, D., et al. (2006). Genome-wide mapping of DNase hypersensitive sites using massively parallel signature sequencing (MPSS). *Genome research* 16, 123-131.
- Crossley, M., Tsang, A.P., Bieker, J.J., and Orkin, S.H. (1994). Regulation of the erythroid Kruppel-like factor (EKLF) gene promoter by the erythroid transcription factor GATA-1. *The Journal of biological chemistry* 269, 15440-15444.
- Curtis, D.J., Hall, M.A., Van Stekelenburg, L.J., Robb, L., Jane, S.M., and Begley, C.G. (2004). SCL is required for normal function of short-term repopulating hematopoietic stem cells. *Blood* 103, 3342-3348.

- Cuthbert, G.L., Daujat, S., Snowden, A.W., Erdjument-Bromage, H., Hagiwara, T., Yamada, M., Schneider, R., Gregory, P.D., Tempst, P., Bannister, A.J., et al. (2004). Histone deimination antagonizes arginine methylation. *Cell* 118, 545-553.
- Dahl, J.A., and Collas, P. (2007). Q2ChIP, a quick and quantitative chromatin immunoprecipitation assay, unravels epigenetic dynamics of developmentally regulated genes in human carcinoma cells. *Stem Cells* 25, 1037-1046.
- Dahmus, M.E. (1995). Phosphorylation of the C-terminal domain of RNA polymerase II. *Biochimica et biophysica acta* 1261, 171-182.
- Daniel, J.A., Torok, M.S., Sun, Z.W., Schieltz, D., Allis, C.D., Yates, J.R., 3rd, and Grant, P.A. (2004). Deubiquitination of histone H2B by a yeast acetyltransferase complex regulates transcription. *The Journal of biological chemistry* 279, 1867-1871.
- Das, P.M., Ramachandran, K., vanWert, J., and Singal, R. (2004). Chromatin immunoprecipitation assay. *BioTechniques* 37, 961-969.
- Davuluri, R.V., Grosse, I., and Zhang, M.Q. (2001). Computational identification of promoters and first exons in the human genome. *Nature genetics* 29, 412-417.
- De Gobbi, M., Anguita, E., Hughes, J., Sloane-Stanley, J.A., Sharpe, J.A., Koch, C.M., Dunham, I., Gibbons, R.J., Wood, W.G., and Higgs, D.R. (2007). Tissue-specific histone modification and transcription factor binding in alpha globin gene expression. *Blood* 110, 4503-4510.
- de Jonge, J., Holtrop, M., Wilschut, J., and Huckriede, A. (2006). Reconstituted influenza virus envelopes as an efficient carrier system for cellular delivery of small-interfering RNAs. *Gene therapy* 13, 400-411.
- Dekker, J., Rippe, K., Dekker, M., and Kleckner, N. (2002). Capturing chromosome conformation. *Science (New York, NY)* 295, 1306-1311.
- Delabesse, E., Ogilvy, S., Chapman, M.A., Piltz, S.G., Gottgens, B., and Green, A.R. (2005). Transcriptional regulation of the SCL locus: identification of an enhancer that targets the primitive erythroid lineage in vivo. *Molecular and cellular biology* 25, 5215-5225.
- Delassus, S., Titley, I., and Enver, T. (1999). Functional and molecular analysis of hematopoietic progenitors derived from the aorta-gonad-mesonephros region of the mouse embryo. *Blood* 94, 1495-1503.
- DeRisi, J.L., Iyer, V.R., and Brown, P.O. (1997). Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science (New York, NY)* 278, 680-686.
- Dermitzakis, E.T., and Clark, A.G. (2002). Evolution of transcription factor binding sites in Mammalian gene regulatory regions: conservation and turnover. *Molecular biology and evolution* 19, 1114-1121.
- Dhami, P. (2005). The SCL gene and transcriptional control of haematopoiesis. PhD Thesis, University of Cambridge.
- Dhami, P., Coffey, A.J., Abbs, S., Vermeesch, J.R., Dumanski, J.P., Woodward, K.J., Andrews, R.M., Langford, C., and Vetrie, D. (2005). Exon array CGH: detection of copy-number changes at the resolution of individual exons in the human genome. *Am J Hum Genet* 76, 750-762.
- Diakos, C., Krapf, G., Gerner, C., Inthal, A., Lemberger, C., Ban, J., Dohnal, A.M., and Panzer-Gruemayer, E.R. (2007). RNAi-mediated silencing of TEL/AML1 reveals a heat-shock protein- and survivin-dependent mechanism for survival. *Blood* 109, 2607-2610.
- Dickins, R.A., Hemann, M.T., Zilfou, J.T., Simpson, D.R., Ibarra, I., Hannon, G.J., and Lowe, S.W. (2005). Probing tumor phenotypes using stable and regulated synthetic microRNA precursors. *Nature genetics* 37, 1289-1295.
- Dobbin KK, Kawasaki ES, Petersen DW, Simon RM. (2005). Characterizing dye bias in microarray experiments. *Bioinformatics* 21, 2430-2437.
- Dorschner, M.O., Hawrylycz, M., Humbert, R., Wallace, J.C., Shafer, A., Kawamoto, J., Mack, J., Hall, R., Goldy, J., Sabo, P.J., et al. (2004). High-throughput localization of functional elements by quantitative chromatin profiling. *Nature methods* 1, 219-225.
- Down, T.A., and Hubbard, T.J. (2002). Computational detection and location of transcription start sites in mammalian genomic DNA. *Genome research* 12, 458-461.
- Down, T.A., and Hubbard, T.J. (2005). NestedMICA: sensitive inference of over-represented motifs in nucleic acid sequence. *Nucleic acids research* 33, 1445-1453.

- D'Souza, S.L., Elefanty, A.G., and Keller, G. (2005). SCL/Tal-1 is essential for hematopoietic commitment of the hemangioblast but not for its development. *Blood* 105, 3862-3870.
- Echeverri, C.J., Beachy, P.A., Baum, B., Boutros, M., Buchholz, F., Chanda, S.K., Downward, J., Ellenberg, J., Fraser, A.G., Hacohen, N., et al. (2006). Minimizing the risk of reporting false positives in large-scale RNAi screens. *Nature methods* 3, 777-779.
- Eisbacher, M., Holmes, M.L., Newton, A., Hogg, P.J., Khachigian, L.M., Crossley, M., and Chong, B.H. (2003). Protein-protein interaction between Fli-1 and GATA-1 mediates synergistic expression of megakaryocyte-specific genes through cooperative DNA binding. *Molecular and cellular biology* 23, 3427-3441.
- Eisen, M.B., Spellman, P.T., Brown, P.O., and Botstein, D. (1998). Cluster analysis and display of genome-wide expression patterns. *Proceedings of the National Academy of Sciences of the United States of America* 95, 14863-14868.
- Elbashir, S.M., Harborth, J., Lendeckel, W., Yalcin, A., Weber, K., and Tuschl, T. (2001). Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells. *Nature* 411, 494-498.
- Elbashir, S.M., Harborth, J., Weber, K., and Tuschl, T. (2002). Analysis of gene function in somatic mammalian cells using small interfering RNAs. *Methods (San Diego, Calif)* 26, 199-213.
- Elefanty, A.G., Begley, C.G., Hartley, L., Papaevangelidou, B., and Robb, L. (1999). SCL expression in the mouse embryo detected with a targeted lacZ reporter gene demonstrates its localization to hematopoietic, vascular, and neural tissues. *Blood* 94, 3754-3763.
- Elefanty, A.G., Robb, L., Birner, R., and Begley, C.G. (1997). Hematopoietic-specific genes are not induced during in vitro differentiation of scl-null embryonic stem cells. *Blood* 90, 1435-1447.
- Elkon, R., Rashi-Elkeles, S., Lerenthal, Y., Linhart, C., Tenne, T., Amariglio, N., Rechavi, G., Shamir, R., and Shiloh, Y. (2005). Dissection of a DNA-damage-induced transcriptional network using a combination of microarrays, RNA interference and computational promoter analysis. *Genome biology* 6, R43.
- Elwood, N.J., Green, A.R., Melder, A., Begley, C.G., and Nicola, N. (1994). The SCL protein displays cell-specific heterogeneity in size. *Leukemia* 8, 106-114.
- Endoh, M., Ogawa, M., Orkin, S., and Nishikawa, S. (2002). SCL/tal-1-dependent process determines a competence to select the definitive hematopoietic lineage prior to endothelial differentiation. *Embo J* 21, 6700-6708.
- Euskirchen, G., Royce, T.E., Bertone, P., Martone, R., Rinn, J.L., Nelson, F.K., Sayward, F., Luscombe, N.M., Miller, P., Gerstein, M., et al. (2004). CREB binds to multiple loci on human chromosome 22. *Molecular and cellular biology* 24, 3804-3814.
- Evans, T., Reitman, M., and Felsenfeld, G. (1988). An erythrocyte-specific DNA-binding factor recognizes a regulatory sequence common to all chicken globin genes. *Proceedings of the National Academy of Sciences of the United States of America* 85, 5976-5980.
- Fedorov, Y., Anderson, E.M., Birmingham, A., Reynolds, A., Karpilow, J., Robinson, K., Leake, D., Marshall, W.S., and Khvorova, A. (2006). Off-target effects by siRNA can induce toxic phenotype. *RNA (New York, NY)* 12, 1188-1196.
- Ferrando, A.A., Neuberg, D.S., Staunton, J., Loh, M.L., Huard, C., Raimondi, S.C., Behm, F.G., Pui, C.H., Downing, J.R., Gilliland, D.G., et al. (2002). Gene expression signatures define novel oncogenic pathways in T cell acute lymphoblastic leukemia. *Cancer cell* 1, 75-87.
- Ferreira, R., Ohneda, K., Yamamoto, M., and Philipsen, S. (2005). GATA1 function, a paradigm for transcription factors in hematopoiesis. *Molecular and cellular biology* 25, 1215-1227.
- Fire, A., Albertson, D., Harrison, S.W., and Moerman, D.G. (1991). Production of antisense RNA leads to effective and specific inhibition of gene expression in *C. elegans* muscle. *Development (Cambridge, England)* 113, 503-514.
- Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E., and Mello, C.C. (1998). Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391, 806-811.
- Follows, G.A., Dhami, P., Gottgens, B., Bruce, A.W., Campbell, P.J., Dillon, S.C., Smith, A.M., Koch, C., Donaldson, I.J., Scott, M.A., et al. (2006). Identifying gene regulatory elements by genomic microarray mapping of DNaseI hypersensitive sites. *Genome research* 16, 1310-1319.

- Follows, G.A., Janes, M.E., Vallier, L., Green, A.R., and Gottgens, B. (2007). Real-time PCR mapping of DNaseI-hypersensitive sites using a novel ligation-mediated amplification technique. *Nucleic acids research* 35, e56.
- Fordham, J.L., Gottgens, B., McLaughlin, F., and Green, A.R. (1999). Chromatin structure and transcriptional regulation of the stem cell leukaemia (SCL) gene in mast cells. *Leukemia* 13, 750-759.
- Fox, A.H., Liew, C., Holmes, M., Kowalski, K., Mackay, J., and Crossley, M. (1999). Transcriptional cofactors of the FOG family interact with GATA proteins by means of multiple zinc fingers. *Embo J* 18, 2812-2822.
- Freeman, L.A., and Garrard, W.T. (1992). DNA supercoiling in chromatin structure and gene expression. *Critical reviews in eukaryotic gene expression* 2, 165-209.
- Freson, K., Devriendt, K., Matthijs, G., Van Hoof, A., De Vos, R., Thys, C., Minner, K., Hoylaerts, M.F., Vermynen, J., and Van Geet, C. (2001). Platelet characteristics in patients with X-linked macrothrombocytopenia because of a novel GATA1 mutation. *Blood* 98, 85-92.
- Freson, K., Matthijs, G., Thys, C., Marien, P., Hoylaerts, M.F., Vermynen, J., and Van Geet, C. (2002). Different substitutions at residue D218 of the X-linked transcription factor GATA1 lead to altered clinical severity of macrothrombocytopenia and anemia and are associated with variable skewed X inactivation. *Human molecular genetics* 11, 147-152.
- Fujiwara, Y., Browne, C.P., Cunniff, K., Goff, S.C., and Orkin, S.H. (1996). Arrested development of embryonic red cell precursors in mouse embryos lacking transcription factor GATA-1. *Proceedings of the National Academy of Sciences of the United States of America* 93, 12355-12358.
- Galas, D.J., and Schmitz, A. (1978). DNase footprinting: a simple method for the detection of protein-DNA binding specificity. *Nucleic acids research* 5, 3157-3170.
- Gao, X., Cui, Y., Levenson, R.M., Chung, L.W., and Nie, S. (2004). In vivo cancer targeting and imaging with semiconductor quantum dots. *Nature biotechnology* 22, 969-976.
- Garner, M.M., and Revzin, A. (1981). A gel electrophoresis method for quantifying the binding of proteins to specific DNA regions: application to components of the Escherichia coli lactose operon regulatory system. *Nucleic acids research* 9, 3047-3060.
- Ge, Y., Dombkowski, A.A., LaFiura, K.M., Tatman, D., Yedidi, R.S., Stout, M.L., Buck, S.A., Massey, G., Becton, D.L., Weinstein, H.J., et al. (2006). Differential gene expression, GATA1 target genes, and the chemotherapy sensitivity of Down syndrome megakaryocytic leukemia. *Blood* 107, 1570-1581.
- Gelmetti, V., Zhang, J., Fanelli, M., Minucci, S., Pelicci, P.G., and Lazar, M.A. (1998). Aberrant recruitment of the nuclear receptor corepressor-histone deacetylase complex by the acute myeloid leukemia fusion partner ETO. *Molecular and cellular biology* 18, 7185-7191.
- Gershenzon, N.I., and Ioshikhes, I.P. (2005). Synergy of human Pol II core promoter elements revealed by statistical sequence analysis. *Bioinformatics (Oxford, England)* 21, 1295-1300.
- Gerton, J.L., DeRisi, J., Shroff, R., Lichten, M., Brown, P.O., and Petes, T.D. (2000). Inaugural article: global mapping of meiotic recombination hotspots and coldspots in the yeast *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences of the United States of America* 97, 11383-11390.
- Giroux, S., Kaushik, A.L., Capron, C., Jalil, A., Kelaidi, C., Sablitzky, F., Dumenil, D., Albagli, O., and Godin, I. (2007). *lyl-1* and *tal-1/scl*, two genes encoding closely related bHLH transcription factors, display highly overlapping expression patterns during cardiovascular and hematopoietic ontogeny. *Gene Expr Patterns* 7, 215-226.
- Goardon, N., Lambert, J.A., Rodriguez, P., Nissaire, P., Herblot, S., Thibault, P., Dumenil, D., Strouboulis, J., Romeo, P.H., and Hoang, T. (2006). ETO2 coordinates cellular proliferation and differentiation during erythropoiesis. *Embo J* 25, 357-366.
- Godin, I., and Cumano, A. (2002). The hare and the tortoise: an embryonic haematopoietic race. *Nat Rev Immunol* 2, 593-604.
- Godin, I., Garcia-Porrero, J.A., Dieterlen-Lievre, F., and Cumano, A. (1999). Stem cell emergence and hemopoietic activity are incompatible in mouse intraembryonic sites. *The Journal of experimental medicine* 190, 43-52.
- Goldfarb, A.N., Goueli, S., Mickelson, D., and Greenberg, J.M. (1992). T-cell acute lymphoblastic leukemia-the associated gene SCL/tal codes for a 42-Kd nuclear phosphoprotein. *Blood* 80, 2858-2866.

- Gonczy, P., Echeverri, C., Oegema, K., Coulson, A., Jones, S.J., Copley, R.R., Duperon, J., Oegema, J., Brehm, M., Cassin, E., et al. (2000). Functional genomic analysis of cell division in *C. elegans* using RNAi of genes on chromosome III. *Nature* 408, 331-336.
- Gonzalez, G.A., and Montminy, M.R. (1989). Cyclic AMP stimulates somatostatin gene transcription by phosphorylation of CREB at serine 133. *Cell* 59, 675-680.
- Gottgens, B., Barton, L.M., Gilbert, J.G., Bench, A.J., Sanchez, M.J., Bahn, S., Mistry, S., Grafham, D., McMurray, A., Vaudin, M., et al. (2000). Analysis of vertebrate SCL loci identifies conserved enhancers. *Nature biotechnology* 18, 181-186.
- Gottgens, B., Broccardo, C., Sanchez, M.J., Deveaux, S., Murphy, G., Gothert, J.R., Kotsopoulou, E., Kinston, S., Delaney, L., Piltz, S., et al. (2004). The scl +18/19 stem cell enhancer is not required for hematopoiesis: identification of a 5' bifunctional hematopoietic-endothelial enhancer bound by Fli-1 and Elf-1. *Molecular and cellular biology* 24, 1870-1883.
- Gottgens, B., McLaughlin, F., Bockamp, E.O., Fordham, J.L., Begley, C.G., Kosmopoulos, K., Elefanty, A.G., and Green, A.R. (1997). Transcription of the SCL gene in erythroid and CD34 positive primitive myeloid cells is controlled by a complex network of lineage-restricted chromatin-dependent and chromatin-independent regulatory elements. *Oncogene* 15, 2419-2428.
- Gottgens, B., Nastos, A., Kinston, S., Piltz, S., Delabesse, E.C., Stanley, M., Sanchez, M.J., Ciau-Uitz, A., Patient, R., and Green, A.R. (2002). Establishing the transcriptional programme for blood: the SCL stem cell enhancer is regulated by a multiprotein complex containing Ets and GATA factors. *Embo J* 21, 3039-3050.
- Gou, D., Jin, N., and Liu, L. (2003). Gene silencing in mammalian cells by PCR-based short hairpin RNA. *FEBS letters* 548, 113-118.
- Grant, P.A., Eberharter, A., John, S., Cook, R.G., Turner, B.M., and Workman, J.L. (1999). Expanded lysine acetylation specificity of Gcn5 in native complexes. *The Journal of biological chemistry* 274, 5895-5900.
- Grass, J.A., Boyer, M.E., Pal, S., Wu, J., Weiss, M.J., and Bresnick, E.H. (2003). GATA-1-dependent transcriptional repression of GATA-2 via disruption of positive autoregulation and domain-wide chromatin remodeling. *Proceedings of the National Academy of Sciences of the United States of America* 100, 8811-8816.
- Green, A.R., Lints, T., Visvader, J., Harvey, R., and Begley, C.G. (1992). SCL is coexpressed with GATA-1 in hemopoietic cells but is also expressed in developing brain. *Oncogene* 7, 653-660.
- Green, A.R., Rockman, S., DeLuca, E., and Begley, C.G. (1993). Induced myeloid differentiation of K562 cells with downregulation of erythroid and megakaryocytic transcription factors: a novel experimental model for hemopoietic lineage restriction. *Exp Hematol* 21, 525-531.
- Green, A.R., Salvaris, E., and Begley, C.G. (1991). Erythroid expression of the 'helix-loop-helix' gene, SCL. *Oncogene* 6, 475-479.
- Greenbaum, S., and Zhuang, Y. (2002). Identification of E2A target genes in B lymphocyte development by using a gene tagging-based chromatin immunoprecipitation system. *Proceedings of the National Academy of Sciences of the United States of America* 99, 15030-15035.
- Gregory, R.C., Taxman, D.J., Seshasayee, D., Kensinger, M.H., Bieker, J.J., and Wojchowski, D.M. (1996). Functional interaction of GATA1 with erythroid Kruppel-like factor and Sp1 at defined erythroid promoters. *Blood* 87, 1793-1801.
- Grimm, D., Streetz, K.L., Jopling, C.L., Storm, T.A., Pandey, K., Davis, C.R., Marion, P., Salazar, F., and Kay, M.A. (2006). Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature* 441, 537-541.
- Gross, D.S., and Garrard, W.T. (1988). Nuclease hypersensitive sites in chromatin. *Annu Rev Biochem* 57, 159-197.
- Grosveld, F., van Assendelft, G.B., Greaves, D.R., and Kollias, G. (1987). Position-independent, high-level expression of the human beta-globin gene in transgenic mice. *Cell* 51, 975-985.
- Grutz, G.G., Bucher, K., Lavenir, I., Larson, T., Larson, R., and Rabbitts, T.H. (1998). The oncogenic T cell LIM-protein Lmo2 forms part of a DNA-binding complex specifically in immature T cells. *Embo J* 17, 4594-4605.
- Gu, W., and Roeder, R.G. (1997). Activation of p53 sequence-specific DNA binding by acetylation of the p53 C-terminal domain. *Cell* 90, 595-606.

- Guenther, M.G., Levine, S.S., Boyer, L.A., Jaenisch, R., and Young, R.A. (2007). A chromatin landmark and transcription initiation at most promoters in human cells. *Cell* 130, 77-88.
- Gupta, S., Schoer, R.A., Egan, J.E., Hannon, G.J., and Mittal, V. (2004). Inducible, reversible, and stable RNA interference in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America* 101, 1927-1932.
- Gyorffy, B., Surowiak, P., Kiesslich, O., Denkert, C., Schafer, R., Dietel, M., and Lage, H. (2005). Gene expression profiling of 30 cancer cell lines predicts resistance towards 11 anticancer drugs at clinically achieved concentrations. *Int J Cancer*.
- Hahn, S. (2004). Structure and mechanism of the RNA polymerase II transcription machinery. *Nat Struct Mol Biol* 11, 394-403.
- Hamlett I, Draper J, Strouboulis J, Iborra F, Porcher C, Vyas P. Characterization of megakaryocyte GATA1-interacting proteins: the corepressor ETO2 and GATA1 interact to regulate terminal megakaryocyte maturation. (2008). *Blood* 112, 2738-2749.
- Hall, M.A., Curtis, D.J., Metcalf, D., Elefanty, A.G., Sourris, K., Robb, L., Gothert, J.R., Jane, S.M., and Begley, C.G. (2003). The critical regulator of embryonic hematopoiesis, SCL, is vital in the adult for megakaryopoiesis, erythropoiesis, and lineage choice in CFU-S12. *Proceedings of the National Academy of Sciences of the United States of America* 100, 992-997.
- Hall, M.A., Slater, N.J., Begley, C.G., Salmon, J.M., Van Stekelenburg, L.J., McCormack, M.P., Jane, S.M., and Curtis, D.J. (2005). Functional but abnormal adult erythropoiesis in the absence of the stem cell leukemia gene. *Molecular and cellular biology* 25, 6355-6362.
- Hao, H., Liu, H., Gonye, G., and Schwaber, J.S. (2008). A fast carrier chromatin immunoprecipitation method applicable to microdissected tissue samples. *Journal of neuroscience methods* 172, 38-42.
- Harris, M.B., Mostecky, J., and Rothman, P.B. (2005). Repression of an interleukin-4-responsive promoter requires cooperative BCL-6 function. *The Journal of biological chemistry* 280, 13114-13121.
- Hassa, P.O., Haenni, S.S., Elser, M., and Hottiger, M.O. (2006). Nuclear ADP-ribosylation reactions in mammalian cells: where are we today and where are we going? *Microbiol Mol Biol Rev* 70, 789-829.
- Hassan, A.H., Neely, K.E., and Workman, J.L. (2001). Histone acetyltransferase complexes stabilize swi/snf binding to promoter nucleosomes. *Cell* 104, 817-827.
- Heemskerk, M.H., Blom, B., Nolan, G., Stegmann, A.P., Bakker, A.Q., Weijer, K., Res, P.C., and Spits, H. (1997). Inhibition of T cell and promotion of natural killer cell development by the dominant negative helix loop helix factor Id3. *The Journal of experimental medicine* 186, 1597-1602.
- Heinemeyer, T., Wingender, E., Reuter, I., Hermjakob, H., Kel, A.E., Kel, O.V., Ignatieva, E.V., Ananko, E.A., Podkolodnaya, O.A., Kolpakov, F.A., et al. (1998). Databases on transcriptional regulation: TRANSFAC, TRRD and COMPEL. *Nucleic acids research* 26, 362-367.
- Heinzel, T., Lavinsky, R.M., Mullen, T.M., Soderstrom, M., Laherty, C.D., Torchia, J., Yang, W.M., Brard, G., Ngo, S.D., Davie, J.R., et al. (1997). A complex containing N-CoR, mSin3 and histone deacetylase mediates transcriptional repression. *Nature* 387, 43-48.
- Hirasawa, R., Shimizu, R., Takahashi, S., Osawa, M., Takayanagi, S., Kato, Y., Onodera, M., Minegishi, N., Yamamoto, M., Fukao, K., et al. (2002). Essential and instructive roles of GATA factors in eosinophil development. *The Journal of experimental medicine* 195, 1379-1386.
- Holen, T., Amarzguioui, M., Wiiger, M.T., Babaie, E., and Prydz, H. (2002). Positional effects of short interfering RNAs targeting the human coagulation trigger Tissue Factor. *Nucleic acids research* 30, 1757-1766.
- Holmes, M., Turner, J., Fox, A., Chisholm, O., Crossley, M., and Chong, B. (1999). hFOG-2, a novel zinc finger protein, binds the co-repressor mCtBP2 and modulates GATA-mediated activation. *The Journal of biological chemistry* 274, 23491-23498.
- Hope, I.A., Mahadevan, S., and Struhl, K. (1988). Structural and functional characterization of the short acidic transcriptional activation region of yeast GCN4 protein. *Nature* 333, 635-640.
- Hoppe, K.L., and Francone, O.L. (1998). Binding and functional effects of transcription factors Sp1 and Sp3 on the proximal human lecithin:cholesterol acyltransferase promoter. *Journal of lipid research* 39, 969-977.

- Horak, C.E., Mahajan, M.C., Luscombe, N.M., Gerstein, M., Weissman, S.M., and Snyder, M. (2002). GATA-1 binding sites mapped in the beta-globin locus by using mammalian ChIP-chip analysis. *Proceedings of the National Academy of Sciences of the United States of America* 99, 2924-2929.
- Horner, M.A., Quintin, S., Domeier, M.E., Kimble, J., Labouesse, M., and Mango, S.E. (1998). pha-4, an HNF-3 homolog, specifies pharyngeal organ identity in *Caenorhabditis elegans*. *Genes & development* 12, 1947-1952.
- Hsieh, A.C., Bo, R., Manola, J., Vazquez, F., Bare, O., Khvorova, A., Scaringe, S., and Sellers, W.R. (2004). A library of siRNA duplexes targeting the phosphoinositide 3-kinase pathway: determinants of gene silencing for use in cell-based screens. *Nucleic acids research* 32, 893-901.
- Hsu, H.L., Cheng, J.T., Chen, Q., and Baer, R. (1991). Enhancer-binding activity of the tal-1 oncoprotein in association with the E47/E12 helix-loop-helix proteins. *Molecular and cellular biology* 11, 3037-3042.
- Hsu, H.L., Huang, L., Tsan, J.T., Funk, W., Wright, W.E., Hu, J.S., Kingston, R.E., and Baer, R. (1994). Preferred sequences for DNA recognition by the TAL1 helix-loop-helix proteins. *Molecular and cellular biology* 14, 1256-1265.
- Hsu, H.L., Wadman, I., and Baer, R. (1994). Formation of in vivo complexes between the TAL1 and E2A polypeptides of leukemic T cells. *Proceedings of the National Academy of Sciences of the United States of America* 91, 3181-3185.
- Hsu, H.L., Wadman, I., Tsan, J.T., and Baer, R. (1994). Positive and negative transcriptional control by the TAL1 helix-loop-helix protein. *Proceedings of the National Academy of Sciences of the United States of America* 91, 5947-5951.
- Huang, D.Y., Kuo, Y.Y., and Chang, Z.F. (2005). GATA-1 mediates auto-regulation of Gfi-1B transcription in K562 cells. *Nucleic acids research* 33, 5331-5342.
- Huang, D.Y., Kuo, Y.Y., Lai, J.S., Suzuki, Y., Sugano, S., and Chang, Z.F. (2004). GATA-1 and NF-Y cooperate to mediate erythroid-specific transcription of Gfi-1B gene. *Nucleic acids research* 32, 3935-3946.
- Hughes, T.R., Mao, M., Jones, A.R., Burchard, J., Marton, M.J., Shannon, K.W., Lefkowitz, S.M., Ziman, M., Schelter, J.M., Meyer, M.R., et al. (2001). Expression profiling using microarrays fabricated by an ink-jet oligonucleotide synthesizer. *Nature biotechnology* 19, 342-347.
- Huo, X.F., Yu, J., Peng, H., Du, Z.W., Liu, X.L., Ma, Y.N., Zhang, X., Zhang, Y., Zhao, H.L., and Zhang, J.W. (2006). Differential expression changes in K562 cells during the hemin-induced erythroid differentiation and the phorbol myristate acetate (PMA)-induced megakaryocytic differentiation. *Molecular and cellular biochemistry* 292, 155-167.
- Hwang, L.Y., Siegelman, M., Davis, L., Oppenheimer-Marks, N., and Baer, R. (1993). Expression of the TAL1 proto-oncogene in cultured endothelial cells and blood vessels of the spleen. *Oncogene* 8, 3043-3046.
- Ikawa, T., Kawamoto, H., Wright, L.Y., and Murre, C. (2004). Long-term cultured E2A-deficient hematopoietic progenitor cells are pluripotent. *Immunity* 20, 349-360.
- Imagawa, S., Suzuki, N., Ohmine, K., Obara, N., Mukai, H.Y., Ozawa, K., Yamamoto, M., and Nagasawa, T. (2002). GATA suppresses erythropoietin gene expression through GATA site in mouse erythropoietin gene promoter. *International journal of hematology* 75, 376-381.
- Imagawa, S., Yamamoto, M., and Miura, Y. (1997). Negative regulation of the erythropoietin gene expression by the GATA transcription factors. *Blood* 89, 1430-1439.
- Inaba, T., Roberts, W.M., Shapiro, L.H., Jolly, K.W., Raimondi, S.C., Smith, S.D., and Look, A.T. (1992). Fusion of the leucine zipper gene HLF to the E2A gene in human acute B-lineage leukemia. *Science (New York, NY)* 257, 531-534.
- Irizarry, R.A., Bolstad, B.M., Collin, F., Cope, L.M., Hobbs, B., and Speed, T.P. (2003). Summaries of Affymetrix GeneChip probe level data. *Nucleic acids research* 31, e15.
- Irizarry, R.A., Hobbs, B., Collin, F., Beazer-Barclay, Y.D., Antonellis, K.J., Scherf, U., and Speed, T.P. (2003). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4, 249-264.
- Iwasaki, H., Mizuno, S., Wells, R.A., Cantor, A.B., Watanabe, S., and Akashi, K. (2003). GATA-1 converts lymphoid and myelomonocytic progenitors into the megakaryocyte/erythrocyte lineages. *Immunity* 19, 451-462.

- Iyer, V.R., Horak, C.E., Scafe, C.S., Botstein, D., Snyder, M., and Brown, P.O. (2001). Genomic binding sites of the yeast cell-cycle transcription factors SBF and MBF. *Nature* 409, 533-538.
- Jackson, A.L., Bartz, S.R., Schelter, J., Kobayashi, S.V., Burchard, J., Mao, M., Li, B., Cavet, G., and Linsley, P.S. (2003). Expression profiling reveals off-target gene regulation by RNAi. *Nature biotechnology* 21, 635-637.
- Jackson, A.L., Burchard, J., Leake, D., Reynolds, A., Schelter, J., Guo, J., Johnson, J.M., Lim, L., Karpilow, J., Nichols, K., et al. (2006). Position-specific chemical modification of siRNAs reduces "off-target" transcript silencing. *RNA (New York, NY)* 12, 1197-1205.
- Jackson, A.L., Burchard, J., Schelter, J., Chau, B.N., Cleary, M., Lim, L., and Linsley, P.S. (2006). Widespread siRNA "off-target" transcript silencing mediated by seed region sequence complementarity. *RNA (New York, NY)* 12, 1179-1187.
- Jazag, A., Ijichi, H., Kanai, F., Imamura, T., Guleng, B., Ohta, M., Imamura, J., Tanaka, Y., Tateishi, K., Ikenoue, T., et al. (2005). Smad4 silencing in pancreatic cancer cell lines using stable RNA interference and gene expression profiles induced by transforming growth factor-beta. *Oncogene* 24, 662-671.
- Jenuwein, T., and Allis, C.D. (2001). Translating the histone code. *Science (New York, NY)* 293, 1074-1080.
- Jiang, J., Chan, Y.S., Loh, Y.H., Cai, J., Tong, G.Q., Lim, C.A., Robson, P., Zhong, S., and Ng, H.H. (2008). A core Klf circuitry regulates self-renewal of embryonic stem cells. *Nature cell biology* 10, 353-360.
- Johnson, D.S., Mortazavi, A., Myers, R.M., and Wold, B. (2007). Genome-wide mapping of in vivo protein-DNA interactions. *Science (New York, NY)* 316, 1497-1502.
- Johnson, E.S. (2004). Protein modification by SUMO. *Annu Rev Biochem* 73, 355-382.
- Johnson, K.D., Grass, J.A., Boyer, M.E., Kiekhäfer, C.M., Blobel, G.A., Weiss, M.J., and Bresnick, E.H. (2002). Cooperative activities of hematopoietic regulators recruit RNA polymerase II to a tissue-specific chromatin domain. *Proceedings of the National Academy of Sciences of the United States of America* 99, 11760-11765.
- Jongeneel, C.V., Delorenzi, M., Iseli, C., Zhou, D., Haudenschild, C.D., Khrebtukova, I., Kuznetsov, D., Stevenson, B.J., Strausberg, R.L., Simpson, A.J., et al. (2005). An atlas of human gene expression from massively parallel signature sequencing (MPSS). *Genome research* 15, 1007-1014.
- Judge, A.D., Sood, V., Shaw, J.R., Fang, D., McClintock, K., and MacLachlan, I. (2005). Sequence-dependent stimulation of the mammalian innate immune response by synthetic siRNA. *Nature biotechnology* 23, 457-462.
- Jurata, L.W., and Gill, G.N. (1997). Functional analysis of the nuclear LIM domain interactor NLI. *Molecular and cellular biology* 17, 5688-5698.
- Kallianpur, A.R., Jordan, J.E., and Brandt, S.J. (1994). The SCL/TAL-1 gene is expressed in progenitors of both the hematopoietic and vascular systems during embryogenesis. *Blood* 83, 1200-1208.
- Kappel, A., Schlaeger, T.M., Flamme, I., Orkin, S.H., Risau, W., and Breier, G. (2000). Role of SCL/Tal-1, GATA, and ets transcription factor binding sites for the regulation of flk-1 expression during murine vascular development. *Blood* 96, 3078-3085.
- Kawasaki, H., and Taira, K. (2004). Induction of DNA methylation and gene silencing by short interfering RNAs in human cells. *Nature* 431, 211-217.
- Kee, B.L., and Murre, C. (1998). Induction of early B cell factor (EBF) and multiple B lineage genes by the basic helix-loop-helix transcription factor E12. *The Journal of experimental medicine* 188, 699-713.
- Kel, A.E., Gossling, E., Reuter, I., Cheremushkin, E., Kel-Margoulis, O.V., and Wingender, E. (2003). MATCH: A tool for searching transcription factor binding sites in DNA sequences. *Nucleic acids research* 31, 3576-3579.
- Kennedy, M., Firpo, M., Choi, K., Wall, C., Robertson, S., Kabrun, N., and Keller, G. (1997). A common precursor for primitive erythropoiesis and definitive haematopoiesis. *Nature* 386, 488-493.
- Kennerdell, J.R., and Carthew, R.W. (1998). Use of dsRNA-mediated genetic interference to demonstrate that frizzled and frizzled 2 act in the wingless pathway. *Cell* 95, 1017-1026.
- Kennerdell, J.R., and Carthew, R.W. (2000). Heritable gene silencing in *Drosophila* using double-stranded RNA. *Nature biotechnology* 18, 896-898.
- Kim, J., Bhinge, A.A., Morgan, X.C., and Iyer, V.R. (2005). Mapping DNA-protein interactions in large genomes by sequence tag analysis of genomic enrichment. *Nature methods* 2, 47-53.

- Kim, J., and Iyer, V.R. (2004). Global role of TATA box-binding protein recruitment to promoters in mediating gene expression profiles. *Molecular and cellular biology* 24, 8104-8112.
- Kim, J.B., Porreca, G.J., Song, L., Greenway, S.C., Gorham, J.M., Church, G.M., Seidman, C.E., and Seidman, J.G. (2007). Polony multiplex analysis of gene expression (PMAGE) in mouse hypertrophic cardiomyopathy. *Science (New York, NY)* 316, 1481-1484.
- Kim, T.H., Barrera, L.O., Qu, C., Van Calcar, S., Trinklein, N.D., Cooper, S.J., Luna, R.M., Glass, C.K., Rosenfeld, M.G., Myers, R.M., et al. (2005). Direct isolation and identification of promoters in the human genome. *Genome research* 15, 830-839.
- Kim, T.H., Barrera, L.O., Zheng, M., Qu, C., Singer, M.A., Richmond, T.A., Wu, Y., Green, R.D., and Ren, B. (2005). A high-resolution map of active promoters in the human genome. *Nature* 436, 876-880.
- Kittler, R., Pelletier, L., Heninger, A.K., Slabicki, M., Theis, M., Miroslaw, L., Poser, I., Lawo, S., Grabner, H., Kozak, K., et al. (2007). Genome-scale RNAi profiling of cell division in human tissue culture cells. *Nature cell biology* 9, 1401-1412.
- Kittler, R., Putz, G., Pelletier, L., Poser, I., Heninger, A.K., Drechsel, D., Fischer, S., Konstantinova, I., Habermann, B., Grabner, H., et al. (2004). An endoribonuclease-prepared siRNA screen in human cells identifies genes essential for cell division. *Nature* 432, 1036-1040.
- Kloosterman, W.P., and Plasterk, R.H. (2006). The diverse functions of microRNAs in animal development and disease. *Developmental cell* 11, 441-450.
- Kobayashi, M., Nishikawa, K., and Yamamoto, M. (2001). Hematopoietic regulatory domain of gata1 gene is positively regulated by GATA1 protein in zebrafish embryos. *Development (Cambridge, England)* 128, 2341-2350.
- Koch, C.M., Andrews, R.M., Flicek, P., Dillon, S.C., Karaoz, U., Clelland, G.K., Wilcox, S., Beare, D.M., Fowler, J.C., Couttet, P., et al. (2007). The landscape of histone modifications across 1% of the human genome in five human cell lines. *Genome research* 17, 691-707.
- Kodius, R., Kojima, M., Nishiyori, H., Nakamura, M., Fukuda, S., Tagami, M., Sasaki, D., Imamura, K., Kai, C., Harbers, M., et al. (2006). CAGE: cap analysis of gene expression. *Nature methods* 3, 211-222.
- Koizumi, K., Higashida, H., Yoo, S., Islam, M.S., Ivanov, A.I., Guo, V., Pozzi, P., Yu, S.H., Rovescalli, A.C., Tang, D., et al. (2007). RNA interference screen to identify genes required for Drosophila embryonic nervous system development. *Proceedings of the National Academy of Sciences of the United States of America* 104, 5626-5631.
- Kolfschoten, I.G., van Leeuwen, B., Berns, K., Mullenders, J., Beijersbergen, R.L., Bernards, R., Voorhoeve, P.M., and Agami, R. (2005). A genetic screen identifies PITX1 as a suppressor of RAS activity and tumorigenicity. *Cell* 121, 849-858.
- Krosl, G., He, G., Lefrancois, M., Charron, F., Romeo, P.H., Jolicoeur, P., Kirsch, I.R., Nemer, M., and Hoang, T. (1998). Transcription factor SCL is required for c-kit expression and c-Kit function in hemopoietic cells. *The Journal of experimental medicine* 188, 439-450.
- Kuhl, C., Atzberger, A., Iborra, F., Nieswandt, B., Porcher, C., and Vyas, P. (2005). GATA1-mediated megakaryocyte differentiation and growth control can be uncoupled and mapped to different domains in GATA1. *Molecular and cellular biology* 25, 8592-8606.
- Kuhn, K., Baker, S.C., Chudin, E., Lieu, M.H., Oeser, S., Bennett, H., Rigault, P., Barker, D., McDaniel, T.K., and Chee, M.S. (2004). A novel, high-performance random array platform for quantitative gene expression profiling. *Genome research* 14, 2347-2356.
- Kulesa, H., Frampton, J., and Graf, T. (1995). GATA-1 reprograms avian myelomonocytic cell lines into eosinophils, thromboblats, and erythroblats. *Genes & development* 9, 1250-1262.
- Kumaravelu, P., Hook, L., Morrison, A.M., Ure, J., Zhao, S., Zuyev, S., Ansell, J., and Medvinsky, A. (2002). Quantitative developmental anatomy of definitive haematopoietic stem cells/long-term repopulating units (HSC/RUs): role of the aorta-gonad-mesonephros (AGM) region and the yolk sac in colonisation of the mouse embryonic liver. *Development (Cambridge, England)* 129, 4891-4899.
- Kurdistani, S.K., Tavazoie, S., and Grunstein, M. (2004). Mapping global histone acetylation patterns to gene expression. *Cell* 117, 721-733.
- Labastie, M.C., Cortes, F., Romeo, P.H., Dulac, C., and Peault, B. (1998). Molecular identity of hematopoietic precursor cells emerging in the human embryo. *Blood* 92, 3624-3635.

- Lachner, M., and Jenuwein, T. (2002). The many faces of histone lysine methylation. *Curr Opin Cell Biol* 14, 286-298.
- Lahlil, R., Lecuyer, E., Herblot, S., and Hoang, T. (2004). SCL assembles a multifactorial complex that determines glycoporphin A expression. *Molecular and cellular biology* 24, 1439-1452.
- Lamond, A.I., and Earnshaw, W.C. (1998). Structure and function in the nucleus. *Science (New York, NY)* 280, 547-553.
- Landry, J.R., Kinston, S., Knezevic, K., de Bruijn, M.F., Wilson, N., Nottingham, W.T., Peitz, M., Edenhofer, F., Pimanda, J.E., Ottersbach, K., et al. (2008). Runx genes are direct targets of Scl/Tal1 in the yolk sac and fetal liver. *Blood* 111, 3005-3014.
- Larson, R.C., Lavenir, I., Larson, T.A., Baer, R., Warren, A.J., Wadman, I., Nottage, K., and Rabbitts, T.H. (1996). Protein dimerization between Lmo2 (Rbtn2) and Tal1 alters thymocyte development and potentiates T cell tumorigenesis in transgenic mice. *Embo J* 15, 1021-1027.
- Lassar, A.B., Davis, R.L., Wright, W.E., Kadesch, T., Murre, C., Voronova, A., Baltimore, D., and Weintraub, H. (1991). Functional activity of myogenic HLH proteins requires hetero-oligomerization with E12/E47-like proteins in vivo. *Cell* 66, 305-315.
- Lazrak, M., Deleuze, V., Noel, D., Haouzi, D., Chalhoub, E., Dohet, C., Robbins, I., and Mathieu, D. (2004). The bHLH TAL-1/SCL regulates endothelial cell migration and morphogenesis. *Journal of cell science* 117, 1161-1171.
- Lecointe, N., Bernard, O., Naert, K., Joulin, V., Larsen, C.J., Romeo, P.H., and Mathieu-Mahul, D. (1994). GATA- and SP1-binding sites are required for the full activity of the tissue-specific promoter of the tal-1 gene. *Oncogene* 9, 2623-2632.
- Lecuyer, E., Herblot, S., Saint-Denis, M., Martin, R., Begley, C.G., Porcher, C., Orkin, S.H., and Hoang, T. (2002). The SCL complex regulates c-kit expression in hematopoietic cells through functional interaction with Sp1. *Blood* 100, 2430-2440.
- Lee, D.H., Gershenson, N., Gupta, M., Ioshikhes, I.P., Reinberg, D., and Lewis, B.A. (2005). Functional characterization of core promoter elements: the downstream core element is recognized by TAF1. *Molecular and cellular biology* 25, 9674-9686.
- Lee, D.Y., Teyssier, C., Strahl, B.D., and Stallcup, M.R. (2005). Role of protein methylation in regulation of transcription. *Endocrine reviews* 26, 147-170.
- Lee, R.C., Feinbaum, R.L., and Ambros, V. (1993). The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 75, 843-854.
- Lee, T.I., Jenner, R.G., Boyer, L.A., Guenther, M.G., Levine, S.S., Kumar, R.M., Chevalier, B., Johnstone, S.E., Cole, M.F., Isono, K., et al. (2006). Control of developmental regulators by Polycomb in human embryonic stem cells. *Cell* 125, 301-313.
- Lee, T.I., Rinaldi, N.J., Robert, F., Odom, D.T., Bar-Joseph, Z., Gerber, G.K., Hannett, N.M., Harbison, C.T., Thompson, C.M., Simon, I., et al. (2002). Transcriptional regulatory networks in *Saccharomyces cerevisiae*. *Science (New York, NY)* 298, 799-804.
- Lee, T.I., and Young, R.A. (2000). Transcription of eukaryotic protein-coding genes. *Annual review of genetics* 34, 77-137.
- LeRoy, G., Rickards, B., and Flint, S.J. (2008). The double bromodomain proteins Brd2 and Brd3 couple histone acetylation to transcription. *Mol Cell* 30, 51-60.
- Leroy-Viard, K., Vinit, M.A., Lecointe, N., Mathieu-Mahul, D., and Romeo, P.H. (1994). Distinct DNase-I hypersensitive sites are associated with TAL-1 transcription in erythroid and T-cell lines. *Blood* 84, 3819-3827.
- Letting, D.L., Chen, Y.Y., Rakowski, C., Reedy, S., and Blobel, G.A. (2004). Context-dependent regulation of GATA-1 by friend of GATA-1. *Proceedings of the National Academy of Sciences of the United States of America* 101, 476-481.
- Leuther, K.K., Salmeron, J.M., and Johnston, S.A. (1993). Genetic evidence that an activation domain of GAL4 does not require acidity and may form a beta sheet. *Cell* 72, 575-585.
- Li, M., Li, H., and Rossi, J.J. (2006). RNAi in combination with a ribozyme and TAR decoy for treatment of HIV infection in hematopoietic cell gene therapy. *Annals of the New York Academy of Sciences* 1082, 172-179.

- Li, Q., Peterson, K.R., Fang, X., and Stamatoyannopoulos, G. (2002). Locus control regions. *Blood* 100, 3077-3086.
- Li, Z., Van Calcar, S., Qu, C., Cavenee, W.K., Zhang, M.Q., and Ren, B. (2003). A global transcriptional regulatory role for c-Myc in Burkitt's lymphoma cells. *Proceedings of the National Academy of Sciences of the United States of America* 100, 8164-8169.
- Liang, P., and Pardee, A.B. (1992). Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science (New York, NY)* 257, 967-971.
- Lieb, J.D., Liu, X., Botstein, D., and Brown, P.O. (2001). Promoter-specific binding of Rap1 revealed by genome-wide maps of protein-DNA association. *Nature genetics* 28, 327-334.
- Lieu, P.T., Jozsi, P., Gilles, P., and Peterson, T. (2005). Development of a DNA-labeling system for array-based comparative genomic hybridization. *J Biomol Tech* 16, 104-111.
- Lim, C.Y., Santoso, B., Boulay, T., Dong, E., Ohler, U., and Kadonaga, J.T. (2004). The MTE, a new core promoter element for transcription by RNA polymerase II. *Genes & development* 18, 1606-1617.
- Lin, Y.W., and Aplan, P.D. (2007). Gene expression profiling of precursor T-cell lymphoblastic leukemia/lymphoma identifies oncogenic pathways that are potential therapeutic targets. *Leukemia* 21, 1276-1284.
- Lipshutz, R.J., Fodor, S.P., Gingeras, T.R., and Lockhart, D.J. (1999). High density synthetic oligonucleotide arrays. *Nature genetics* 21, 20-24.
- Liu, X., Noll, D.M., Lieb, J.D., and Clarke, N.D. (2005). DIP-chip: rapid and accurate determination of DNA-binding specificity. *Genome research* 15, 421-427.
- Liu, X.S., Brutlag, D.L., and Liu, J.S. (2002). An algorithm for finding protein-DNA binding sites with applications to chromatin-immunoprecipitation microarray experiments. *Nature biotechnology* 20, 835-839.
- Loh, Y.H., Wu, Q., Chew, J.L., Vega, V.B., Zhang, W., Chen, X., Bourque, G., George, J., Leong, B., Liu, J., et al. (2006). The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nature genetics* 38, 431-440.
- Longabaugh, W.J., Davidson, E.H., and Bolouri, H. (2005). Computational representation of developmental genetic regulatory networks. *Developmental biology* 283, 1-16.
- Lorch, Y., Zhang, M., and Kornberg, R.D. (1999). Histone octamer transfer by a chromatin-remodeling complex. *Cell* 96, 389-392.
- Louie, M.C., Yang, H.Q., Ma, A.H., Xu, W., Zou, J.X., Kung, H.J., and Chen, H.W. (2003). Androgen-induced recruitment of RNA polymerase II to a nuclear receptor-p160 coactivator complex. *Proceedings of the National Academy of Sciences of the United States of America* 100, 2226-2230.
- Loukinov, D.I., Pugacheva, E., Vatolin, S., Pack, S.D., Moon, H., Chernukhin, I., Mannan, P., Larsson, E., Kanduri, C., Vostrov, A.A., et al. (2002). BORIS, a novel male germ-line-specific protein associated with epigenetic reprogramming events, shares the same 11-zinc-finger domain with CTCF, the insulator protein involved in reading imprinting marks in the soma. *Proceedings of the National Academy of Sciences of the United States of America* 99, 6806-6811.
- Loyola, A., Huang, J.Y., LeRoy, G., Hu, S., Wang, Y.H., Donnelly, R.J., Lane, W.S., Lee, S.C., and Reinberg, D. (2003). Functional analysis of the subunits of the chromatin assembly factor RSF. *Molecular and cellular biology* 23, 6759-6768.
- Lozzio, B.B., and Lozzio, C.B. (1977). Properties of the K562 cell line derived from a patient with chronic myeloid leukemia. *Int J Cancer* 19, 136.
- Lozzio, B.B., Lozzio, C.B., Bamberger, E.G., and Feliu, A.S. (1981). A multipotential leukemia cell line (K-562) of human origin. *Proceedings of the Society for Experimental Biology and Medicine Society for Experimental Biology and Medicine (New York, NY)* 166, 546-550.
- Lugus, J.J., Chung, Y.S., Mills, J.C., Kim, S.I., Grass, J., Kyba, M., Doherty, J.M., Bresnick, E.H., and Choi, K. (2007). GATA2 functions at multiple steps in hemangioblast development and differentiation. *Development (Cambridge, England)* 134, 393-405.
- Lum, L., Yao, S., Mozer, B., Rovescalli, A., Von Kessler, D., Nirenberg, M., and Beachy, P.A. (2003). Identification of Hedgehog pathway components by RNAi in *Drosophila* cultured cells. *Science (New York, NY)* 299, 2039-2045.

- Lupberger, J., Kreuzer, K.A., Baskaynak, G., Peters, U.R., le Coutre, P., and Schmidt, C.A. (2002). Quantitative analysis of beta-actin, beta-2-microglobulin and porphobilinogen deaminase mRNA and their comparison as control transcripts for RT-PCR. *Molecular and cellular probes* 16, 25-30.
- Mackay, J.P., Kowalski, K., Fox, A.H., Czolij, R., King, G.F., and Crossley, M. (1998). Involvement of the N-finger in the self-association of GATA-1. *The Journal of biological chemistry* 273, 30560-30567.
- MacKeigan, J.P., Murphy, L.O., and Blenis, J. (2005). Sensitized RNAi screen of human kinases and phosphatases identifies new regulators of apoptosis and chemoresistance. *Nature cell biology* 7, 591-600.
- Mahadevan, L.C., Willis, A.C., and Barratt, M.J. (1991). Rapid histone H3 phosphorylation in response to growth factors, phorbol esters, okadaic acid, and protein synthesis inhibitors. *Cell* 65, 775-783.
- Mansson, R., Tsapogas, P., Akerlund, M., Lagergren, A., Gisler, R., and Sigvardsson, M. (2004). Pearson correlation analysis of microarray data allows for the identification of genetic targets for early B-cell factor. *The Journal of biological chemistry* 279, 17905-17913.
- Mao, D.Y., Watson, J.D., Yan, P.S., Barysytte-Lovejoy, D., Khosravi, F., Wong, W.W., Farnham, P.J., Huang, T.H., and Penn, L.Z. (2003). Analysis of Myc bound loci identified by CpG island arrays shows that Max is essential for Myc-dependent repression. *Curr Biol* 13, 882-886.
- Margulies, M., Egholm, M., Altman, W.E., Attiya, S., Bader, J.S., Bemben, L.A., Berka, J., Braverman, M.S., Chen, Y.J., Chen, Z., et al. (2005). Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437, 376-380.
- Marodon, G., Mouly, E., Blair, E.J., Frisen, C., Lemoine, F.M., and Klatzmann, D. (2003). Specific transgene expression in human and mouse CD4+ cells using lentiviral vectors with regulatory sequences from the CD4 gene. *Blood* 101, 3416-3423.
- Marques, J.T., Devosse, T., Wang, D., Zamanian-Daryoush, M., Serbinowski, P., Hartmann, R., Fujita, T., Behlke, M.A., and Williams, B.R. (2006). A structural basis for discriminating between self and nonself double-stranded RNAs in mammalian cells. *Nature biotechnology* 24, 559-565.
- Martin, D.I., and Orkin, S.H. (1990). Transcriptional activation and DNA binding by the erythroid factor GF-1/NF-E1/Eryf 1. *Genes & development* 4, 1886-1898.
- Martin, D.I., Zon, L.I., Mutter, G., and Orkin, S.H. (1990). Expression of an erythroid transcription factor in megakaryocytic and mast cell lineages. *Nature* 344, 444-447.
- Martin, P., and Papayannopoulou, T. (1982). HEL cells: a new human erythroleukemia cell line with spontaneous and induced globin expression. *Science (New York, NY)* 216, 1233-1235.
- Marton, M.J., DeRisi, J.L., Bennett, H.A., Iyer, V.R., Meyer, M.R., Roberts, C.J., Stoughton, R., Burchard, J., Slade, D., Dai, H., et al. (1998). Drug target validation and identification of secondary drug target effects using DNA microarrays. *Nature medicine* 4, 1293-1301.
- Martone, R., Euskirchen, G., Bertone, P., Hartman, S., Royce, T.E., Luscombe, N.M., Rinn, J.L., Nelson, F.K., Miller, P., Gerstein, M., et al. (2003). Distribution of NF-kappaB-binding sites across human chromosome 22. *Proceedings of the National Academy of Sciences of the United States of America* 100, 12247-12252.
- Massari, M.E., Grant, P.A., Pray-Grant, M.G., Berger, S.L., Workman, J.L., and Murre, C. (1999). A conserved motif present in a class of helix-loop-helix proteins activates transcription by direct recruitment of the SAGA complex. *Mol Cell* 4, 63-73.
- Matsukura, S., Jones, P.A., and Takai, D. (2003). Establishment of conditional vectors for hairpin siRNA knockdowns. *Nucleic acids research* 31, e77.
- Matsumoto, N., Kubo, A., Liu, H., Akita, K., Laub, F., Ramirez, F., Keller, G., and Friedman, S.L. (2006). Developmental regulation of yolk sac hematopoiesis by Kruppel-like factor 6. *Blood* 107, 1357-1365.
- Matsumura, I., Kawasaki, A., Tanaka, H., Sonoyama, J., Ezoe, S., Minegishi, N., Nakajima, K., Yamamoto, M., and Kanakura, Y. (2000). Biologic significance of GATA-1 activities in Ras-mediated megakaryocytic differentiation of hematopoietic cell lines. *Blood* 96, 2440-2450.
- McArthur, M., Gerum, S., and Stamatoyannopoulos, G. (2001). Quantification of DNaseI-sensitivity by real-time PCR: quantitative analysis of DNaseI-hypersensitivity of the mouse beta-globin LCR. *Journal of molecular biology* 313, 27-34.

- Medina, K.L., Pongubala, J.M., Reddy, K.L., Lancki, D.W., Dekoter, R., Kieslinger, M., Grosschedl, R., and Singh, H. (2004). Assembling a gene regulatory network for specification of the B cell fate. *Developmental cell* 7, 607-617.
- Medvinsky, A., and Dzierzak, E. (1996). Definitive hematopoiesis is autonomously initiated by the AGM region. *Cell* 86, 897-906.
- Meersseman, G., Pennings, S., and Bradbury, E.M. (1992). Mobile nucleosomes--a general behavior. *Embo J* 11, 2951-2959.
- Mehaffey, M.G., Newton, A.L., Gandhi, M.J., Crossley, M., and Drachman, J.G. (2001). X-linked thrombocytopenia caused by a novel mutation of GATA-1. *Blood* 98, 2681-2688.
- Meister, G., Landthaler, M., Patkaniowska, A., Dorsett, Y., Teng, G., and Tuschl, T. (2004). Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol Cell* 15, 185-197.
- Mellentin, J.D., Smith, S.D., and Cleary, M.L. (1989). Iyl-1, a novel gene altered by chromosomal translocation in T cell leukemia, codes for a protein with a helix-loop-helix DNA binding motif. *Cell* 58, 77-83.
- Meng, A., Tang, H., Yuan, B., Ong, B.A., Long, Q., and Lin, S. (1999). Positive and negative cis-acting elements are required for hematopoietic expression of zebrafish GATA-1. *Blood* 93, 500-508.
- Merika, M., and Orkin, S.H. (1995). Functional synergy and physical interactions of the erythroid transcription factor GATA-1 with the Kruppel family proteins Sp1 and EKLF. *Molecular and cellular biology* 15, 2437-2447.
- Mermod, N., O'Neill, E.A., Kelly, T.J., and Tjian, R. (1989). The proline-rich transcriptional activator of CTF/NF-I is distinct from the replication and DNA binding domain. *Cell* 58, 741-753.
- Metzger, E., Wissmann, M., Yin, N., Muller, J.M., Schneider, R., Peters, A.H., Gunther, T., Buettner, R., and Schule, R. (2005). LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription. *Nature* 437, 436-439.
- Migliaccio, A.R., Rana, R.A., Sanchez, M., Lorenzini, R., Centurione, L., Bianchi, L., Vannucchi, A.M., Migliaccio, G., and Orkin, S.H. (2003). GATA-1 as a regulator of mast cell differentiation revealed by the phenotype of the GATA-1^{low} mouse mutant. *The Journal of experimental medicine* 197, 281-296.
- Mikkola, H.K., Klintman, J., Yang, H., Hock, H., Schlaeger, T.M., Fujiwara, Y., and Orkin, S.H. (2003). Haematopoietic stem cells retain long-term repopulating activity and multipotency in the absence of stem-cell leukaemia SCL/tal-1 gene. *Nature* 421, 547-551.
- Miller, I.J., and Bieker, J.J. (1993). A novel, erythroid cell-specific murine transcription factor that binds to the CACCC element and is related to the Kruppel family of nuclear proteins. *Molecular and cellular biology* 13, 2776-2786.
- Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., and Alon, U. (2002). Network motifs: simple building blocks of complex networks. *Science (New York, NY)* 298, 824-827.
- Milyavsky, M., Tabach, Y., Shats, I., Erez, N., Cohen, Y., Tang, X., Kalis, M., Kogan, I., Buganim, Y., Goldfinger, N., et al. (2005). Transcriptional programs following genetic alterations in p53, INK4A, and H-Ras genes along defined stages of malignant transformation. *Cancer research* 65, 4530-4543.
- Mineno, J., Okamoto, S., Ando, T., Sato, M., Chono, H., Izu, H., Takayama, M., Asada, K., Mirochnitchenko, O., Inouye, M., et al. (2006). The expression profile of microRNAs in mouse embryos. *Nucleic acids research* 34, 1765-1771.
- Moffat, J., Grueneberg, D.A., Yang, X., Kim, S.Y., Kloepfer, A.M., Hinkle, G., Piqani, B., Eisenhaure, T.M., Luo, B., Grenier, J.K., et al. (2006). A lentiviral RNAi library for human and mouse genes applied to an arrayed viral high-content screen. *Cell* 124, 1283-1298.
- Molkentin, J.D. (2000). The zinc finger-containing transcription factors GATA-4, -5, and -6. Ubiquitously expressed regulators of tissue-specific gene expression. *The Journal of biological chemistry* 275, 38949-38952.
- Moore, M.A., and Metcalf, D. (1970). Ontogeny of the haemopoietic system: yolk sac origin of in vivo and in vitro colony forming cells in the developing mouse embryo. *British journal of haematology* 18, 279-296.
- Morin, R.D., O'Connor, M.D., Griffith, M., Kuchenbauer, F., Delaney, A., Prabhu, A.L., Zhao, Y., McDonald, H., Zeng, T., Hirst, M., et al. (2008). Application of massively parallel sequencing to microRNA profiling and discovery in human embryonic stem cells. *Genome research*.

- Moroni, E., Mastrangelo, T., Razzini, R., Cairns, L., Moi, P., Ottolenghi, S., and Giglioni, B. (2000). Regulation of mouse p45 NF-E2 transcription by an erythroid-specific GATA-dependent intronic alternative promoter. *The Journal of biological chemistry* 275, 10567-10576.
- Morris, K.V., Chan, S.W., Jacobsen, S.E., and Looney, D.J. (2004). Small interfering RNA-induced transcriptional gene silencing in human cells. *Science (New York, NY)* 305, 1289-1292.
- Morrison, T.B., Weis, J.J., and Wittwer, C.T. (1998). Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. *BioTechniques* 24, 954-958, 960, 962.
- Mouthon, M.A., Bernard, O., Mitjavila, M.T., Romeo, P.H., Vainchenker, W., and Mathieu-Mahul, D. (1993). Expression of tal-1 and GATA-binding proteins during human hematopoiesis. *Blood* 81, 647-655.
- Mughal, T.I., and Goldman, J.M. (2006). Chronic myeloid leukemia: why does it evolve from chronic phase to blast transformation? *Front Biosci* 11, 198-208.
- Mukhopadhyay, M., Teufel, A., Yamashita, T., Agulnick, A.D., Chen, L., Downs, K.M., Schindler, A., Grinberg, A., Huang, S.P., Dorward, D., et al. (2003). Functional ablation of the mouse *Ldb1* gene results in severe patterning defects during gastrulation. *Development (Cambridge, England)* 130, 495-505.
- Muntean, A.G., and Crispino, J.D. (2005). Differential requirements for the activation domain and FOG-interaction surface of GATA-1 in megakaryocyte gene expression and development. *Blood* 106, 1223-1231.
- Murre, C., McCaw, P.S., and Baltimore, D. (1989). A new DNA binding and dimerization motif in immunoglobulin enhancer binding, daughterless, MyoD, and myc proteins. *Cell* 56, 777-783.
- Myers, J.W., Jones, J.T., Meyer, T., and Ferrell, J.E., Jr. (2003). Recombinant Dicer efficiently converts large dsRNAs into siRNAs suitable for gene silencing. *Nature biotechnology* 21, 324-328.
- Myers, L.C., and Kornberg, R.D. (2000). Mediator of transcriptional regulation. *Annu Rev Biochem* 69, 729-749.
- Nagy, P.L., Griesenbeck, J., Kornberg, R.D., and Cleary, M.L. (2002). A trithorax-group complex purified from *Saccharomyces cerevisiae* is required for methylation of histone H3. *Proceedings of the National Academy of Sciences of the United States of America* 99, 90-94.
- Nakamura, Y., Umehara, T., Nakano, K., Jang, M.K., Shirouzu, M., Morita, S., Uda-Tochio, H., Hamana, H., Terada, T., Adachi, N., et al. (2007). Crystal structure of the human BRD2 bromodomain: insights into dimerization and recognition of acetylated histone H4. *The Journal of biological chemistry* 282, 4193-4201.
- Neale, G.A., Rehg, J.E., and Goorha, R.M. (1995). Ectopic expression of rhombotin-2 causes selective expansion of CD4-CD8- lymphocytes in the thymus and T-cell tumors in transgenic mice. *Blood* 86, 3060-3071.
- Nelson, C.J., Santos-Rosa, H., and Kouzarides, T. (2006). Proline isomerization of histone H3 regulates lysine methylation and gene expression. *Cell* 126, 905-916.
- Ngo, V.N., Davis, R.E., Lamy, L., Yu, X., Zhao, H., Lenz, G., Lam, L.T., Dave, S., Yang, L., Powell, J., et al. (2006). A loss-of-function RNA interference screen for molecular targets in cancer. *Nature* 441, 106-110.
- Nichols, K.E., Crispino, J.D., Poncz, M., White, J.G., Orkin, S.H., Maris, J.M., and Weiss, M.J. (2000). Familial dyserythropoietic anaemia and thrombocytopenia due to an inherited mutation in GATA1. *Nature genetics* 24, 266-270.
- Nicolis, S., Bertini, C., Ronchi, A., Crotta, S., Lanfranco, L., Moroni, E., Giglioni, B., and Ottolenghi, S. (1991). An erythroid specific enhancer upstream to the gene encoding the cell-type specific transcription factor GATA-1. *Nucleic acids research* 19, 5285-5291.
- Nilsson, R., Bajic, V.B., Suzuki, H., di Bernardo, D., Bjorkegren, J., Katayama, S., Reid, J.F., Sweet, M.J., Gariboldi, M., Carninci, P., et al. (2006). Transcriptional network dynamics in macrophage activation. *Genomics* 88, 133-142.
- Nishikawa, K., Kobayashi, M., Masumi, A., Lyons, S.E., Weinstein, B.M., Liu, P.P., and Yamamoto, M. (2003). Self-association of Gata1 enhances transcriptional activity in vivo in zebra fish embryos. *Molecular and cellular biology* 23, 8295-8305.
- Nishimura, S., Takahashi, S., Kuroha, T., Suwabe, N., Nagasawa, T., Trainor, C., and Yamamoto, M. (2000). A GATA box in the GATA-1 gene hematopoietic enhancer is a critical element in the network of GATA factors and sites that regulate this gene. *Molecular and cellular biology* 20, 713-723.
- Noble, W.S., Kuehn, S., Thurman, R., Yu, M., and Stamatoyannopoulos, J. (2005). Predicting the in vivo signature of human gene regulatory sequences. *Bioinformatics (Oxford, England)* 21 Suppl 1, i338-343.

- Nordhoff, E., Krogsdam, A.M., Jorgensen, H.F., Kallipolitis, B.H., Clark, B.F., Roepstorff, P., and Kristiansen, K. (1999). Rapid identification of DNA-binding proteins by mass spectrometry. *Nature biotechnology* 17, 884-888.
- Nottingham, W.T., Jarratt, A., Burgess, M., Speck, C.L., Cheng, J.F., Prabhakar, S., Rubin, E.M., Li, P.S., Sloane-Stanley, J., Kong, A.S.J., et al. (2007). Runx1-mediated hematopoietic stem-cell emergence is controlled by a Gata/Ets/SCL-regulated enhancer. *Blood* 110, 4188-4197.
- Nourse, J., Mellentin, J.D., Galili, N., Wilkinson, J., Stanbridge, E., Smith, S.D., and Cleary, M.L. (1990). Chromosomal translocation t(1;19) results in synthesis of a homeobox fusion mRNA that codes for a potential chimeric transcription factor. *Cell* 60, 535-545.
- Nuez, B., Michalovich, D., Bygrave, A., Ploemacher, R., and Grosveld, F. (1995). Defective haematopoiesis in fetal liver resulting from inactivation of the EKLf gene. *Nature* 375, 316-318.
- Nunez, E., Kwon, Y.S., Hutt, K.R., Hu, Q., Cardamone, M.D., Ohgi, K.A., Garcia-Bassets, I., Rose, D.W., Glass, C.K., Rosenfeld, M.G., et al. (2008). Nuclear receptor-enhanced transcription requires motor- and LSD1-dependent gene networking in interchromatin granules. *Cell* 134, 189.
- Oberley, M.J., Inman, D.R., and Farnham, P.J. (2003). E2F6 negatively regulates BRCA1 in human cancer cells without methylation of histone H3 on lysine 9. *The Journal of biological chemistry* 278, 42466-42476.
- Oda, K., Kimura, T., Matsuoka, Y., Funahashi, A., Muramatsu, M., and Kitano, H. (2004). Molecular Interaction Map of Macrophage. *AfCS Research Reports* 2.
- Odom, D.T., Dowell, R.D., Jacobsen, E.S., Gordon, W., Danford, T.W., MacIsaac, K.D., Rolfe, P.A., Conboy, C.M., Gifford, D.K., and Fraenkel, E. (2007). Tissue-specific transcriptional regulation has diverged significantly between human and mouse. *Nature genetics* 39, 730-732.
- Odom, D.T., Zizlsperger, N., Gordon, D.B., Bell, G.W., Rinaldi, N.J., Murray, H.L., Volkert, T.L., Schreiber, J., Rolfe, P.A., Gifford, D.K., et al. (2004). Control of pancreas and liver gene expression by HNF transcription factors. *Science (New York, NY)* 303, 1378-1381.
- Ogilvy, S., Ferreira, R., Piltz, S.G., Bowen, J.M., Gottgens, B., and Green, A.R. (2007). The SCL +40 enhancer targets the midbrain together with primitive and definitive hematopoiesis and is regulated by SCL and GATA proteins. *Molecular and cellular biology* 27, 7206-7219.
- Oliva, A., Rosebrock, A., Ferrezuelo, F., Pyne, S., Chen, H., Skiena, S., Fitcher, B., and Leatherwood, J. (2005). The cell cycle-regulated genes of *Schizosaccharomyces pombe*. *PLoS biology* 3, e225.
- Olsen, P.H., and Ambros, V. (1999). The lin-4 regulatory RNA controls developmental timing in *Caenorhabditis elegans* by blocking LIN-14 protein synthesis after the initiation of translation. *Developmental biology* 216, 671-680.
- O'Neil, J., Shank, J., Cusson, N., Murre, C., and Kelliher, M. (2004). TAL1/SCL induces leukemia by inhibiting the transcriptional activity of E47/HEB. *Cancer cell* 5, 587-596.
- O'Neill, L.P., and Turner, B.M. (1996). Immunoprecipitation of chromatin. *Methods in enzymology* 274, 189-197.
- O'Neill, L.P., VerMilyea, M.D., and Turner, B.M. (2006). Epigenetic characterization of the early embryo with a chromatin immunoprecipitation protocol applicable to small cell populations. *Nature genetics* 38, 835-841.
- Onizuka, T., Endo, S., Hirano, M., Kanai, S., and Akiyama, H. (2002). Design of a fluorescent electrophoretic mobility shift assay improved for the quantitative and multiple analysis of protein-DNA complexes. *Bioscience, biotechnology, and biochemistry* 66, 2732-2734.
- Ono, Y., Fukuhara, N., and Yoshie, O. (1997). Transcriptional activity of TAL1 in T cell acute lymphoblastic leukemia (T-ALL) requires RBTN1 or -2 and induces TALLA1, a highly specific tumor marker of T-ALL. *The Journal of biological chemistry* 272, 4576-4581.
- Ono, Y., Fukuhara, N., and Yoshie, O. (1998). TAL1 and LIM-only proteins synergistically induce retinaldehyde dehydrogenase 2 expression in T-cell acute lymphoblastic leukemia by acting as cofactors for GATA3. *Molecular and cellular biology* 18, 6939-6950.
- Onodera, K., Takahashi, S., Nishimura, S., Ohta, J., Motohashi, H., Yomogida, K., Hayashi, N., Engel, J.D., and Yamamoto, M. (1997). GATA-1 transcription is controlled by distinct regulatory mechanisms during primitive and definitive erythropoiesis. *Proceedings of the National Academy of Sciences of the United States of America* 94, 4487-4492.

- Onodera, K., Yomogida, K., Suwabe, N., Takahashi, S., Muraosa, Y., Hayashi, N., Ito, E., Gu, L., Rassoulzadegan, M., Engel, J.D., et al. (1997). Conserved structure, regulatory elements, and transcriptional regulation from the GATA-1 gene testis promoter. *Journal of biochemistry* 121, 251-263.
- Orian, A., van Steensel, B., Delrow, J., Bussemaker, H.J., Li, L., Sawado, T., Williams, E., Loo, L.W., Cowley, S.M., Yost, C., et al. (2003). Genomic binding by the Drosophila Myc, Max, Mad/Mnt transcription factor network. *Genes & development* 17, 1101-1114.
- Orlando, V., Strutt, H., and Paro, R. (1997). Analysis of chromatin structure by in vivo formaldehyde cross-linking. *Methods (San Diego, Calif)* 11, 205-214.
- Pabo, C.O., and Sauer, R.T. (1992). Transcription factors: structural families and principles of DNA recognition. *Annu Rev Biochem* 61, 1053-1095.
- Paddison, P.J., Caudy, A.A., Bernstein, E., Hannon, G.J., and Conklin, D.S. (2002). Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells. *Genes & development* 16, 948-958.
- Pal, S., Cantor, A.B., Johnson, K.D., Moran, T.B., Boyer, M.E., Orkin, S.H., and Bresnick, E.H. (2004). Coregulator-dependent facilitation of chromatin occupancy by GATA-1. *Proceedings of the National Academy of Sciences of the United States of America* 101, 980-985.
- Palomero, T., Odom, D.T., O'Neil, J., Ferrando, A.A., Margolin, A., Neuberg, D.S., Winter, S.S., Larson, R.S., Li, W., Liu, X.S., et al. (2006). Transcriptional regulatory networks downstream of TAL1/SCL in T-cell acute lymphoblastic leukemia. *Blood* 108, 986-992.
- Parry, D.H., Xu, J., and Ruvkun, G. (2007). A whole-genome RNAi Screen for *C. elegans* miRNA pathway genes. *Curr Biol* 17, 2013-2022.
- Pebernard, S., and Iggo, R.D. (2004). Determinants of interferon-stimulated gene induction by RNAi vectors. *Differentiation; research in biological diversity* 72, 103-111.
- Pennacchio, L.A., Ahituv, N., Moses, A.M., Prabhakar, S., Nobrega, M.A., Shoukry, M., Minovitsky, S., Dubchak, I., Holt, A., Lewis, K.D., et al. (2006). In vivo enhancer analysis of human conserved non-coding sequences. *Nature* 444, 499-502.
- Pevny, L., Lin, C.S., D'Agati, V., Simon, M.C., Orkin, S.H., and Costantini, F. (1995). Development of hematopoietic cells lacking transcription factor GATA-1. *Development (Cambridge, England)* 121, 163-172.
- Pevny, L., Simon, M.C., Robertson, E., Klein, W.H., Tsai, S.F., D'Agati, V., Orkin, S.H., and Costantini, F. (1991). Erythroid differentiation in chimaeric mice blocked by a targeted mutation in the gene for transcription factor GATA-1. *Nature* 349, 257-260.
- Pokholok, D.K., Hannett, N.M., and Young, R.A. (2002). Exchange of RNA polymerase II initiation and elongation factors during gene expression in vivo. *Mol Cell* 9, 799-809.
- Pokholok, D.K., Harbison, C.T., Levine, S., Cole, M., Hannett, N.M., Lee, T.I., Bell, G.W., Walker, K., Rolfe, P.A., Herbolsheimer, E., et al. (2005). Genome-wide map of nucleosome acetylation and methylation in yeast. *Cell* 122, 517-527.
- Poot, R.A., Dellaire, G., Hulsmann, B.B., Grimaldi, M.A., Corona, D.F., Becker, P.B., Bickmore, W.A., and Varga-Weisz, P.D. (2000). HuCHRAC, a human ISWI chromatin remodelling complex contains hACF1 and two novel histone-fold proteins. *Embo J* 19, 3377-3387.
- Porcher, C., Swat, W., Rockwell, K., Fujiwara, Y., Alt, F.W., and Orkin, S.H. (1996). The T cell leukemia oncoprotein SCL/tal-1 is essential for development of all hematopoietic lineages. *Cell* 86, 47-57.
- Privalsky, M.L. (2004). The role of corepressors in transcriptional regulation by nuclear hormone receptors. *Annual review of physiology* 66, 315-360.
- Ptashne, M., and Gann, A. (1997). Transcriptional activation by recruitment. *Nature* 386, 569-577.
- Pulford, K., Lecoite, N., Leroy-Viard, K., Jones, M., Mathieu-Mahul, D., and Mason, D.Y. (1995). Expression of TAL-1 proteins in human tissues. *Blood* 85, 675-684.
- Qiu, Y., Sharma, A., and Stein, R. (1998). p300 mediates transcriptional stimulation by the basic helix-loop-helix activators of the insulin gene. *Molecular and cellular biology* 18, 2957-2964.
- Rea, S., Eisenhaber, F., O'Carroll, D., Strahl, B.D., Sun, Z.W., Schmid, M., Opravil, S., Mechtler, K., Ponting, C.P., Allis, C.D., et al. (2000). Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature* 406, 593-599.

- Rekhtman, N., Radparvar, F., Evans, T., and Skoultschi, A.I. (1999). Direct interaction of hematopoietic transcription factors PU.1 and GATA-1: functional antagonism in erythroid cells. *Genes & development* 13, 1398-1411.
- Ren, B., Cam, H., Takahashi, Y., Volkert, T., Terragni, J., Young, R.A., and Dynlacht, B.D. (2002). E2F integrates cell cycle progression with DNA repair, replication, and G(2)/M checkpoints. *Genes & development* 16, 245-256.
- Ren, B., Robert, F., Wyrick, J.J., Aparicio, O., Jennings, E.G., Simon, I., Zeitlinger, J., Schreiber, J., Hannett, N., Kanin, E., et al. (2000). Genome-wide location and function of DNA binding proteins. *Science (New York, NY)* 290, 2306-2309.
- Reynaud, D., Ravet, E., Titeux, M., Mazurier, F., Renia, L., Dubart-Kupperschmitt, A., Romeo, P.H., and Pflumio, F. (2005). SCL/TAL1 expression level regulates human hematopoietic stem cell self-renewal and engraftment. *Blood* 106, 2318-2328.
- Rippe, K., von Hippel, P.H., and Langowski, J. (1995). Action at a distance: DNA-looping and initiation of transcription. *Trends in biochemical sciences* 20, 500-506.
- Robb, L., Elwood, N.J., Elefanty, A.G., Kontgen, F., Li, R., Barnett, L.D., and Begley, C.G. (1996). The scl gene product is required for the generation of all hematopoietic lineages in the adult mouse. *Embo J* 15, 4123-4129.
- Robb, L., Lyons, I., Li, R., Hartley, L., Kontgen, F., Harvey, R.P., Metcalf, D., and Begley, C.G. (1995). Absence of yolk sac hematopoiesis from mice with a targeted disruption of the scl gene. *Proceedings of the National Academy of Sciences of the United States of America* 92, 7075-7079.
- Robert, F., Pokholok, D.K., Hannett, N.M., Rinaldi, N.J., Chandy, M., Rolfe, A., Workman, J.L., Gifford, D.K., and Young, R.A. (2004). Global position and recruitment of HATs and HDACs in the yeast genome. *Mol Cell* 16, 199-209.
- Robertson, S.M., Kennedy, M., Shannon, J.M., and Keller, G. (2000). A transitional stage in the commitment of mesoderm to hematopoiesis requiring the transcription factor SCL/tal-1. *Development (Cambridge, England)* 127, 2447-2459.
- Roby, D., Suka, Y., Xenarios, I., Kurdistani, S.K., Wang, A., Suka, N., and Grunstein, M. (2002). Microarray deacetylation maps determine genome-wide functions for yeast histone deacetylases. *Cell* 109, 437-446.
- Robzyk, K., Recht, J., and Osley, M.A. (2000). Rad6-dependent ubiquitination of histone H2B in yeast. *Science (New York, NY)* 287, 501-504.
- Roth, F.P., Hughes, J.D., Estep, P.W., and Church, G.M. (1998). Finding DNA regulatory motifs within unaligned noncoding sequences clustered by whole-genome mRNA quantitation. *Nature biotechnology* 16, 939-945.
- Rylski, M., Welch, J.J., Chen, Y.Y., Letting, D.L., Diehl, J.A., Chodosh, L.A., Blobel, G.A., and Weiss, M.J. (2003). GATA-1-mediated proliferation arrest during erythroid maturation. *Molecular and cellular biology* 23, 5031-5042.
- Saha, A., Wittmeyer, J., and Cairns, B.R. (2006). Chromatin remodelling: the industrial revolution of DNA around histones. *Nature reviews* 7, 437-447.
- Salghetti, S.E., Caudy, A.A., Chenoweth, J.G., and Tansey, W.P. (2001). Regulation of transcriptional activation domain function by ubiquitin. *Science (New York, NY)* 293, 1651-1653.
- Sanchez, M., Gottgens, B., Sinclair, A.M., Stanley, M., Begley, C.G., Hunter, S., and Green, A.R. (1999). An SCL 3' enhancer targets developing endothelium together with embryonic and adult haematopoietic progenitors. *Development (Cambridge, England)* 126, 3891-3904.
- Sanchez, M.J., Bockamp, E.O., Miller, J., Gambardella, L., and Green, A.R. (2001). Selective rescue of early haematopoietic progenitors in Scl(-/-) mice by expressing Scl under the control of a stem cell enhancer. *Development (Cambridge, England)* 128, 4815-4827.
- Sandelin, A., Alkema, W., Engstrom, P., Wasserman, W.W., and Lenhard, B. (2004). JASPAR: an open-access database for eukaryotic transcription factor binding profiles. *Nucleic acids research* 32, D91-94.
- Sassone-Corsi, P., Mizzen, C.A., Cheung, P., Crosio, C., Monaco, L., Jacquot, S., Hanauer, A., and Allis, C.D. (1999). Requirement of Rsk-2 for epidermal growth factor-activated phosphorylation of histone H3. *Science (New York, NY)* 285, 886-891.

- Scacheri, P.C., Rozenblatt-Rosen, O., Caplen, N.J., Wolfsberg, T.G., Umayam, L., Lee, J.C., Hughes, C.M., Shanmugam, K.S., Bhattacharjee, A., Meyerson, M., et al. (2004). Short interfering RNAs can induce unexpected and divergent changes in the levels of untargeted proteins in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America* 101, 1892-1897.
- Schena, M., Shalon, D., Davis, R.W., and Brown, P.O. (1995). Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science (New York, NY)* 270, 467-470.
- Scherf, M., Klingenhoff, A., and Werner, T. (2000). Highly specific localization of promoter regions in large genomic sequences by PromoterInspector: a novel context analysis approach. *Journal of molecular biology* 297, 599-606.
- Schiebel, W., Pelissier, T., Riedel, L., Thalmeir, S., Schiebel, R., Kempe, D., Lottspeich, F., Sanger, H.L., and Wassenegger, M. (1998). Isolation of an RNA-directed RNA polymerase-specific cDNA clone from tomato. *The Plant cell* 10, 2087-2101.
- Schlabach, M.R., Luo, J., Solimini, N.L., Hu, G., Xu, Q., Li, M.Z., Zhao, Z., Smogorzewska, A., Sowa, M.E., Ang, X.L., et al. (2008). Cancer proliferation gene discovery through functional genomics. *Science (New York, NY)* 319, 620-624.
- Schmitter, D., Filkowski, J., Sewer, A., Pillai, R.S., Oakeley, E.J., Zavolan, M., Svoboda, P., and Filipowicz, W. (2006). Effects of Dicer and Argonaute down-regulation on mRNA levels in human HEK293 cells. *Nucleic acids research* 34, 4801-4815.
- Schones, D.E., Cui, K., Cuddapah, S., Roh, T.Y., Barski, A., Wang, Z., Wei, G., and Zhao, K. (2008). Dynamic regulation of nucleosome positioning in the human genome. *Cell* 132, 887-898.
- Schug, J., Overton, G.C. (1997). TESS: Transcriptional Element Search Software on the WWW. Technical Report CBIL-TR-1997-1001-v0.0. . Computational Biology and Informatics Laboratory, School of Medicine, University of Pennsylvania.
- Schuh, A.H., Tipping, A.J., Clark, A.J., Hamlett, I., Guyot, B., Iborra, F.J., Rodriguez, P., Strouboulis, J., Enver, T., Vyas, P., et al. (2005). ETO-2 associates with SCL in erythroid cells and megakaryocytes and provides repressor functions in erythropoiesis. *Molecular and cellular biology* 25, 10235-10250.
- Segal, E., Shapira, M., Regev, A., Pe'er, D., Botstein, D., Koller, D., and Friedman, N. (2003). Module networks: identifying regulatory modules and their condition-specific regulators from gene expression data. *Nature genetics* 34, 166-176.
- Semizarov, D., Frost, L., Sarthy, A., Kroeger, P., Halbert, D.N., and Fesik, S.W. (2003). Specificity of short interfering RNA determined through gene expression signatures. *Proceedings of the National Academy of Sciences of the United States of America* 100, 6347-6352.
- Sen, G., Wehrman, T.S., Myers, J.W., and Blau, H.M. (2004). Restriction enzyme-generated siRNA (REGS) vectors and libraries. *Nature genetics* 36, 183-189.
- Sertil, O., Kapoor, R., Cohen, B.D., Abramova, N., and Lowry, C.V. (2003). Synergistic repression of anaerobic genes by Mot3 and Rox1 in *Saccharomyces cerevisiae*. *Nucleic acids research* 31, 5831-5837.
- Shabalina, S.A., and Spiridonov, N.A. (2004). The mammalian transcriptome and the function of non-coding DNA sequences. *Genome biology* 5, 105.
- Shannon, M.F., and Rao, S. (2002). Transcription. Of chips and ChIPs. *Science (New York, NY)* 296, 666-669.
- Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research* 13, 2498-2504.
- Shen-Orr, S.S., Milo, R., Mangan, S., and Alon, U. (2002). Network motifs in the transcriptional regulation network of *Escherichia coli*. *Nature genetics* 31, 64-68.
- Shi, Y., Lan, F., Matson, C., Mulligan, P., Whetstine, J.R., Cole, P.A., Casero, R.A., and Shi, Y. (2004). Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119, 941-953.
- Shi, Y.J., Matson, C., Lan, F., Iwase, S., Baba, T., and Shi, Y. (2005). Regulation of LSD1 histone demethylase activity by its associated factors. *Mol Cell* 19, 857-864.
- Shimizu, R., Trainor, C.D., Nishikawa, K., Kobayashi, M., Ohneda, K., and Yamamoto, M. (2007). GATA-1 self-association controls erythroid development in vivo. *The Journal of biological chemistry* 282, 15862-15871.

- Shipp, M.A., Ross, K.N., Tamayo, P., Weng, A.P., Kutok, J.L., Aguiar, R.C., Gaasenbeek, M., Angelo, M., Reich, M., Pinkus, G.S., et al. (2002). Diffuse large B-cell lymphoma outcome prediction by gene-expression profiling and supervised machine learning. *Nature medicine* 8, 68-74.
- Shiraki, T., Kondo, S., Katayama, S., Waki, K., Kasukawa, T., Kawaji, H., Kodzius, R., Watahiki, A., Nakamura, M., Arakawa, T., et al. (2003). Cap analysis gene expression for high-throughput analysis of transcriptional starting point and identification of promoter usage. *Proceedings of the National Academy of Sciences of the United States of America* 100, 15776-15781.
- Shivdasani, R.A., Fujiwara, Y., McDevitt, M.A., and Orkin, S.H. (1997). A lineage-selective knockout establishes the critical role of transcription factor GATA-1 in megakaryocyte growth and platelet development. *Embo J* 16, 3965-3973.
- Shivdasani, R.A., Mayer, E.L., and Orkin, S.H. (1995). Absence of blood formation in mice lacking the T-cell leukaemia oncogene tal-1/SCL. *Nature* 373, 432-434.
- Siddiqui, A.S., Delaney, A.D., Schnerch, A., Griffith, O.L., Jones, S.J., and Marra, M.A. (2006). Sequence biases in large scale gene expression profiling data. *Nucleic acids research* 34, e83.
- Siepel, A., Bejerano, G., Pedersen, J.S., Hinrichs, A.S., Hou, M., Rosenbloom, K., Clawson, H., Spieth, J., Hillier, L.W., Richards, S., et al. (2005). Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes. *Genome research* 15, 1034-1050.
- Sijen, T., Fleenor, J., Simmer, F., Thijssen, K.L., Parrish, S., Timmons, L., Plasterk, R.H., and Fire, A. (2001). On the role of RNA amplification in dsRNA-triggered gene silencing. *Cell* 107, 465-476.
- Silberstein, L., Sanchez, M.J., Socolovsky, M., Liu, Y., Hoffman, G., Kinston, S., Piltz, S., Bowen, M., Gambardella, L., Green, A.R., et al. (2005). Transgenic analysis of the stem cell leukemia +19 stem cell enhancer in adult and embryonic hematopoietic and endothelial cells. *Stem Cells* 23, 1378-1388.
- Silva, J.M., Li, M.Z., Chang, K., Ge, W., Golding, M.C., Rickles, R.J., Siolas, D., Hu, G., Paddison, P.J., Schlabach, M.R., et al. (2005). Second-generation shRNA libraries covering the mouse and human genomes. *Nature genetics* 37, 1281-1288.
- Silva, J.M., Marran, K., Parker, J.S., Silva, J., Golding, M., Schlabach, M.R., Elledge, S.J., Hannon, G.J., and Chang, K. (2008). Profiling essential genes in human mammary cells by multiplex RNAi screening. *Science* (New York, NY) 319, 617-620.
- Silver, L., and Palis, J. (1997). Initiation of murine embryonic erythropoiesis: a spatial analysis. *Blood* 89, 1154-1164.
- Singh-Gasson, S., Green, R.D., Yue, Y., Nelson, C., Blattner, F., Sussman, M.R., and Cerrina, F. (1999). Maskless fabrication of light-directed oligonucleotide microarrays using a digital micromirror array. *Nature biotechnology* 17, 974-978.
- Sledz, C.A., Holko, M., de Veer, M.J., Silverman, R.H., and Williams, B.R. (2003). Activation of the interferon system by short-interfering RNAs. *Nature cell biology* 5, 834-839.
- Smale, S.T., and Kadonaga, J.T. (2003). The RNA polymerase II core promoter. *Annu Rev Biochem* 72, 449-479.
- Smardon, A., Spoerke, J.M., Stacey, S.C., Klein, M.E., Mackin, N., and Maine, E.M. (2000). EGO-1 is related to RNA-directed RNA polymerase and functions in germ-line development and RNA interference in *C. elegans*. *Curr Biol* 10, 169-178.
- Smith, E.M., Gisler, R., and Sigvardsson, M. (2002). Cloning and characterization of a promoter flanking the early B cell factor (EBF) gene indicates roles for E-proteins and autoregulation in the control of EBF expression. *J Immunol* 169, 261-270.
- Song, E., Zhu, P., Lee, S.K., Chowdhury, D., Kussman, S., Dykxhoorn, D.M., Feng, Y., Palliser, D., Weiner, D.B., Shankar, P., et al. (2005). Antibody mediated in vivo delivery of small interfering RNAs via cell-surface receptors. *Nature biotechnology* 23, 709-717.
- Song, S.H., Hou, C., and Dean, A. (2007). A positive role for NLI/Ldb1 in long-range beta-globin locus control region function. *Mol Cell* 28, 810-822.
- Sonnichsen, B., Koski, L.B., Walsh, A., Marschall, P., Neumann, B., Brehm, M., Alleaume, A.M., Artelt, J., Bettencourt, P., Cassin, E., et al. (2005). Full-genome RNAi profiling of early embryogenesis in *Caenorhabditis elegans*. *Nature* 434, 462-469.

- Spicuglia, S., Kumar, S., Yeh, J.H., Vachez, E., Chasson, L., Gorbach, S., Cautres, J., and Ferrier, P. (2002). Promoter activation by enhancer-dependent and -independent loading of activator and coactivator complexes. *Mol Cell* 10, 1479-1487.
- Spilianakis, C.G., Lalioti, M.D., Town, T., Lee, G.R., and Flavell, R.A. (2005). Interchromosomal associations between alternatively expressed loci. *Nature* 435, 637-645.
- Stewart, S.A., Dykxhoorn, D.M., Palliser, D., Mizuno, H., Yu, E.Y., An, D.S., Sabatini, D.M., Chen, I.S., Hahn, W.C., Sharp, P.A., et al. (2003). Lentivirus-delivered stable gene silencing by RNAi in primary cells. *RNA* (New York, NY) 9, 493-501.
- Stielow, B., Sapetschnig, A., Kruger, I., Kunert, N., Brehm, A., Boutros, M., and Suske, G. (2008). Identification of SUMO-dependent chromatin-associated transcriptional repression components by a genome-wide RNAi screen. *Mol Cell* 29, 742-754.
- Stopka, T., Amanatullah, D.F., Papetti, M., and Skoultchi, A.I. (2005). PU.1 inhibits the erythroid program by binding to GATA-1 on DNA and creating a repressive chromatin structure. *Embo J* 24, 3712-3723.
- Stopka T, Skoultchi AI. The ISWI ATPase Snf2h is required for early mouse development. (2003). *Proc Natl Acad Sci U S A*. 100, 14097-102.
- Stopka T, Zakova D, Fuchs O, Kubrova O, Blafkova J, Jelinek J, Necas E, Zivny J. (2000). Chromatin remodeling gene SMARCA5 is dysregulated in primitive hematopoietic cells of acute leukemia. *Leukemia* 14, 1247-1252.
- Studitsky, V.M., Clark, D.J., and Felsenfeld, G. (1994). A histone octamer can step around a transcribing polymerase without leaving the template. *Cell* 76, 371-382.
- Sun, X.H., and Baltimore, D. (1991). An inhibitory domain of E12 transcription factor prevents DNA binding in E12 homodimers but not in E12 heterodimers. *Cell* 64, 459-470.
- Suzuki, Y., Tsunoda, T., Sese, J., Taira, H., Mizushima-Sugano, J., Hata, H., Ota, T., Isogai, T., Tanaka, T., Nakamura, Y., et al. (2001). Identification and characterization of the potential promoter regions of 1031 kinds of human genes. *Genome research* 11, 677-684.
- Swiers, G., Patient, R., and Loose, M. (2006). Genetic regulatory networks programming hematopoietic stem cells and erythroid lineage specification. *Developmental biology* 294, 525-540.
- Szekvolgyi, L., Balint, B.L., Imre, L., Goda, K., Szabo, M., Nagy, L., and Szabo, G. (2006). Chip-on-beads: flow-cytometric evaluation of chromatin immunoprecipitation. *Cytometry A* 69, 1086-1091.
- Tabara, H., Grishok, A., and Mello, C.C. (1998). RNAi in *C. elegans*: soaking in the genome sequence. *Science* (New York, NY) 282, 430-431.
- Tavazoie, S., Hughes, J.D., Campbell, M.J., Cho, R.J., and Church, G.M. (1999). Systematic determination of genetic network architecture. *Nature genetics* 22, 281-285.
- Tavernarakis, N., Wang, S.L., Dorovkov, M., Ryazanov, A., and Driscoll, M. (2000). Heritable and inducible genetic interference by double-stranded RNA encoded by transgenes. *Nature genetics* 24, 180-183.
- Thomson, S., Clayton, A.L., Hazzalin, C.A., Rose, S., Barratt, M.J., and Mahadevan, L.C. (1999). The nucleosomal response associated with immediate-early gene induction is mediated via alternative MAP kinase cascades: MSK1 as a potential histone H3/HMG-14 kinase. *Embo J* 18, 4779-4793.
- Thut, C.J., Chen, J.L., Klemm, R., and Tjian, R. (1995). p53 transcriptional activation mediated by coactivators TAFII40 and TAFII60. *Science* (New York, NY) 267, 100-104.
- Timmons, L., Court, D.L., and Fire, A. (2001). Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene* 263, 103-112.
- Timmons, L., and Fire, A. (1998). Specific interference by ingested dsRNA. *Nature* 395, 854.
- Torres, T.T., Metta, M., Ottenwalder, B., and Schlotterer, C. (2008). Gene expression profiling by massively parallel sequencing. *Genome research* 18, 172-177.
- Tsai, F.Y., and Orkin, S.H. (1997). Transcription factor GATA-2 is required for proliferation/survival of early hematopoietic cells and mast cell formation, but not for erythroid and myeloid terminal differentiation. *Blood* 89, 3636-3643.
- Tsai, S.F., Martin, D.I., Zon, L.I., D'Andrea, A.D., Wong, G.G., and Orkin, S.H. (1989). Cloning of cDNA for the major DNA-binding protein of the erythroid lineage through expression in mammalian cells. *Nature* 339, 446-451.

- Tsang, A.P., Visvader, J.E., Turner, C.A., Fujiwara, Y., Yu, C., Weiss, M.J., Crossley, M., and Orkin, S.H. (1997). FOG, a multitype zinc finger protein, acts as a cofactor for transcription factor GATA-1 in erythroid and megakaryocytic differentiation. *Cell* 90, 109-119.
- Tse, C., Sera, T., Wolffe, A.P., and Hansen, J.C. (1998). Disruption of higher-order folding by core histone acetylation dramatically enhances transcription of nucleosomal arrays by RNA polymerase III. *Molecular and cellular biology* 18, 4629-4638.
- Tuerk, C., and Gold, L. (1990). Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science (New York, NY)* 249, 505-510.
- Tusher, V.G., Tibshirani, R., and Chu, G. (2001). Significance analysis of microarrays applied to the ionizing radiation response. *Proceedings of the National Academy of Sciences of the United States of America* 98, 5116-5121.
- Utley, R.T., Ikeda, K., Grant, P.A., Cote, J., Steger, D.J., Eberharter, A., John, S., and Workman, J.L. (1998). Transcriptional activators direct histone acetyltransferase complexes to nucleosomes. *Nature* 394, 498-502.
- Valverde-Garduno, V., Guyot, B., Anguita, E., Hamlett, I., Porcher, C., and Vyas, P. (2004). Differences in the chromatin structure and cis-element organization of the human and mouse GATA1 loci: implications for cis-element identification. *Blood* 104, 3106-3116.
- van de Wetering, M., Oving, I., Muncan, V., Pon Fong, M.T., Brantjes, H., van Leenen, D., Holstege, F.C., Brummelkamp, T.R., Agami, R., and Clevers, H. (2003). Specific inhibition of gene expression using a stably integrated, inducible small-interfering-RNA vector. *EMBO reports* 4, 609-615.
- van Eekelen, J.A., Bradley, C.K., Gothert, J.R., Robb, L., Elefanty, A.G., Begley, C.G., and Harvey, A.R. (2003). Expression pattern of the stem cell leukaemia gene in the CNS of the embryonic and adult mouse. *Neuroscience* 122, 421-436.
- van Haaften, G., Romeijn, R., Pothof, J., Koole, W., Mullenders, L.H., Pastink, A., Plasterk, R.H., and Tijsterman, M. (2006). Identification of conserved pathways of DNA-damage response and radiation protection by genome-wide RNAi. *Curr Biol* 16, 1344-1350.
- van Steensel, B., and Henikoff, S. (2000). Identification of in vivo DNA targets of chromatin proteins using tethered dam methyltransferase. *Nature biotechnology* 18, 424-428.
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A., and Speleman, F. (2002). Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome biology* 3, RESEARCH0034.
- Velculescu, V.E., Zhang, L., Vogelstein, B., and Kinzler, K.W. (1995). Serial analysis of gene expression. *Science (New York, NY)* 270, 484-487.
- Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., Smith, H.O., Yandell, M., Evans, C.A., Holt, R.A., et al. (2001). The sequence of the human genome. *Science (New York, NY)* 291, 1304-1351.
- Vernimmen, D., De Gobbi, M., Sloane-Stanley, J.A., Wood, W.G., and Higgs, D.R. (2007). Long-range chromosomal interactions regulate the timing of the transition between poised and active gene expression. *Embo J* 26, 2041-2051.
- Viens, A., Mechold, U., Lehrmann, H., Harel-Bellan, A., and Ogryzko, V. (2004). Use of protein biotinylation in vivo for chromatin immunoprecipitation. *Analytical biochemistry* 325, 68-76.
- Vilar, J.M., and Saiz, L. (2005). DNA looping in gene regulation: from the assembly of macromolecular complexes to the control of transcriptional noise. *Current opinion in genetics & development* 15, 136-144.
- Visel, A., Minovitsky, S., Dubchak, I., and Pennacchio, L.A. (2007). VISTA Enhancer Browser--a database of tissue-specific human enhancers. *Nucleic acids research* 35, D88-92.
- Visvader, J., Begley, C.G., and Adams, J.M. (1991). Differential expression of the LYL, SCL and E2A helix-loop-helix genes within the hemopoietic system. *Oncogene* 6, 187-194.
- Visvader, J.E., Fujiwara, Y., and Orkin, S.H. (1998). Unsuspected role for the T-cell leukemia protein SCL/tal-1 in vascular development. *Genes & development* 12, 473-479.
- Visvader, J.E., Mao, X., Fujiwara, Y., Hahm, K., and Orkin, S.H. (1997). The LIM-domain binding protein Ldb1 and its partner LMO2 act as negative regulators of erythroid differentiation. *Proceedings of the National Academy of Sciences of the United States of America* 94, 13707-13712.

- Vitelli, L., Condorelli, G., Lulli, V., Hoang, T., Luchetti, L., Croce, C.M., and Peschle, C. (2000). A pentamer transcriptional complex including tal-1 and retinoblastoma protein downmodulates c-kit expression in normal erythroblasts. *Molecular and cellular biology* 20, 5330-5342.
- Vogel, M.J., Guelen, L., de Wit, E., Peric-Hupkes, D., Loden, M., Talhout, W., Feenstra, M., Abbas, B., Classen, A.K., and van Steensel, B. (2006). Human heterochromatin proteins form large domains containing KRAB-ZNF genes. *Genome research* 16, 1493-1504.
- Vogel, M.J., Peric-Hupkes, D., and van Steensel, B. (2007). Detection of in vivo protein-DNA interactions using DamID in mammalian cells. *Nature protocols* 2, 1467-1478.
- Vokes, S.A., Ji, H., McCuine, S., Tenzen, T., Giles, S., Zhong, S., Longabaugh, W.J., Davidson, E.H., Wong, W.H., and McMahon, A.P. (2007). Genomic characterization of Gli-activator targets in sonic hedgehog-mediated neural patterning. *Development (Cambridge, England)* 134, 1977-1989.
- Vyas, P., Ault, K., Jackson, C.W., Orkin, S.H., and Shivdasani, R.A. (1999). Consequences of GATA-1 deficiency in megakaryocytes and platelets. *Blood* 93, 2867-2875.
- Vyas, P., McDevitt, M.A., Cantor, A.B., Katz, S.G., Fujiwara, Y., and Orkin, S.H. (1999). Different sequence requirements for expression in erythroid and megakaryocytic cells within a regulatory element upstream of the GATA-1 gene. *Development (Cambridge, England)* 126, 2799-2811.
- Wadman, I., Li, J., Bash, R.O., Forster, A., Osada, H., Rabbitts, T.H., and Baer, R. (1994). Specific in vivo association between the bHLH and LIM proteins implicated in human T cell leukemia. *Embo J* 13, 4831-4839.
- Wadman, I.A., Osada, H., Grutz, G.G., Agulnick, A.D., Westphal, H., Forster, A., and Rabbitts, T.H. (1997). The LIM-only protein Lmo2 is a bridging molecule assembling an erythroid, DNA-binding complex which includes the TAL1, E47, GATA-1 and Ldb1/NLI proteins. *Embo J* 16, 3145-3157.
- Wakabayashi, J., Yomogida, K., Nakajima, O., Yoh, K., Takahashi, S., Engel, J.D., Ohneda, K., and Yamamoto, M. (2003). GATA-1 testis activation region is essential for Sertoli cell-specific expression of GATA-1 gene in transgenic mouse. *Genes Cells* 8, 619-630.
- Walker, G.T. (1993). Empirical aspects of strand displacement amplification. *PCR methods and applications* 3, 1-6.
- Wall, L., deBoer, E., and Grosveld, F. (1988). The human beta-globin gene 3' enhancer contains multiple binding sites for an erythroid-specific protein. *Genes & development* 2, 1089-1100.
- Wang, H., Wang, L., Erdjument-Bromage, H., Vidal, M., Tempst, P., Jones, R.S., and Zhang, Y. (2004). Role of histone H2A ubiquitination in Polycomb silencing. *Nature* 431, 873-878.
- Wang, J., Hoshino, T., Redner, R.L., Kajigaya, S., and Liu, J.M. (1998). ETO, fusion partner in t(8;21) acute myeloid leukemia, represses transcription by interaction with the human N-CoR/mSin3/HDAC1 complex. *Proceedings of the National Academy of Sciences of the United States of America* 95, 10860-10865.
- Wang, Y., Wysocka, J., Sayegh, J., Lee, Y.H., Perlin, J.R., Leonelli, L., Sonbuchner, L.S., McDonald, C.H., Cook, R.G., Dou, Y., et al. (2004). Human PAD4 regulates histone arginine methylation levels via demethylimination. *Science (New York, NY)* 306, 279-283.
- Warren, A.J., Colledge, W.H., Carlton, M.B., Evans, M.J., Smith, A.J., and Rabbitts, T.H. (1994). The oncogenic cysteine-rich LIM domain protein rbtn2 is essential for erythroid development. *Cell* 78, 45-57.
- Wasserman, W.W., and Fickett, J.W. (1998). Identification of regulatory regions which confer muscle-specific gene expression. *Journal of molecular biology* 278, 167-181.
- Waterhouse, P.M., Wang, M.B., and Lough, T. (2001). Gene silencing as an adaptive defence against viruses. *Nature* 411, 834-842.
- Weber, F., de Villiers, J., and Schaffner, W. (1984). An SV40 "enhancer trap" incorporates exogenous enhancers or generates enhancers from its own sequences. *Cell* 36, 983-992.
- Wechsler, J., Greene, M., McDevitt, M.A., Anastasi, J., Karp, J.E., Le Beau, M.M., and Crispino, J.D. (2002). Acquired mutations in GATA1 in the megakaryoblastic leukemia of Down syndrome. *Nature genetics* 32, 148-152.
- Wei, C.L., Wu, Q., Vega, V.B., Chiu, K.P., Ng, P., Zhang, T., Shahab, A., Yong, H.C., Fu, Y., Weng, Z., et al. (2006). A global map of p53 transcription-factor binding sites in the human genome. *Cell* 124, 207-219.

- Weinberg, M.S., Villeneuve, L.M., Ehsani, A., Amarzguioui, M., Aagaard, L., Chen, Z.X., Riggs, A.D., Rossi, J.J., and Morris, K.V. (2006). The antisense strand of small interfering RNAs directs histone methylation and transcriptional gene silencing in human cells. *RNA (New York, NY)* 12, 256-262.
- Weinmann, A.S., Bartley, S.M., Zhang, T., Zhang, M.Q., and Farnham, P.J. (2001). Use of chromatin immunoprecipitation to clone novel E2F target promoters. *Molecular and cellular biology* 21, 6820-6832.
- Weinmann, A.S., Yan, P.S., Oberley, M.J., Huang, T.H., and Farnham, P.J. (2002). Isolating human transcription factor targets by coupling chromatin immunoprecipitation and CpG island microarray analysis. *Genes & development* 16, 235-244.
- Weiss, M.J., Keller, G., and Orkin, S.H. (1994). Novel insights into erythroid development revealed through in vitro differentiation of GATA-1 embryonic stem cells. *Genes & development* 8, 1184-1197.
- Weiss, M.J., and Orkin, S.H. (1995). Transcription factor GATA-1 permits survival and maturation of erythroid precursors by preventing apoptosis. *Proceedings of the National Academy of Sciences of the United States of America* 92, 9623-9627.
- Welch, J.J., Watts, J.A., Vakoc, C.R., Yao, Y., Wang, H., Hardison, R.C., Blobel, G.A., Chodosh, L.A., and Weiss, M.J. (2004). Global regulation of erythroid gene expression by transcription factor GATA-1. *Blood* 104, 3136-3147.
- Wells, J., Yan, P.S., Cechvala, M., Huang, T., and Farnham, P.J. (2003). Identification of novel pRb binding sites using CpG microarrays suggests that E2F recruits pRb to specific genomic sites during S phase. *Oncogene* 22, 1445-1460.
- West, A.G., Gaszner, M., and Felsenfeld, G. (2002). Insulators: many functions, many mechanisms. *Genes & development* 16, 271-288.
- Westbrook, T.F., Martin, E.S., Schlabach, M.R., Leng, Y., Liang, A.C., Feng, B., Zhao, J.J., Roberts, T.M., Mandel, G., Hannon, G.J., et al. (2005). A genetic screen for candidate tumor suppressors identifies REST. *Cell* 121, 837-848.
- White, K.P., Rifkin, S.A., Hurban, P., and Hogness, D.S. (1999). Microarray analysis of *Drosophila* development during metamorphosis. *Science (New York, NY)* 286, 2179-2184.
- Wightman, B., Ha, I., and Ruvkun, G. (1993). Posttranscriptional regulation of the heterochronic gene *lin-14* by *lin-4* mediates temporal pattern formation in *C. elegans*. *Cell* 75, 855-862.
- Williams, A.O., Isaacs, R.J., and Stowell, K.M. (2007). Down-regulation of human topoisomerase IIalpha expression correlates with relative amounts of specificity factors Sp1 and Sp3 bound at proximal and distal promoter regions. *BMC molecular biology* 8, 36.
- Williams, B.R. (1997). Role of the double-stranded RNA-activated protein kinase (PKR) in cell regulation. *Biochem Soc Trans* 25, 509-513.
- Williams, R.R., Azuara, V., Perry, P., Sauer, S., Dvorkina, M., Jorgensen, H., Roix, J., McQueen, P., Misteli, T., Merkenschlager, M., et al. (2006). Neural induction promotes large-scale chromatin reorganisation of the *Mash1* locus. *Journal of cell science* 119, 132-140.
- Wolfrum, C., Shi, S., Jayaprakash, K.N., Jayaraman, M., Wang, G., Pandey, R.K., Rajeev, K.G., Nakayama, T., Charrise, K., Ndungo, E.M., et al. (2007). Mechanisms and optimization of in vivo delivery of lipophilic siRNAs. *Nature biotechnology* 25, 1149-1157.
- Wolstein, O., Silkov, A., Revach, M., and Dikstein, R. (2000). Specific interaction of TAFII105 with OCA-B is involved in activation of octamer-dependent transcription. *The Journal of biological chemistry* 275, 16459-16465.
- Wong, J.M., and Bateman, E. (1994). TBP-DNA interactions in the minor groove discriminate between A:T and T:A base pairs. *Nucleic acids research* 22, 1890-1896.
- Wood, A., Krogan, N.J., Dover, J., Schneider, J., Heidt, J., Boateng, M.A., Dean, K., Golshani, A., Zhang, Y., Greenblatt, J.F., et al. (2003). Bre1, an E3 ubiquitin ligase required for recruitment and substrate selection of Rad6 at a promoter. *Mol Cell* 11, 267-274.
- Woolfe, A., Goodson, M., Goode, D.K., Snell, P., McEwen, G.K., Vavouri, T., Smith, S.F., North, P., Callaway, H., Kelly, K., et al. (2005). Highly conserved non-coding sequences are associated with vertebrate development. *PLoS biology* 3, e7.
- Wright, W.E., Binder, M., and Funk, W. (1991). Cyclic amplification and selection of targets (CASTing) for the myogenin consensus binding site. *Molecular and cellular biology* 11, 4104-4110.

- Wu, C., Wong, Y.C., and Elgin, S.C. (1979). The chromatin structure of specific genes: II. Disruption of chromatin structure during gene activity. *Cell* 16, 807-814.
- Wu, Z., and Irizarry, R.A. (2005). Stochastic models inspired by hybridization theory for short oligonucleotide arrays. *J Comput Biol* 12, 882-893.
- Wyrick, J.J., Aparicio, J.G., Chen, T., Barnett, J.D., Jennings, E.G., Young, R.A., Bell, S.P., and Aparicio, O.M. (2001). Genome-wide distribution of ORC and MCM proteins in *S. cerevisiae*: high-resolution mapping of replication origins. *Science (New York, NY)* 294, 2357-2360.
- Xu, Z., Huang, S., Chang, L.S., Agulnick, A.D., and Brandt, S.J. (2003). Identification of a TAL1 target gene reveals a positive role for the LIM domain-binding protein Ldb1 in erythroid gene expression and differentiation. *Molecular and cellular biology* 23, 7585-7599.
- Xu, Z., Meng, X., Cai, Y., Koury, M.J., and Brandt, S.J. (2006). Recruitment of the SWI/SNF protein Brg1 by a multiprotein complex effects transcriptional repression in murine erythroid progenitors. *The Biochemical journal* 399, 297-304.
- Yamada, Y., Pannell, R., Forster, A., and Rabbitts, T.H. (2000). The oncogenic LIM-only transcription factor Lmo2 regulates angiogenesis but not vasculogenesis in mice. *Proceedings of the National Academy of Sciences of the United States of America* 97, 320-324.
- Yamada, Y., Warren, A.J., Dobson, C., Forster, A., Pannell, R., and Rabbitts, T.H. (1998). The T cell leukemia LIM protein Lmo2 is necessary for adult mouse hematopoiesis. *Proceedings of the National Academy of Sciences of the United States of America* 95, 3890-3895.
- Yamaguchi, Y., Ackerman, S.J., Minegishi, N., Takiguchi, M., Yamamoto, M., and Suda, T. (1998). Mechanisms of transcription in eosinophils: GATA-1, but not GATA-2, transactivates the promoter of the eosinophil granule major basic protein gene. *Blood* 91, 3447-3458.
- Yamaguchi, Y., Nishio, H., Kishi, K., Ackerman, S.J., and Suda, T. (1999). C/EBPbeta and GATA-1 synergistically regulate activity of the eosinophil granule major basic protein promoter: implication for C/EBPbeta activity in eosinophil gene expression. *Blood* 94, 1429-1439.
- Yang, D., Buchholz, F., Huang, Z., Goga, A., Chen, C.Y., Brodsky, F.M., and Bishop, J.M. (2002). Short RNA duplexes produced by hydrolysis with *Escherichia coli* RNase III mediate effective RNA interference in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America* 99, 9942-9947.
- Yi, Z.C., Wang, Z., Li, H.X., Liu, M.J., Wu, R.C., and Wang, X.H. (2004). Effects of chebulinic acid on differentiation of human leukemia K562 cells. *Acta pharmacologica Sinica* 25, 231-238.
- Yoder, M.C., Hiatt, K., and Mukherjee, P. (1997). In vivo repopulating hematopoietic stem cells are present in the murine yolk sac at day 9.0 postcoitus. *Proceedings of the National Academy of Sciences of the United States of America* 94, 6776-6780.
- Yokomizo, T., Takahashi, S., Mochizuki, N., Kuroha, T., Ema, M., Wakamatsu, A., Shimizu, R., Ohneda, O., Osato, M., Okada, H., et al. (2007). Characterization of GATA-1(+) hemangioblastic cells in the mouse embryo. *Embo J* 26, 184-196.
- Yomogida, K., Ohtani, H., Harigae, H., Ito, E., Nishimune, Y., Engel, J.D., and Yamamoto, M. (1994). Developmental stage- and spermatogenic cycle-specific expression of transcription factor GATA-1 in mouse Sertoli cells. *Development (Cambridge, England)* 120, 1759-1766.
- Yoo, J., Herman, L.E., Li, C., Krantz, S.B., and Tuan, D. (1996). Dynamic changes in the locus control region of erythroid progenitor cells demonstrated by polymerase chain reaction. *Blood* 87, 2558-2567.
- Yoo, J.Y., Kim, J.H., Kim, J., Huang, J.H., Zhang, S.N., Kang, Y.A., Kim, H., and Yun, C.O. (2008). Short hairpin RNA-expressing oncolytic adenovirus-mediated inhibition of IL-8: effects on antiangiogenesis and tumor growth inhibition. *Gene therapy* 15, 635-651.
- Yu, C., Cantor, A.B., Yang, H., Browne, C., Wells, R.A., Fujiwara, Y., and Orkin, S.H. (2002). Targeted deletion of a high-affinity GATA-binding site in the GATA-1 promoter leads to selective loss of the eosinophil lineage in vivo. *The Journal of experimental medicine* 195, 1387-1395.
- Yu, C., Niakan, K.K., Matsushita, M., Stamatoyannopoulos, G., Orkin, S.H., and Raskind, W.H. (2002). X-linked thrombocytopenia with thalassemia from a mutation in the amino finger of GATA-1 affecting DNA binding rather than FOG-1 interaction. *Blood* 100, 2040-2045.

- Zeller, K.I., Zhao, X., Lee, C.W., Chiu, K.P., Yao, F., Yustein, J.T., Ooi, H.S., Orlov, Y.L., Shahab, A., Yong, H.C., et al. (2006). Global mapping of c-Myc binding sites and target gene networks in human B cells. *Proceedings of the National Academy of Sciences of the United States of America* 103, 17834-17839.
- Zeng, P.Y., Vakoc, C.R., Chen, Z.C., Blobel, G.A., and Berger, S.L. (2006). In vivo dual cross-linking for identification of indirect DNA-associated proteins by chromatin immunoprecipitation. *BioTechniques* 41, 694, 696, 698.
- Zeng, Y., Wagner, E.J., and Cullen, B.R. (2002). Both natural and designed micro RNAs can inhibit the expression of cognate mRNAs when expressed in human cells. *Mol Cell* 9, 1327-1333.
- Zenzie-Gregory, B., Khachi, A., Garraway, I.P., and Smale, S.T. (1993). Mechanism of initiator-mediated transcription: evidence for a functional interaction between the TATA-binding protein and DNA in the absence of a specific recognition sequence. *Molecular and cellular biology* 13, 3841-3849.
- Zhang, P., Zhang, X., Iwama, A., Yu, C., Smith, K.A., Mueller, B.U., Narravula, S., Torbett, B.E., Orkin, S.H., and Tenen, D.G. (2000). PU.1 inhibits GATA-1 function and erythroid differentiation by blocking GATA-1 DNA binding. *Blood* 96, 2641-2648.
- Zhang, W., Yang, H., Kong, X., Mohapatra, S., San Juan-Vergara, H., Hellermann, G., Behera, S., Singam, R., Lockey, R.F., and Mohapatra, S.S. (2005). Inhibition of respiratory syncytial virus infection with intranasal siRNA nanoparticles targeting the viral NS1 gene. *Nature medicine* 11, 56-62.
- Zhang, Y., Payne, K.J., Zhu, Y., Price, M.A., Parrish, Y.K., Zielinska, E., Barsky, L.W., and Crooks, G.M. (2005). SCL expression at critical points in human hematopoietic lineage commitment. *Stem Cells* 23, 852-860.
- Zhao, H., and Dean, A. (2004). An insulator blocks spreading of histone acetylation and interferes with RNA polymerase II transfer between an enhancer and gene. *Nucleic acids research* 32, 4903-4919.
- Zhao, Z., Tavoosidana, G., Sjolinder, M., Gondor, A., Mariano, P., Wang, S., Kanduri, C., Lezcano, M., Sandhu, K.S., Singh, U., et al. (2006). Circular chromosome conformation capture (4C) uncovers extensive networks of epigenetically regulated intra- and interchromosomal interactions. *Nature genetics* 38, 1341-1347.
- Zhu, B., Zheng, Y., Pham, A.D., Mandal, S.S., Erdjument-Bromage, H., Tempst, P., and Reinberg, D. (2005). Monoubiquitination of human histone H2B: the factors involved and their roles in HOX gene regulation. *Mol Cell* 20, 601-611.
- Zhu, P., Zhou, W., Wang, J., Puc, J., Ohgi, K.A., Erdjument-Bromage, H., Tempst, P., Glass, C.K., and Rosenfeld, M.G. (2007). A histone H2A deubiquitinase complex coordinating histone acetylation and H1 dissociation in transcriptional regulation. *Mol Cell* 27, 609-621.
- Zhuang, Y., Jackson, A., Pan, L., Shen, K., and Dai, M. (2004). Regulation of E2A gene expression in B-lymphocyte development. *Molecular immunology* 40, 1165-1177.
- Zhuang, Y., Soriano, P., and Weintraub, H. (1994). The helix-loop-helix gene E2A is required for B cell formation. *Cell* 79, 875-884.
- Zimmermann, T.S., Lee, A.C., Akinc, A., Bramlage, B., Bumcrot, D., Fedoruk, M.N., Harborth, J., Heyes, J.A., Jeffs, L.B., John, M., et al. (2006). RNAi-mediated gene silencing in non-human primates. *Nature* 441, 111-114.
- Zon, L.I., Tsai, S.F., Burgess, S., Matsudaira, P., Bruns, G.A., and Orkin, S.H. (1990). The major human erythroid DNA-binding protein (GF-1): primary sequence and localization of the gene to the X chromosome. *Proceedings of the National Academy of Sciences of the United States of America* 87, 668-672.
- Zon, L.I., Yamaguchi, Y., Yee, K., Albee, E.A., Kimura, A., Bennett, J.C., Orkin, S.H., and Ackerman, S.J. (1993). Expression of mRNA for the GATA-binding proteins in human eosinophils and basophils: potential role in gene transcription. *Blood* 81, 3234-3241.
- Zon, L.I., Youssoufian, H., Mather, C., Lodish, H.F., and Orkin, S.H. (1991). Activation of the erythropoietin receptor promoter by transcription factor GATA-1. *Proceedings of the National Academy of Sciences of the United States of America* 88, 10638-10641.
- Zweidler-Mckay, P.A., Grimes, H.L., Flubacher, M.M., and Tschlis, P.N. (1996). Gfi-1 encodes a nuclear zinc finger protein that binds DNA and functions as a transcriptional repressor. *Molecular and cellular biology* 16, 4024-4034.