

REFERENCES

Aasen T, Raya A, Barrero MJ, Garreta E, Consiglio A, Gonzalez F, Vassena R, Bilic J, Pekarik V, Tiscornia G et al: Efficient and rapid generation of induced pluripotent stem cells from human keratinocytes. *Nat Biotechnol* 2008, 26(11):1276-1284.

Abdelrahman B, Fiddler M, Rappolee D, Pergament E: Expression of Transcription Regulating Genes in Human Preimplantation Embryos. *Hum Reprod* 1995, 10(10):2787-2792.

Agarwal S, Loh YH, McLoughlin EM, Huang J, Park IH, Miller JD, Huo H, Okuka M, Dos Reis RM, Loewer S et al: Telomere elongation in induced pluripotent stem cells from dyskeratosis congenita patients. *Nature* 2010, 464(7286):292-296.

Aiuti A, Cattaneo F, Galimberti S, Benninghoff U, Cassani B, Callegaro L, Scaramuzza S, Andolfi G, Mirolo M, Brigida I et al: Gene Therapy for Immunodeficiency Due to Adenosine Deaminase Deficiency. *New Engl J Med* 2009, 360(5):447-458.

Akagi T, Ito T, Kato M, Jin Z, Cheng Y, Kan T, Yamamoto G, Oлару A, Kawamata N, Boulton J et al: Chromosomal abnormalities and novel disease-related regions in progression from Barrett's esophagus to esophageal adenocarcinoma. *International journal of cancer Journal international du cancer* 2009, 125(10):2349-2359.

Albright CD, Salganik RI, Van Dyke T: Dietary depletion of vitamin E and vitamin A inhibits mammary tumor growth and metastasis in transgenic mice. *J Nutr* 2004, 134(5):1139-1144.

Alsayed Y, Uddin S, Mahmud N, Lekmine F, Kalvakolanu DV, Minucci S, Bokoch G, Plataniotis LC: Activation of Rac1 and the p38 mitogen-activated protein kinase pathway in response to all-trans-retinoic acid. *The Journal of biological chemistry* 2001, 276(6):4012-4019.

Ampas K, Andrews PW, Anyfantis G, Armstrong L, Avery S, Baharvand H, Baker J, Baker D, Munoz MB, Beil S et al: Screening ethnically diverse human embryonic stem

cells identifies a chromosome 20 minimal amplicon conferring growth advantage. *Nat Biotechnol* 2011, 29(12):1132-1144.

Amsterdam A, Nissen RM, Sun Z, Swindell EC, Farrington S, Hopkins N: Identification of 315 genes essential for early zebrafish development. *P Natl Acad Sci USA* 2004, 101(35):12792-12797.

An MC, Zhang NZ, Scott G, Montoro D, Wittkop T, Mooney S, Melov S, Ellerby LM: Genetic Correction of Huntington's Disease Phenotypes in Induced Pluripotent Stem Cells. *Cell stem cell* 2012, 11(2):253-263.

Andre E, Gawlas K, Becker-Andre M: A novel isoform of the orphan nuclear receptor RORbeta is specifically expressed in pineal gland and retina. *Gene* 1998, 216(2):277-283.

Ang HL, Duester G: Initiation of retinoid signaling in primitive streak mouse embryos: spatiotemporal expression patterns of receptors and metabolic enzymes for ligand synthesis. *Dev Dyn* 1997, 208(4):536-543.

Anokye-Danso F, Trivedi CM, Jühr D, Gupta M, Cui Z, Tian Y, Zhang Y, Yang W, Gruber PJ, Epstein JA et al: Highly efficient miRNA-mediated reprogramming of mouse and human somatic cells to pluripotency. *Cell stem cell* 2011, 8(4):376-388.

Aoi T, Yae K, Nakagawa M, Ichisaka T, Okita K, Takahashi K, Chiba T, Yamanaka S: Generation of pluripotent stem cells from adult mouse liver and stomach cells. *Science* 2008, 321(5889):699-702.

Aristotle: *Generation of Animals*. Cambridge, MA: Harvard University Press; 1979.

Avilion AA, Nicolis SK, Pevny LH, Perez L, Vivian N, Lovell-Badge R: Multipotent cell lineages in early mouse development depend on SOX2 function. *Gene Dev* 2003, 17(1):126-140.

Balciunas D, Wangensteen KJ, Wilber A, Bell J, Geurts A, Sivasubbu S, Wang X, Hackett PB, Largaespada DA, McIvor RS et al: Harnessing a high cargo-capacity transposon for genetic applications in vertebrates. *PLoS genetics* 2006, 2(11):e169.

Balmer JE, Blomhoff R: Gene expression regulation by retinoic acid. *Journal of lipid research* 2002, 43(11):1773-1808.

Bao S, Tang F, Li X, Hayashi K, Gillich A, Lao K, Surani MA: Epigenetic reversion of post-implantation epiblast to pluripotent embryonic stem cells. *Nature* 2009, 461(7268):1292-1295.

Bard-Chapeau EA, Jeyakani J, Kok CH, Muller J, Chua BQ, Gunaratne J, Batagov A, Jenjaroenpun P, Kuznetsov VA, Wei CL et al: Ecotopic viral integration site 1 (EV1) regulates multiple cellular processes important for cancer and is a synergistic partner for FOS protein in invasive tumors. *P Natl Acad Sci USA* 2012, 109(6):2168-2173.

Barnea E, Bergman Y: Synergy of SF1 and RAR in activation of Oct-3/4 promoter. *The Journal of biological chemistry* 2000, 275(9):6608-6619.

Bastien J, Adam-Stitah S, Riedl T, Egly JM, Chambon P, Rochette-Egly C: TFIIF interacts with the retinoic acid receptor gamma and phosphorylates its AF-1-activating domain through cdk7. *The Journal of biological chemistry* 2000, 275(29):21896-21904.

Beattie GM, Lopez AD, Bucay N, Hinton A, Firpo MT, King CC, Hayek A: Activin A maintains pluripotency of human embryonic stem cells in the absence of feeder layers. *Stem Cells* 2005, 23(4):489-495.

Becker M, Becker A, Miyara F, Han Z, Kihara M, Brown DT, Hager GL, Latham K, Adashi EY, Misteli T: Differential in vivo binding dynamics of somatic and oocyte-specific linker histones in oocytes and during ES cell nuclear transfer. *Molecular biology of the cell* 2005, 16(8):3887-3895.

Becker-Andre M, Andre E, DeLamarter JF: Identification of nuclear receptor mRNAs by RT-PCR amplification of conserved zinc-finger motif sequences. *Biochem Biophys Res Commun* 1993, 194(3):1371-1379.

Belay E, Dastidar S, VandenDriessche T, Chuah MKL: Transposon-Mediated Gene Transfer into Adult and Induced Pluripotent Stem Cells. *Curr Gene Ther* 2011, 11(5):406-413.

Benbrook D, Lernhardt E, Pfahl M: A new retinoic acid receptor identified from a hepatocellular carcinoma. *Nature* 1988, 333(6174):669-672.

Bendall AJ, Sturm RA, Danoy PA, Molloy PL: Broad binding-site specificity and affinity properties of octamer 1 and brain octamer-binding proteins. *European journal of biochemistry / FEBS* 1993, 217(3):799-811.

Ben-Shushan E, Sharir H, Pikarsky E, Bergman Y: A dynamic balance between ARP-1/COUP-TFII, EAR-3/COUP-TFI, and retinoic acid receptor:retinoid X receptor heterodimers regulates Oct-3/4 expression in embryonal carcinoma cells. *Mol Cell Biol* 1995, 15(2):1034-1048.

Bessereau JL, Wright A, Williams DC, Schuske K, Davis MW, Jorgensen EM: Mobilization of a *Drosophila* transposon in the *Caenorhabditis elegans* germ line. *Nature* 2001, 413(6851):70-74.

Bhutani N, Brady JJ, Damian M, Sacco A, Corbel SY, Blau HM: Reprogramming towards pluripotency requires AID-dependent DNA demethylation. *Nature* 2010, 463(7284):1042-1047.

Bingham PM, Kidwell MG, Rubin GM: The molecular basis of P-M hybrid dysgenesis: the role of the P element, a P-strain-specific transposon family. *Cell* 1982, 29(3):995-1004.

Blaese RM, Culver KW, Miller AD, Carter CS, Fleisher T, Clerici M, Shearer G, Chang L, Chiang YW, Tolstoshev P et al: T-Lymphocyte-Directed Gene-Therapy for *Ada(-)* Scid - Initial Trial Results after 4 Years. *Science* 1995, 270(5235):475-480.

Blau HM, Chiu CP, Webster C: Cytoplasmic activation of human nuclear genes in stable heterocaryons. *Cell* 1983, 32(4):1171-1180.

Bouchard C, Dittrich O, Kiermaier A, Dohmann K, Menkel A, Eilers M, Luscher B: Regulation of cyclin D2 gene expression by the *Myc/Max/Mad* network: *Myc*-dependent TRRAP recruitment and histone acetylation at the cyclin D2 promoter. *Gene Dev* 2001, 15(16):2042-2047.

Boyer LA, Lee TI, Cole MF, Johnstone SE, Levine SS, Zucker JP, Guenther MG, Kumar RM, Murray HL, Jenner RG et al: Core transcriptional regulatory circuitry in human embryonic stem cells. *Cell* 2005, 122(6):947-956.

Bradley A, Evans M, Kaufman MH, Robertson E: Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature* 1984, 309(5965):255-256.

Brand N, Petkovich M, Krust A, Chambon P, de The H, Marchio A, Tiollais P, Dejean A: Identification of a second human retinoic acid receptor. *Nature* 1988, 332(6167):850-853.

Brehm A, Ohbo K, Scholer H: The carboxy-terminal transactivation domain of Oct-4 acquires cell specificity through the POU domain. *Mol Cell Biol* 1997, 17(1):154-162.

Bretscher A, Lynch W: Identification and Localization of Immunoreactive Forms of Caldesmon in Smooth and Nonmuscle Cells - a Comparison with the Distributions of Tropomyosin and Alpha-Actinin. *J Cell Biol* 1985, 100(5):1656-1663.

Brett BT, Berquam-Vrieze KE, Nannapaneni K, Huang J, Scheetz TE, Dupuy AJ: Novel molecular and computational methods improve the accuracy of insertion site analysis in Sleeping Beauty-induced tumors. *PloS one* 2011, 6(9):e24668.

Briggs R, King TJ: Transplantation of Living Nuclei From Blastula Cells into Enucleated Frogs' Eggs. *P Natl Acad Sci USA* 1952, 38(5):455-463.

Brimble SN, Zeng X, Weiler DA, Luo Y, Liu Y, Lyons IG, Freed WJ, Robins AJ, Rao MS, Schulz TC: Karyotypic stability, genotyping, differentiation, feeder-free maintenance, and gene expression sampling in three human embryonic stem cell lines derived prior to August 9, 2001. *Stem cells and development* 2004, 13(6):585-597.

Brons IG, Smithers LE, Trotter MW, Rugg-Gunn P, Sun B, Chuva de Sousa Lopes SM, Howlett SK, Clarkson A, Ahrlund-Richter L, Pedersen RA et al: Derivation of pluripotent epiblast stem cells from mammalian embryos. *Nature* 2007, 448(7150):191-195.

Bruck N, Vitoux D, Ferry C, Duong V, Bauer A, de The H, Rochette-Egly C: A coordinated phosphorylation cascade initiated by p38MAPK/MSK1 directs RARalpha to target promoters. *The EMBO journal* 2009, 28(1):34-47.

Brumbaugh J, Hou ZG, Russell JD, Howden SE, Yu PZ, Ledvina AR, Coon JJ, Thomson JA: Phosphorylation regulates human OCT4. *P Natl Acad Sci USA* 2012, 109(19):7162-7168.

Buecker C, Chen HH, Polo JM, Daheron L, Bu L, Barakat TS, Okwieka P, Porter A, Gribnau J, Hochedlinger K et al: A murine ESC-like state facilitates transgenesis and homologous recombination in human pluripotent stem cells. *Cell stem cell* 2010, 6(6):535-546.

Buehr M, Meek S, Blair K, Yang J, Ure J, Silva J, McLay R, Hall J, Ying QL, Smith A: Capture of authentic embryonic stem cells from rat blastocysts. *Cell* 2008, 135(7):1287-1298.

Bultmann S, Morbitzer R, Schmidt CS, Thanisch K, Spada F, Elsaesser J, Lahaye T, Leonhardt H: Targeted transcriptional activation of silent oct4 pluripotency gene by combining designer TALEs and inhibition of epigenetic modifiers. *Nucleic Acids Res* 2012.

Burdon T, Stracey C, Chambers I, Nichols J, Smith A: Suppression of SHP-2 and ERK signalling promotes self-renewal of mouse embryonic stem cells. *Developmental biology* 1999, 210(1):30-43.

Buzzard JJ, Gough NM, Crook JM, Colman A: Karyotype of human ES cells during extended culture. *Nat Biotechnol* 2004, 22(4):381-382; author reply 382.

Byrne JA, Simonsson S, Western PS, Gurdon JB: Nuclei of adult mammalian somatic cells are directly reprogrammed to oct-4 stem cell gene expression by amphibian oocytes. *Curr Biol* 2003, 13(14):1206-1213.

Cadinanos J, Bradley A: Generation of an inducible and optimized piggyBac transposon system. *Nucleic Acids Res* 2007, 35(12):e87.

Carette JE, Guimaraes CP, Varadarajan M, Park AS, Wuethrich I, Godarova A, Kotecki M, Cochran BH, Spooner E, Ploegh HL et al: Haploid Genetic Screens in Human Cells Identify Host Factors Used by Pathogens. *Science* 2009, 326(5957):1231-1235.

Carey BW, Markoulaki S, Hanna J, Saha K, Gao Q, Mitalipova M, Jaenisch R: Reprogramming of murine and human somatic cells using a single polycistronic vector. *P Natl Acad Sci USA* 2009, 106(1):157-162.

Carey BW, Markoulaki S, Hanna JH, Faddah DA, Buganim Y, Kim J, Ganz K, Steine EJ, Cassady JP, Creighton MP et al: Reprogramming factor stoichiometry influences the epigenetic state and biological properties of induced pluripotent stem cells. *Cell stem cell* 2011, 9(6):588-598.

Carlberg C, Hooft van Huijsduijnen R, Staple JK, DeLamarter JF, Becker-Andre M: RZR_s, a new family of retinoid-related orphan receptors that function as both monomers and homodimers. *Mol Endocrinol* 1994, 8(6):757-770.

Cartwright P, McLean C, Sheppard A, Rivett D, Jones K, Dalton S: LIF/STAT3 controls ES cell self-renewal and pluripotency by a Myc-dependent mechanism. *Development* 2005, 132(5):885-896.

Casci T: Developmental biology - Retinoic acid passes the morphogen test. *Nat Rev Genet* 2008, 9(1):7-7.

Cauffman G, Liebaers I, Van Steirteghem A, de Velde HV: POU5F1 isoforms show different expression patterns in human embryonic stem cells and preimplantation embryos. *Stem Cells* 2006, 24(12):2685-2691.

Chambers I, Colby D, Robertson M, Nichols J, Lee S, Tweedie S, Smith A: Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell* 2003, 113(5):643-655.

Chambers I, Silva J, Colby D, Nichols J, Nijmeijer B, Robertson M, Vrana J, Jones K, Grotewold L, Smith A: Nanog safeguards pluripotency and mediates germline development. *Nature* 2007, 450(7173):1230-1234.

Chambon P: A decade of molecular biology of retinoic acid receptors. *Faseb J* 1996, 10(9):940-954.

Chen J, Liu J, Yang J, Chen Y, Chen J, Ni S, Song H, Zeng L, Ding K, Pei D: BMPs functionally replace Klf4 and support efficient reprogramming of mouse fibroblasts by Oct4 alone. *Cell research* 2011, 21(1):205-212.

Chen X, Xu H, Yuan P, Fang F, Huss M, Vega VB, Wong E, Orlov YL, Zhang W, Jiang J et al: Integration of external signaling pathways with the core transcriptional network in embryonic stem cells. *Cell* 2008, 133(6):1106-1117.

Chen Y, Yee D, Dains K, Chatterjee A, Cavalcoli J, Schneider E, Om J, Woychik RP, Magnuson T: Genotype-based screen for ENU-induced mutations in mouse embryonic stem cells. *Nat Genet* 2000, 24(3):314-317.

Chen YT, Furushima K, Hou PS, Ku AT, Deng JM, Jang CW, Fang H, Adams HP, Kuo ML, Ho HN et al: PiggyBac transposon-mediated, reversible gene transfer in human embryonic stem cells. *Stem cells and development* 2010, 19(6):763-771.

Cheong HS, Lee HC, Park BL, Kim H, Jang MJ, Han YM, Kim SY, Kim YS, Shin HD: Epigenetic modification of retinoic acid-treated human embryonic stem cells. *Bmb Rep* 2010, 43(12):830-835.

Chew JL, Loh YH, Zhang W, Chen X, Tam WL, Yeap LS, Li P, Ang YS, Lim B, Robson P et al: Reciprocal transcriptional regulation of Pou5f1 and Sox2 via the Oct4/Sox2 complex in embryonic stem cells. *Mol Cell Biol* 2005, 25(14):6031-6046.

Chia NY, Chan YS, Feng B, Lu X, Orlov YL, Moreau D, Kumar P, Yang L, Jiang J, Lau MS et al: A genome-wide RNAi screen reveals determinants of human embryonic stem cell identity. *Nature* 2010, 468(7321):316-320.

Chicas A, Kapoor A, Wang X, Aksoy O, Evertts AG, Zhang MQ, Garcia BA, Bernstein E, Lowe SW: H3K4 demethylation by Jarid1a and Jarid1b contributes to retinoblastoma-mediated gene silencing during cellular senescence. *P Natl Acad Sci USA* 2012, 109(23):8971-8976.

Chickarmane V, Enver T, Peterson C: Computational modeling of the hematopoietic erythroid-myeloid switch reveals insights into cooperativity, priming, and irreversibility. *PLoS computational biology* 2009, 5(1):e1000268.

Chiu CP, Blau HM: 5-Azacytidine permits gene activation in a previously noninducible cell type. *Cell* 1985, 40(2):417-424.

Claassen GF, Hann SR: A role for transcriptional repression of p21CIP1 by c-Myc in overcoming transforming growth factor beta -induced cell-cycle arrest. *P Natl Acad Sci USA* 2000, 97(17):9498-9503.

Conlon RA, Rossant J: Exogenous retinoic acid rapidly induces anterior ectopic expression of murine Hox-2 genes in vivo. *Development* 1992, 116(2):357-368.

Consortium Hi: Induced Pluripotent Stem Cells from Patients with Huntington's Disease Show CAG-Repeat-Expansion-Associated Phenotypes. *Cell stem cell* 2012, 11(2):264-278.

Cooley L, Kelley R, Spradling A: Insertional mutagenesis of the *Drosophila* genome with single P elements. *Science* 1988, 239(4844):1121-1128.

Cowan CA, Atienza J, Melton DA, Eggan K: Nuclear reprogramming of somatic cells after fusion with human embryonic stem cells. *Science* 2005, 309(5739):1369-1373.

Daheron L, Opitz SL, Zaehres H, Lensch MW, Andrews PW, Itskovitz-Eldor J, Daley GQ: LIF/STAT3 signaling fails to maintain self-renewal of human embryonic stem cells. *Stem Cells* 2004, 22(5):770-778.

De Los Angeles A, Loh YH, Tesar PJ, Daley GQ: Accessing naive human pluripotency. *Current opinion in genetics & development* 2012.

Deb-Rinker P, Ly D, Jezierski A, Sikorska M, Walker PR: Sequential DNA methylation of the Nanog and Oct-4 upstream regions in human NT2 cells during neuronal differentiation. *J Biol Chem* 2005, 280(8):6257-6260.

Dejean A, Bougueleret L, Grzeschik KH, Tiollais P: Hepatitis B virus DNA integration in a sequence homologous to v-erb-A and steroid receptor genes in a hepatocellular carcinoma. *Nature* 1986, 322(6074):70-72.

Dejosez M, Krumenacker JS, Zitur LJ, Passeri M, Chu LF, Songyang Z, Thomson JA, Zwaka TP: Ronin is essential for embryogenesis and the pluripotency of mouse embryonic stem cells. *Cell* 2008, 133(7):1162-1174.

Delacroix L, Moutier E, Altobelli G, Legras S, Poch O, Choukrallah MA, Bertin I, Jost B, Davidson I: Cell-specific interaction of retinoic acid receptors with target genes in mouse embryonic fibroblasts and embryonic stem cells. *Mol Cell Biol* 2010, 30(1):231-244.

Di Matteo M, Belay E, Chuah MK, Vandendriessche T: Recent developments in transposon-mediated gene therapy. *Expert opinion on biological therapy* 2012, 12(7):841-858.

DiBerardino MA, Hoffner NJ: Gene reactivation in erythrocytes: nuclear transplantation in oocytes and eggs of *Rana*. *Science* 1983, 219(4586):862-864.

Dickman ED, Thaller C, Smith SM: Temporally-regulated retinoic acid depletion produces specific neural crest, ocular and nervous system defects. *Development* 1997, 124(16):3111-3121.

Dimos JT, Rodolfa KT, Niakan KK, Weisenthal LM, Mitsumoto H, Chung W, Croft GF, Saphier G, Leibel R, Golland R et al: Induced pluripotent stem cells generated from patients with ALS can be differentiated into motor neurons. *Science* 2008, 321(5893):1218-1221.

Ding S, Wu X, Li G, Han M, Zhuang Y, Xu T: Efficient transposition of the piggyBac (PB) transposon in mammalian cells and mice. *Cell* 2005, 122(3):473-483.

Do JT, Scholer HR: Cell fusion-induced reprogramming. *Methods in molecular biology* (Clifton, NJ 2010, 636:179-190.

Doetschman T, Williams P, Maeda N: Establishment of hamster blastocyst-derived embryonic stem (ES) cells. *Developmental biology* 1988, 127(1):224-227.

Dolle P: Developmental expression of retinoic acid receptors (RARs). *Nuclear receptor signaling* 2009, 7:e006.

Doulazmi M, Frederic F, Capone F, Becker-Andre M, Delhay-Bouchaud N, Mariani J: A comparative study of Purkinje cells in two RORalpha gene mutant mice: staggerer and RORalpha(-/-). *Brain research Developmental brain research* 2001, 127(2):165-174.

Doulazmi M, Frederic F, Lemaigre-Dubreuil Y, Hadj-Sahraoui N, Delhay-Bouchaud N, Mariani J: Cerebellar Purkinje cell loss during life span of the heterozygous staggerer mouse (Rora(+)/Rora(sg)) is gender-related. *The Journal of comparative neurology* 1999, 411(2):267-273.

Driesch H: *Analytische theorie der organischen entwicklung*: Leipzig, W. Engelmann; 1894.

Dupuy AJ, Akagi K, Largaespada DA, Copeland NG, Jenkins NA: Mammalian mutagenesis using a highly mobile somatic Sleeping Beauty transposon system. *Nature* 2005, 436(7048):221-226.

Dupuy AJ, Rogers LM, Kim J, Nannapaneni K, Starr TK, Liu P, Largaespada DA, Scheetz TE, Jenkins NA, Copeland NG: A modified sleeping beauty transposon system that can be used to model a wide variety of human cancers in mice. *Cancer research* 2009, 69(20):8150-8156.

Dussault I, Fawcett D, Matthyssen A, Bader JA, Giguere V: Orphan nuclear receptor ROR alpha-deficient mice display the cerebellar defects of staggerer. *Mechanisms of development* 1998, 70(1-2):147-153.

Ebert AD, Yu J, Rose FF, Jr., Mattis VB, Lorson CL, Thomson JA, Svendsen CN: Induced pluripotent stem cells from a spinal muscular atrophy patient. *Nature* 2009, 457(7227):277-280.

- Efe JA, Hilcove S, Kim J, Zhou H, Ouyang K, Wang G, Chen J, Ding S: Conversion of mouse fibroblasts into cardiomyocytes using a direct reprogramming strategy. *Nat Cell Biol* 2011, 13(3):215-U261.
- Eilken HM, Nishikawa S, Schroeder T: Continuous single-cell imaging of blood generation from haemogenic endothelium. *Nature* 2009, 457(7231):896-900.
- Ellinger-Ziegelbauer H, Dreyer C: A retinoic acid receptor expressed in the early development of *Xenopus laevis*. *Gene Dev* 1991, 5(1):94-104.
- Esteban MA, Wang T, Qin B, Yang J, Qin D, Cai J, Li W, Weng Z, Chen J, Ni S et al: Vitamin C enhances the generation of mouse and human induced pluripotent stem cells. *Cell stem cell* 2010, 6(1):71-79.
- Esteban MA, Xu J, Yang J, Peng M, Qin D, Li W, Jiang Z, Chen J, Deng K, Zhong M et al: Generation of induced pluripotent stem cell lines from Tibetan miniature pig. *The Journal of biological chemistry* 2009, 284(26):17634-17640.
- Evans MJ: The isolation and properties of a clonal tissue culture strain of pluripotent mouse teratoma cells. *Journal of embryology and experimental morphology* 1972, 28(1):163-176.
- Evans MJ, Kaufman MH: Establishment in culture of pluripotential cells from mouse embryos. *Nature* 1981, 292(5819):154-156.
- Ezashi T, Das P, Roberts RM: Low O₂ tensions and the prevention of differentiation of hES cells. *P Natl Acad Sci USA* 2005, 102(13):4783-4788.
- Felgner PL, Gadek TR, Holm M, Roman R, Chan HW, Wenz M, Northrop JP, Ringold GM, Danielsen M: Lipofection - a Highly Efficient, Lipid-Mediated DNA-Transfection Procedure. *P Natl Acad Sci USA* 1987, 84(21):7413-7417.
- Feng B, Jiang J, Kraus P, Ng JH, Heng JC, Chan YS, Yaw LP, Zhang W, Loh YH, Han J et al: Reprogramming of fibroblasts into induced pluripotent stem cells with orphan nuclear receptor Esrrb. *Nat Cell Biol* 2009, 11(2):197-203.

Fibrici G: De Formatione Ovi, et Pulli Tractatus Accuratissimus: Patavii [Padua]: ex Officina Aloysii Bencii Bibliopolae; 1621.

Flasza M, Shering AF, Smith K, Andrews PW, Talley P, Johnson PA: Reprogramming in inter-species embryonal carcinoma-somatic cell hybrids induces expression of pluripotency and differentiation markers. *Cloning and stem cells* 2003, 5(4):339-354.

Flynn PJ, Miller WJ, Weisdorf DJ, Arthur DC, Brunning R, Branda RF: Retinoic acid treatment of acute promyelocytic leukemia: in vitro and in vivo observations. *Blood* 1983, 62(6):1211-1217.

Forsyth NR, Musio A, Vezzoni P, Simpson AH, Noble BS, McWhir J: Physiologic oxygen enhances human embryonic stem cell clonal recovery and reduces chromosomal abnormalities. *Cloning and stem cells* 2006, 8(1):16-23.

Foshay KM, Looney TJ, Chari S, Mao FF, Lee JH, Zhang L, Fernandes CJ, Baker SW, Clift KL, Gaetz J et al: Embryonic Stem Cells Induce Pluripotency in Somatic Cell Fusion through Biphasic Reprogramming. *Mol Cell* 2012, 46(2):159-170.

Frank SR, Schroeder M, Fernandez P, Taubert S, Amati B: Binding of c-Myc to chromatin mediates mitogen-induced acetylation of histone H4 and gene activation. *Gene Dev* 2001, 15(16):2069-2082.

Fraser MJ, Ciszczon T, Elick T, Bauser C: Precise excision of TTAA-specific lepidopteran transposons piggyBac (IFP2) and tagalong (TFP3) from the baculovirus genome in cell lines from two species of Lepidoptera. *Insect molecular biology* 1996, 5(2):141-151.

Freberg CT, Dahl JA, Timoskainen S, Collas P: Epigenetic reprogramming of OCT4 and NANOG regulatory regions by embryonal carcinoma cell extract. *Molecular biology of the cell* 2007, 18(5):1543-1553.

French AJ, Adams CA, Anderson LS, Kitchen JR, Hughes MR, Wood SH: Development of human cloned blastocysts following somatic cell nuclear transfer with adult fibroblasts. *Stem Cells* 2008, 26(2):485-493.

Fuhrmann G, Chung AC, Jackson KJ, Hummelke G, Baniahmad A, Sutter J, Sylvester I, Scholer HR, Cooney AJ: Mouse germline restriction of Oct4 expression by germ cell nuclear factor. *Developmental cell* 2001, 1(3):377-387.

Fuhrmann G, Sylvester I, Scholer HR: Repression of Oct-4 during embryonic cell differentiation correlates with the appearance of TRIF, a transiently induced DNA-binding factor. *Cell Mol Biol (Noisy-le-grand)* 1999, 45(5):717-724.

Furst DO, Cross RA, Demey J, Small JV: Caldesmon Is an Elongated, Flexible Molecule Localized in the Actomyosin Domains of Smooth-Muscle. *Embo J* 1986, 5(2):251-257.

Fusaki N, Ban H, Nishiyama A, Saeki K, Hasegawa M: Efficient induction of transgene-free human pluripotent stem cells using a vector based on Sendai virus, an RNA virus that does not integrate into the host genome. *Proceedings of the Japan Academy* 2009, 85(8):348-362.

Gao S, Chung YG, Parseghian MH, King GJ, Adashi EY, Latham KE: Rapid H1 linker histone transitions following fertilization or somatic cell nuclear transfer: evidence for a uniform developmental program in mice. *Developmental biology* 2004, 266(1):62-75.

Gerami-Naini B, Dovzhenko OV, Durning M, Wegner FH, Thomson JA, Golos TG: Trophoblast differentiation in embryoid bodies derived from human embryonic stem cells. *Endocrinology* 2004, 145(4):1517-1524.

Geurts AM, Collier LS, Geurts JL, Oseth LL, Bell ML, Mu D, Lucito R, Godbout SA, Green LE, Lowe SW et al: Gene mutations and genomic rearrangements in the mouse as a result of transposon mobilization from chromosomal concatemers. *PLoS genetics* 2006, 2(9):1413-1423.

Ghyselinck NB, Dupe V, Dierich A, Messaddeq N, Garnier JM, Rochette-Egly C, Chambon P, Mark M: Role of the retinoic acid receptor beta (RARbeta) during mouse development. *The International journal of developmental biology* 1997, 41(3):425-447.

Gianni M, Bauer A, Garattini E, Chambon P, Rochette-Egly C: Phosphorylation by p38MAPK and recruitment of SUG-1 are required for RA-induced RAR gamma degradation and transactivation. *The EMBO journal* 2002, 21(14):3760-3769.

Gianni M, Parrella E, Raska I, Jr., Gaillard E, Nigro EA, Gaudon C, Garattini E, Rochette-Egly C: P38MAPK-dependent phosphorylation and degradation of SRC-3/AIB1 and RARalpha-mediated transcription. *The EMBO journal* 2006, 25(4):739-751.

Giguere V, Beatty B, Squire J, Copeland NG, Jenkins NA: The orphan nuclear receptor ROR alpha (RORA) maps to a conserved region of homology on human chromosome 15q21-q22 and mouse chromosome 9. *Genomics* 1995, 28(3):596-598.

Giguere V, Ong ES, Segui P, Evans RM: Identification of a receptor for the morphogen retinoic acid. *Nature* 1987, 330(6149):624-629.

Giguere V, Tini M, Flock G, Ong E, Evans RM, Otulakowski G: Isoform-specific amino-terminal domains dictate DNA-binding properties of ROR alpha, a novel family of orphan hormone nuclear receptors. *Gene Dev* 1994, 8(5):538-553.

Gillespie RF, Gudas LJ: Retinoic acid receptor isotype specificity in F9 teratocarcinoma stem cells results from the differential recruitment of coregulators to retinoic response elements. *The Journal of biological chemistry* 2007, 282(46):33421-33434.

Gillespie RF, Gudas LJ: Retinoid regulated association of transcriptional co-regulators and the polycomb group protein SUZ12 with the retinoic acid response elements of Hoxa1, RARbeta(2), and Cyp26A1 in F9 embryonal carcinoma cells. *Journal of molecular biology* 2007, 372(2):298-316.

Giorgetti A, Montserrat N, Aasen T, Gonzalez F, Rodriguez-Piza I, Vassena R, Raya A, Boue S, Barrero MJ, Corbella BA et al: Generation of induced pluripotent stem cells from human cord blood using OCT4 and SOX2. *Cell stem cell* 2009, 5(4):353-357.

Gloor GB, Nassif NA, Johnson-Schlitz DM, Preston CR, Engels WR: Targeted gene replacement in *Drosophila* via P element-induced gap repair. *Science* 1991, 253(5024):1110-1117.

Godin I, Wylie C, Heasman J: Genital Ridges Exert Long-Range Effects on Mouse Primordial Germ-Cell Numbers and Direction of Migration in Culture. *Development* 1990, 108(2):357-363.

Goff SP, Witte ON, Gilboa E, Rosenberg N, Baltimore D: Genome structure of Abelson murine leukemia virus variants: proviruses in fibroblasts and lymphoid cells. *Journal of virology* 1981, 38(2):460-468.

Golling G, Amsterdam A, Sun Z, Antonelli M, Maldonado E, Chen W, Burgess S, Haldi M, Artzt K, Farrington S et al: Insertional mutagenesis in zebrafish rapidly identifies genes essential for early vertebrate development. *Nat Genet* 2002, 31(2):135-140.

Gonzalez F, Boue S, Izpisua Belmonte JC: Methods for making induced pluripotent stem cells: reprogramming a la carte. *Nat Rev Genet* 2011, 12(4):231-242.

Gore A, Li Z, Fung HL, Young JE, Agarwal S, Antosiewicz-Bourget J, Canto I, Giorgetti A, Israel MA, Kiskinis E et al: Somatic coding mutations in human induced pluripotent stem cells. *Nature* 2011, 471(7336):63-67.

Gossen M, Bujard H: Tight Control of Gene-Expression in Mammalian-Cells by Tetracycline-Responsive Promoters. *P Natl Acad Sci USA* 1992, 89(12):5547-5551.

Gossen M, Freundlieb S, Bender G, Muller G, Hillen W, Bujard H: Transcriptional Activation by Tetracyclines in Mammalian-Cells. *Science* 1995, 268(5218):1766-1769.

Graves KH, Moreadith RW: Derivation and characterization of putative pluripotential embryonic stem cells from preimplantation rabbit embryos. *Molecular reproduction and development* 1993, 36(4):424-433.

Gresch O, Engel FB, Nesic D, Tran TT, England HM, Hickman ES, Korner I, Gan L, Chen S, Castro-Obregon S et al: New non-viral method for gene transfer into primary cells. *Methods* 2004, 33(2):151-163.

Gu P, Goodwin B, Chung AC, Xu X, Wheeler DA, Price RR, Galardi C, Peng L, Latour AM, Koller BH et al: Orphan nuclear receptor LRH-1 is required to maintain Oct4 expression at the epiblast stage of embryonic development. *Mol Cell Biol* 2005, 25(9):3492-3505.

Guo G, Huang Y, Humphreys P, Wang X, Smith A: A PiggyBac-based recessive screening method to identify pluripotency regulators. *PloS one* 2011, 6(4):e18189.

Guo G, Smith A: A genome-wide screen in EpiSCs identifies Nr5a nuclear receptors as potent inducers of ground state pluripotency. *Development* 2010, 137(19):3185-3192.

Guo G, Yang J, Nichols J, Hall JS, Eyres I, Mansfield W, Smith A: Klf4 reverts developmentally programmed restriction of ground state pluripotency. *Development* 2009, 136(7):1063-1069.

Guo HQ, Wang CLA: Specific disruption of smooth muscle caldesmon expression in mice. *Biochem Bioph Res Co* 2005, 330(4):1132-1137.

Gupta P, Ho PC, Huq MM, Ha SG, Park SW, Khan AA, Tsai NP, Wei LN: Retinoic acid-stimulated sequential phosphorylation, PML recruitment, and SUMOylation of nuclear receptor TR2 to suppress Oct4 expression. *P Natl Acad Sci USA* 2008, 105(32):11424-11429.

Gurdon JB: The developmental capacity of nuclei taken from intestinal epithelium cells of feeding tadpoles. *Journal of embryology and experimental morphology* 1962, 10:622-640.

Gurdon JB, Brennan S, Fairman S, Mohun TJ: Transcription of muscle-specific actin genes in early *Xenopus* development: nuclear transplantation and cell dissociation. *Cell* 1984, 38(3):691-700.

Gurdon JB, Elsdale TR, Fischberg M: Sexually mature individuals of *Xenopus laevis* from the transplantation of single somatic nuclei. *Nature* 1958, 182(4627):64-65.

Gurdon JB, Laskey RA, Reeves OR: The developmental capacity of nuclei transplanted from keratinized skin cells of adult frogs. *Journal of embryology and experimental morphology* 1975, 34(1):93-112.

Hale F: Pigs born without eyeballs. *Journal of Heredity* 1933, 24:105-106.

Hall J, Guo G, Wray J, Eyres I, Nichols J, Grotewold L, Morfopoulou S, Humphreys P, Mansfield W, Walker R et al: Oct4 and LIF/Stat3 additively induce Kruppel factors to sustain embryonic stem cell self-renewal. *Cell stem cell* 2009, 5(6):597-609.

Hall VJ, Compton D, Stojkovic P, Nesbitt M, Herbert M, Murdoch A, Stojkovic M: Developmental competence of human in vitro aged oocytes as host cells for nuclear transfer. *Human reproduction (Oxford, England)* 2007, 22(1):52-62.

Hamanaka RB, Chandel NS: Targeting glucose metabolism for cancer therapy. *The Journal of experimental medicine* 2012, 209(2):211-215.

Hamilton BA, Frankel WN, Kerrebrock AW, Hawkins TL, FitzHugh W, Kusumi K, Russell LB, Mueller KL, van Berkel V, Birren BW et al: Disruption of the nuclear hormone receptor RORalpha in staggerer mice. *Nature* 1996, 379(6567):736-739.

Hammachi F MAM, Sharov A.A., Livigni A., Narayan S., Papapetrou E.P., O'Malley J., Kaji K., Ko M.S.H., Ptashne M., Brickman J.M.: Transcriptional Activation by Oct4 Is Sufficient for the Maintenance and Induction of Pluripotency. *Cell Reports* 2012, 1(2):99-109.

Han DW, Tapia N, Joo JY, Greber B, Arauzo-Bravo MJ, Bernemann C, Ko K, Wu G, Stehling M, Do JT et al: Epiblast stem cell subpopulations represent mouse embryos of distinct pregastrulation stages. *Cell* 2010, 143(4):617-627.

Han J, Yuan P, Yang H, Zhang J, Soh BS, Li P, Lim SL, Cao S, Tay J, Orlov YL et al: Tbx3 improves the germ-line competency of induced pluripotent stem cells. *Nature* 2010, 463(7284):1096-1100.

Hanna J, Cheng AW, Saha K, Kim J, Lengner CJ, Soldner F, Cassady JP, Muffat J, Carey BW, Jaenisch R: Human embryonic stem cells with biological and epigenetic characteristics similar to those of mouse ESCs. *P Natl Acad Sci USA* 2010, 107(20):9222-9227.

Hanna J, Markoulaki S, Schorderet P, Carey BW, Beard C, Wernig M, Creighton MP, Steine EJ, Cassady JP, Foreman R et al: Direct reprogramming of terminally differentiated mature B lymphocytes to pluripotency. *Cell* 2008, 133(2):250-264.

Hanna J, Saha K, Pando B, van Zon J, Lengner CJ, Creighton MP, van Oudenaarden A, Jaenisch R: Direct cell reprogramming is a stochastic process amenable to acceleration. *Nature* 2009, 462(7273):595-601.

Hanna J, Wernig M, Markoulaki S, Sun CW, Meissner A, Cassady JP, Beard C, Brambrink T, Wu LC, Townes TM et al: Treatment of sickle cell anemia mouse model with iPS cells generated from autologous skin. *Science* 2007, 318(5858):1920-1923.

Hansen GM, Markesich DC, Burnett MB, Zhu QH, Dionne KM, Richter LJ, Finnell RH, Sands AT, Zambrowicz BP, Abuin A: Large-scale gene trapping in C57BL/6N mouse embryonic stem cells. *Genome research* 2008, 18(10):1670-1679.

Hansis C, Barreto G, Maltry N, Niehrs C: Nuclear reprogramming of human somatic cells by xenopus egg extract requires BRG1. *Curr Biol* 2004, 14(16):1475-1480.

Hansis C, Grifo JA, Krey LC: Oct-4 expression in inner cell mass and trophectoderm of human blastocysts. *Mol Hum Reprod* 2000, 6(11):999-1004.

Hansis C, Tang YX, Grifo JA, Krey LC: Analysis of Oct-4 expression and ploidy in individual human blastomeres. *Mol Hum Reprod* 2001, 7(2):155-161.

Harris H: Behaviour of differentiated nuclei in heterokaryons of animal cells from different species. *Nature* 1965, 206(984):583-588.

Harris H, Watkins JF, Ford CE, Schoepl GI: Artificial heterokaryons of animal cells from different species. *J Cell Sci* 1966, 1(1):1-30.

Harvey W: *Exercitationes de generatione animalium*: Hagae Comitum, apud A. Leers, 1680.; 1651.

Hasegawa K, Nakatsuji N: Insulators prevent transcriptional interference between two promoters in a double gene construct for transgenesis. *FEBS letters* 2002, 520(1-3):47-52.

Hattori N, Nishino K, Ko YG, Hattori N, Ohgane J, Tanaka S, Shiota K: Epigenetic control of mouse Oct-4 gene expression in embryonic stem cells and trophoblast stem cells. *J Biol Chem* 2004, 279(17):17063-17069.

Hawkins RD, Hon GC, Lee LK, Ngo Q, Lister R, Pelizzola M, Edsall LE, Kuan S, Luu Y, Klugman S et al: Distinct epigenomic landscapes of pluripotent and lineage-committed human cells. *Cell stem cell* 2010, 6(5):479-491.

He YW, Deftos ML, Ojala EW, Bevan MJ: ROR γ t, a novel isoform of an orphan receptor, negatively regulates Fas ligand expression and IL-2 production in T cells. *Immunity* 1998, 9(6):797-806.

Heintzman ND, Stuart RK, Hon G, Fu Y, Ching CW, Hawkins RD, Barrera LO, Van Calcar S, Qu C, Ching KA et al: Distinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome. *Nat Genet* 2007, 39(3):311-318.

Heng JC, Feng B, Han J, Jiang J, Kraus P, Ng JH, Orlov YL, Huss M, Yang L, Lufkin T et al: The nuclear receptor Nr5a2 can replace Oct4 in the reprogramming of murine somatic cells to pluripotent cells. *Cell stem cell* 2010, 6(2):167-174.

Heyman RA, Mangelsdorf DJ, Dyck JA, Stein RB, Eichele G, Evans RM, Thaller C: 9-Cis Retinoic Acid Is a High-Affinity Ligand for the Retinoid-X Receptor. *Cell* 1992, 68(2):397-406.

Hirose T, Smith RJ, Jetten AM: ROR γ : the third member of ROR/RZR orphan receptor subfamily that is highly expressed in skeletal muscle. *Biochem Biophys Res Commun* 1994, 205(3):1976-1983.

Hochedlinger K, Plath K: Epigenetic reprogramming and induced pluripotency. *Development* 2009, 136(4):509-523.

Hochedlinger K, Yamada Y, Beard C, Jaenisch R: Ectopic expression of Oct-4 blocks progenitor-cell differentiation and causes dysplasia in epithelial tissues. *Cell* 2005, 121(3):465-477.

Holder N, Hill J: Retinoic acid modifies development of the midbrain-hindbrain border and affects cranial ganglion formation in zebrafish embryos. *Development* 1991, 113(4):1159-1170.

Hong H, Takahashi K, Ichisaka T, Aoi T, Kanagawa O, Nakagawa M, Okita K, Yamanaka S: Suppression of induced pluripotent stem cell generation by the p53-p21 pathway. *Nature* 2009, 460(7259):1132-1135.

Hu X, Lazar MA: The CoRNR motif controls the recruitment of corepressors by nuclear hormone receptors. *Nature* 1999, 402(6757):93-96.

Huang X, Guo HF, Tammana S, Jung YC, Mellgren E, Bassi P, Cao Q, Tu ZJ, Kim YC, Ekker SC et al: Gene Transfer Efficiency and Genome-Wide Integration Profiling of Sleeping Beauty, Tol2, and PiggyBac Transposons in Human Primary T Cells (vol 18, pg 1803, 2010). *Mol Ther* 2010, 18(11):2038-2038.

Huangfu D, Maehr R, Guo W, Eijkelenboom A, Snitow M, Chen AE, Melton DA: Induction of pluripotent stem cells by defined factors is greatly improved by small-molecule compounds. *Nat Biotechnol* 2008, 26(7):795-797.

Huangfu D, Osafune K, Maehr R, Guo W, Eijkelenboom A, Chen S, Muhlestein W, Melton DA: Induction of pluripotent stem cells from primary human fibroblasts with only Oct4 and Sox2. *Nat Biotechnol* 2008, 26(11):1269-1275.

Huggenvik JI, Collard MW, Kim YW, Sharma RP: Modification of the retinoic acid signaling pathway by the catalytic subunit of protein kinase-A. *Mol Endocrinol* 1993, 7(4):543-550.

Humphrey RK, Beattie GM, Lopez AD, Bucay N, King CC, Firpo MT, Rose-John S, Hayek A: Maintenance of pluripotency in human embryonic stem cells is STAT3 independent. *Stem Cells* 2004, 22(4):522-530.

Hussein SM, Batada NN, Vuoristo S, Ching RW, Autio R, Narva E, Ng S, Sourour M, Hamalainen R, Olsson C et al: Copy number variation and selection during reprogramming to pluripotency. *Nature* 2011, 471(7336):58-62.

Ichida JK, Blanchard J, Lam K, Son EY, Chung JE, Egli D, Loh KM, Carter AC, Di Giorgio FP, Koszka K et al: A small-molecule inhibitor of tgf-Beta signaling replaces sox2 in reprogramming by inducing nanog. *Cell stem cell* 2009, 5(5):491-503.

Ieda M, Fu JD, Delgado-Olguin P, Vedantham V, Hayashi Y, Bruneau BG, Srivastava D: Direct reprogramming of fibroblasts into functional cardiomyocytes by defined factors. *Cell* 2010, 142(3):375-386.

Inoue A, Zhang Y: Replication-dependent loss of 5-hydroxymethylcytosine in mouse preimplantation embryos. *Science* 2011, 334(6053):194.

Itzhaki I, Maizels L, Huber I, Zwi-Dantsis L, Caspi O, Winterstern A, Feldman O, Gepstein A, Arbel G, Hammerman H et al: Modelling the long QT syndrome with induced pluripotent stem cells. *Nature* 2011, 471(7337):225-229.

Iuchi S, Easley K, Matsuzaki K, Weiner L, O'Connor N, Green H: Alternative subcellular locations of keratinocyte basonuclin. *Experimental dermatology* 2000, 9(3):178-184.

Iuchi S, Green H: Nuclear localization of basonuclin in human keratinocytes and the role of phosphorylation. *P Natl Acad Sci USA* 1997, 94(15):7948-7953.

Ivics Z, Hackett PB, Plasterk RH, Izsvak Z: Molecular reconstruction of Sleeping Beauty, a Tc1-like transposon from fish, and its transposition in human cells. *Cell* 1997, 91(4):501-510.

Iwasaki H, Mizuno S, Arinobu Y, Ozawa H, Mori Y, Shigematsu H, Takatsu K, Tenen DG, Akashi K: The order of expression of transcription factors directs hierarchical specification of hematopoietic lineages. *Gene Dev* 2006, 20(21):3010-3021.

Izsvak Z, Ivics Z: Sleeping Beauty hits them all: transposon-mediated saturation mutagenesis in the mouse germline. *Nature methods* 2005, 2(10):735-736.

Jaenisch R: Germ line integration and Mendelian transmission of the exogenous Moloney leukemia virus. *P Natl Acad Sci USA* 1976, 73(4):1260-1264.

James D, Noggle SA, Swigut T, Brivanlou AH: Contribution of human embryonic stem cells to mouse blastocysts. *Developmental biology* 2006, 295(1):90-102.

Jang H, Kim TW, Yoon S, Choi SY, Kang TW, Kim SY, Kwon YW, Cho EJ, Youn HD: O-GlcNAc Regulates Pluripotency and Reprogramming by Directly Acting on Core Components of the Pluripotency Network. *Cell stem cell* 2012, 11(1):62-74.

Jang J, Kang HC, Kim HS, Kim JY, Huh YJ, Kim DS, Yoo JE, Lee JA, Lim B, Lee J et al: Induced pluripotent stem cell models from X-linked adrenoleukodystrophy patients. *Annals of neurology* 2011, 70(3):402-409.

Jetten AM: Retinoid-related orphan receptors (RORs): critical roles in development, immunity, circadian rhythm, and cellular metabolism. *Nuclear receptor signaling* 2009, 7:e003.

Ji Z, Tian B: Reprogramming of 3' untranslated regions of mRNAs by alternative polyadenylation in generation of pluripotent stem cells from different cell types. *PloS one* 2009, 4(12):e8419.

Jiang J, Chan YS, Loh YH, Cai J, Tong GQ, Lim CA, Robson P, Zhong S, Ng HH: A core Klf circuitry regulates self-renewal of embryonic stem cells. *Nat Cell Biol* 2008, 10(3):353-360.

Jiang W, Shi Y, Zhao D, Chen S, Yong J, Zhang J, Qing T, Sun X, Zhang P, Ding M et al: In vitro derivation of functional insulin-producing cells from human embryonic stem cells. *Cell research* 2007, 17(4):333-344.

Jin JP, Arias EE, Chen J, Harper JW, Walter JC: A family of diverse Cul4-Ddb1-interacting proteins includes Cdt2, which is required for S phase destruction of the replication factor Cdt1. *Mol Cell* 2006, 23(5):709-721.

Judson RL, Babiarz JE, Venere M, Blueloch R: Embryonic stem cell-specific microRNAs promote induced pluripotency. *Nat Biotechnol* 2009, 27(5):459-461.

Jullien J, Astrand C, Halley-Stott RP, Garrett N, Gurdon JB: Characterization of somatic cell nuclear reprogramming by oocytes in which a linker histone is required for pluripotency gene reactivation. *P Natl Acad Sci USA* 2010, 107(12):5483-5488.

Kagey MH, Newman JJ, Bilodeau S, Zhan Y, Orlando DA, van Berkum NL, Ebmeier CC, Goossens J, Rahl PB, Levine SS et al: Mediator and cohesin connect gene expression and chromatin architecture. *Nature* 2010, 467(7314):430-435.

Kahan BW, Ephrussi B: Developmental potentialities of clonal in vitro cultures of mouse testicular teratoma. *Journal of the National Cancer Institute* 1970, 44(5):1015-1036.

Kaji K, Norrby K, Paca A, Mileikovsky M, Mohseni P, Woltjen K: Virus-free induction of pluripotency and subsequent excision of reprogramming factors. *Nature* 2009, 458(7239):771-U112.

Kajimura S, Seale P, Kubota K, Lunsford E, Frangioni JV, Gygi SP, Spiegelman BM: Initiation of myoblast to brown fat switch by a PRDM16-C/EBP-beta transcriptional complex. *Nature* 2009, 460(7259):1154-1158.

Kallen J, Schlaeppli JM, Bitsch F, Delhon I, Fournier B: Crystal structure of the human RORalpha Ligand binding domain in complex with cholesterol sulfate at 2.2 Å. *The Journal of biological chemistry* 2004, 279(14):14033-14038.

Kang Y, Zhang XY, Jiang W, Wu CQ, Chen CM, Zheng YF, Gu JR, Xu CJ: Tumor-directed gene therapy in mice using a composite nonviral gene delivery system consisting of the piggyBac transposon and polyethylenimine. *Bmc Cancer* 2009, 9.

Kastner P, Grondona JM, Mark M, Gansmuller A, LeMeur M, Decimo D, Vonesch JL, Dolle P, Chambon P: Genetic analysis of RXR alpha developmental function: convergence of RXR and RAR signaling pathways in heart and eye morphogenesis. *Cell* 1994, 78(6):987-1003.

Kawakami K, Shima A, Kawakami N: Identification of a functional transposase of the Tol2 element, an Ac-like element from the Japanese medaka fish, and its transposition in the zebrafish germ lineage. *P Natl Acad Sci USA* 2000, 97(21):11403-11408.

Kawamura T, Suzuki J, Wang YV, Menendez S, Morera LB, Raya A, Wahl GM, Izpisua Belmonte JC: Linking the p53 tumour suppressor pathway to somatic cell reprogramming. *Nature* 2009, 460(7259):1140-1144.

Kelley MR, Kidd S, Berg RL, Young MW: Restriction of P-Element Insertions at the Notch Locus of *Drosophila-Melanogaster*. *Mol Cell Biol* 1987, 7(4):1545-1548.

Kemler I, Schreiber E, Muller MM, Matthias P, Schaffner W: Octamer transcription factors bind to two different sequence motifs of the immunoglobulin heavy chain promoter. *The EMBO journal* 1989, 8(7):2001-2008.

Keng VW, Villanueva A, Chiang DY, Dupuy AJ, Ryan BJ, Matisse I, Silverstein KA, Sarver A, Starr TK, Akagi K et al: A conditional transposon-based insertional mutagenesis screen for genes associated with mouse hepatocellular carcinoma. *Nat Biotechnol* 2009, 27(3):264-274.

Kile BT, Hentges KE, Clark AT, Nakamura H, Salinger AP, Liu B, Box N, Stockton DW, Johnson RL, Behringer RR et al: Functional genetic analysis of mouse chromosome 11. *Nature* 2003, 425(6953):81-86.

Kim D, Kim CH, Moon JI, Chung YG, Chang MY, Han BS, Ko S, Yang E, Cha KY, Lanza R et al: Generation of human induced pluripotent stem cells by direct delivery of reprogramming proteins. *Cell stem cell* 2009, 4(6):472-476.

Kim J, Chu J, Shen X, Wang J, Orkin SH: An extended transcriptional network for pluripotency of embryonic stem cells. *Cell* 2008, 132(6):1049-1061.

Kim JB, Greber B, Arauzo-Bravo MJ, Meyer J, Park KI, Zaehres H, Scholer HR: Direct reprogramming of human neural stem cells by OCT4. *Nature* 2009, 461(7264):649-643.

Kim JB, Sebastiano V, Wu G, Arauzo-Bravo MJ, Sasse P, Gentile L, Ko K, Ruau D, Ehrlich M, van den Boom D et al: Oct4-induced pluripotency in adult neural stem cells. *Cell* 2009, 136(3):411-419.

Kim K, Doi A, Wen B, Ng K, Zhao R, Cahan P, Kim J, Aryee MJ, Ji H, Ehrlich LI et al: Epigenetic memory in induced pluripotent stem cells. *Nature* 2010, 467(7313):285-290.

Kim TH, Barrera LO, Zheng M, Qu C, Singer MA, Richmond TA, Wu Y, Green RD, Ren B: A high-resolution map of active promoters in the human genome. *Nature* 2005, 436(7052):876-880.

King TJ, Briggs R: Changes in the Nuclei of Differentiating Gastrula Cells, as Demonstrated by Nuclear Transplantation. *P Natl Acad Sci USA* 1955, 41(5):321-325.

Klein G: Specific chromosomal translocations and the genesis of B-cell-derived tumors in mice and men. *Cell* 1983, 32(2):311-315.

Kleinsmith LJ, Pierce GB, Jr.: Multipotentiality of Single Embryonal Carcinoma Cells. *Cancer research* 1964, 24:1544-1551.

Klemm JD, Rould MA, Aurora R, Herr W, Pabo CO: Crystal structure of the Oct-1 POU domain bound to an octamer site: DNA recognition with tethered DNA-binding modules. *Cell* 1994, 77(1):21-32.

Klempnauer KH, Gonda TJ, Bishop JM: Nucleotide sequence of the retroviral leukemia gene v-myb and its cellular progenitor c-myb: the architecture of a transduced oncogene. *Cell* 1982, 31(2 Pt 1):453-463.

Kliwer SA, Umesono K, Mangelsdorf DJ, Evans RM: Retinoid X receptor interacts with nuclear receptors in retinoic acid, thyroid hormone and vitamin D3 signalling. *Nature* 1992, 355(6359):446-449.

Kobayashi T, Yamaguchi T, Hamanaka S, Kato-Itoh M, Yamazaki Y, Ibata M, Sato H, Lee YS, Usui J, Knisely AS et al: Generation of rat pancreas in mouse by interspecific blastocyst injection of pluripotent stem cells. *Cell* 2010, 142(5):787-799.

Kopp JL, Ormsbee BD, Desler M, Rizzino A: Small increases in the level of Sox2 trigger the differentiation of mouse embryonic stem cells. *Stem Cells* 2008, 26(4):903-911.

Krust A, Kastner P, Petkovich M, Zelent A, Chambon P: A third human retinoic acid receptor, hRAR-gamma. *P Natl Acad Sci USA* 1989, 86(14):5310-5314.

Kuehn MR, Bradley A, Robertson EJ, Evans MJ: A potential animal model for Lesch-Nyhan syndrome through introduction of HPRT mutations into mice. *Nature* 1987, 326(6110):295-298.

Kuijk EW, van Tol LT, Van de Velde H, Wubbolts R, Welling M, Geijsen N, Roelen BA: The roles of FGF and MAP kinase signaling in the segregation of the epiblast and hypoblast cell lineages in bovine and human embryos. *Development* 2012, 139(5):871-882.

Kulesa H, Frampton J, Graf T: GATA-1 reprograms avian myelomonocytic cell lines into eosinophils, thromboblats, and erythroblats. *Gene Dev* 1995, 9(10):1250-1262.

Kunath T, Saba-El-Leil MK, Almousailleakh M, Wray J, Meloche S, Smith A: FGF stimulation of the Erk1/2 signalling cascade triggers transition of pluripotent embryonic stem cells from self-renewal to lineage commitment. *Development* 2007, 134(16):2895-2902.

Laiosa CV, Stadtfeld M, Xie H, de Andres-Aguayo L, Graf T: Reprogramming of committed T cell progenitors to macrophages and dendritic cells by C/EBP alpha and PU.1 transcription factors. *Immunity* 2006, 25(5):731-744.

Lammer EJ, Chen DT, Hoar RM, Agnish ND, Benke PJ, Braun JT, Curry CJ, Fernhoff PM, Grix AW, Jr., Lott IT et al: Retinoic acid embryopathy. *The New England journal of medicine* 1985, 313(14):837-841.

Lang MR, Patterson LB, Gordon TN, Johnson SL, Parichy DM: Basonuclin-2 Requirements for Zebrafish Adult Pigment Pattern Development and Female Fertility. *PLoS genetics* 2009, 5(11).

Laskey RA, Gurdon JB: Genetic content of adult somatic cells tested by nuclear transplantation from cultured cells. *Nature* 1970, 228(5278):1332-1334.

Laursen KB, Wong PM, Gudas LJ: Epigenetic regulation by RARalpha maintains ligand-independent transcriptional activity. *Nucleic Acids Res* 2012, 40(1):102-115.

Ledda-Columbano GM, Pibiri M, Molotzu F, Cossu C, Sanna L, Simbula G, Perra A, Columbano A: Induction of hepatocyte proliferation by retinoic acid. *Carcinogenesis* 2004, 25(11):2061-2066.

Lee G, Papapetrou EP, Kim H, Chambers SM, Tomishima MJ, Fasano CA, Ganat YM, Menon J, Shimizu F, Viale A et al: Modelling pathogenesis and treatment of familial dysautonomia using patient-specific iPSCs. *Nature* 2009, 461(7262):402-406.

Lee J, Kim HK, Rho JY, Han YM, Kim J: The human OCT-4 isoforms differ in their ability to confer self-renewal. *J Biol Chem* 2006, 281(44):33554-33565.

Lee JM, Kim IS, Kim H, Lee JS, Kim K, Yim HY, Jeong J, Kim JH, Kim JY, Lee H et al: RORalpha attenuates Wnt/beta-catenin signaling by PKCalpha-dependent phosphorylation in colon cancer. *Mol Cell* 2010, 37(2):183-195.

Leeb M, Wutz A: Derivation of haploid embryonic stem cells from mouse embryos. *Nature* 2011, 479(7371):131-134.

Leeb M, Wutz A: Derivation of haploid embryonic stem cells from mouse embryos. *Nature* 2011, 479(7371):131-134.

Lefort N, Feyeux M, Bas C, Feraud O, Bennaceur-Griscelli A, Tachdjian G, Peschanski M, Perrier AL: Human embryonic stem cells reveal recurrent genomic instability at 20q11.21. *Nat Biotechnol* 2008, 26(12):1364-1366.

Leid M, Kastner P, Lyons R, Nakshatri H, Saunders M, Zacharewski T, Chen JY, Staub A, Garnier JM, Mader S et al: Purification, cloning, and RXR identity of the

HeLa cell factor with which RAR or TR heterodimerizes to bind target sequences efficiently. *Cell* 1992, 68(2):377-395.

Lengner CJ, Gimelbrant AA, Erwin JA, Cheng AW, Guenther MG, Welstead GG, Alagappan R, Frampton GM, Xu P, Muffat J et al: Derivation of pre-X inactivation human embryonic stem cells under physiological oxygen concentrations. *Cell* 2010, 141(5):872-883.

Lenhoff SG, Lenhoff HM: *Hydra and the birth of experimental biology-1744: Abraham Trembley's memoirs concerning the natural history of a type of freshwater polyp with arms shaped like horns.* . Pacific Grove, CA: Boxwood Press; 1986.

Levin AA, Sturzenbecker LJ, Kazmer S, Bosakowski T, Huselton C, Allenby G, Speck J, Kratzeisen C, Rosenberger M, Lovey A et al: 9-cis retinoic acid stereoisomer binds and activates the nuclear receptor RXR alpha. *Nature* 1992, 355(6358):359-361.

Li H, Collado M, Villasante A, Strati K, Ortega S, Canamero M, Blasco MA, Serrano M: The Ink4/Arf locus is a barrier for iPS cell reprogramming. *Nature* 2009, 460(7259):1136-1139.

Li J, Shen H, Himmel KL, Dupuy AJ, Largaespada DA, Nakamura T, Shaughnessy JD, Jr., Jenkins NA, Copeland NG: Leukaemia disease genes: large-scale cloning and pathway predictions. *Nat Genet* 1999, 23(3):348-353.

Li J, Zhang JM, Li X, Suo F, Zhang MJ, Hou W, Han J, Du LL: A piggyBac transposon-based mutagenesis system for the fission yeast *Schizosaccharomyces pombe*. *Nucleic Acids Res* 2011, 39(6):e40.

Li MA, Turner DJ, Ning Z, Yusa K, Liang Q, Eckert S, Rad L, Fitzgerald TW, Craig NL, Bradley A: Mobilization of giant piggyBac transposons in the mouse genome. *Nucleic Acids Res* 2011, 39(22):e148.

Li R, Liang J, Ni S, Zhou T, Qing X, Li H, He W, Chen J, Li F, Zhuang Q et al: A mesenchymal-to-epithelial transition initiates and is required for the nuclear reprogramming of mouse fibroblasts. *Cell stem cell* 2010, 7(1):51-63.

- Li W, Wei W, Zhu S, Zhu J, Shi Y, Lin T, Hao E, Hayek A, Deng H, Ding S: Generation of rat and human induced pluripotent stem cells by combining genetic reprogramming and chemical inhibitors. *Cell stem cell* 2009, 4(1):16-19.
- Liang Q, Kong J, Stalker J, Bradley A: Chromosomal mobilization and reintegration of Sleeping Beauty and PiggyBac transposons. *Genesis* 2009, 47(6):404-408.
- Liao B, Bao X, Liu L, Feng S, Zovoilis A, Liu W, Xue Y, Cai J, Guo X, Qin B et al: MicroRNA cluster 302-367 enhances somatic cell reprogramming by accelerating a mesenchymal-to-epithelial transition. *The Journal of biological chemistry* 2011, 286(19):17359-17364.
- Liao J, Cui C, Chen S, Ren J, Chen J, Gao Y, Li H, Jia N, Cheng L, Xiao H et al: Generation of induced pluripotent stem cell lines from adult rat cells. *Cell stem cell* 2009, 4(1):11-15.
- Lin H, Shabbir A, Molnar M, Lee T: Stem cell regulatory function mediated by expression of a novel mouse Oct4 pseudogene. *Biochem Bioph Res Co* 2007, 355(1):111-116.
- Lin JJ, Li Y, Eppinga RD, Wang Q, Jin JP: Chapter 1: roles of caldesmon in cell motility and actin cytoskeleton remodeling. *International review of cell and molecular biology* 2009, 274:1-68.
- Lin S, Gaiano N, Culp P, Burns JC, Friedmann T, Yee JK, Hopkins N: Integration and germ-line transmission of a pseudotyped retroviral vector in zebrafish. *Science* 1994, 265(5172):666-669.
- Lister R, Pelizzola M, Kida YS, Hawkins RD, Nery JR, Hon G, Antosiewicz-Bourget J, O'Malley R, Castanon R, Klugman S et al: Hotspots of aberrant epigenomic reprogramming in human induced pluripotent stem cells. *Nature* 2011, 471(7336):68-73.
- Liu H, Ye Z, Kim Y, Sharkis S, Jang YY: Generation of endoderm-derived human induced pluripotent stem cells from primary hepatocytes. *Hepatology (Baltimore, Md)* 2010, 51(5):1810-1819.

Liu H, Zhu F, Yong J, Zhang P, Hou P, Li H, Jiang W, Cai J, Liu M, Cui K et al: Generation of induced pluripotent stem cells from adult rhesus monkey fibroblasts. *Cell stem cell* 2008, 3(6):587-590.

Liu XX, Nugoli M, Laferriere J, Saleh SM, Rodrigue-Gervais IG, Saleh M, Park M, Hallett MT, Muller WJ, Giguere V: Stromal retinoic acid receptor beta promotes mammary gland tumorigenesis. *P Natl Acad Sci USA* 2011, 108(2):774-779.

Lluis F, Ombrato L, Pedone E, Pepe S, Merrill BJ, Cosma MP: T-cell factor 3 (Tcf3) deletion increases somatic cell reprogramming by inducing epigenome modifications. *P Natl Acad Sci USA* 2011, 108(29):11912-11917.

Loewer S, Cabili MN, Guttman M, Loh YH, Thomas K, Park IH, Garber M, Curran M, Onder T, Agarwal S et al: Large intergenic non-coding RNA-RoR modulates reprogramming of human induced pluripotent stem cells. *Nat Genet* 2010, 42(12):1113-1117.

Loh YH, Agarwal S, Park IH, Urbach A, Huo H, Heffner GC, Kim K, Miller JD, Ng K, Daley GQ: Generation of induced pluripotent stem cells from human blood. *Blood* 2009, 113(22):5476-5479.

Loh YH, Wu Q, Chew JL, Vega VB, Zhang W, Chen X, Bourque G, George J, Leong B, Liu J et al: The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nat Genet* 2006, 38(4):431-440.

Lohle M, Hermann A, Glass H, Kempe A, Schwarz SC, Kim JB, Poulet C, Ravens U, Schwarz J, Scholer HR et al: Differentiation efficiency of induced pluripotent stem cells depends on the number of reprogramming factors. *Stem Cells* 2012, 30(3):570-579.

Lohnes D, Kastner P, Dierich A, Mark M, LeMeur M, Chambon P: Function of retinoic acid receptor gamma in the mouse. *Cell* 1993, 73(4):643-658.

Lohnes D, Mark M, Mendelsohn C, Dolle P, Dierich A, Gorry P, Gansmuller A, Chambon P: Function of the retinoic acid receptors (RARs) during development (I).

Craniofacial and skeletal abnormalities in RAR double mutants. *Development* 1994, 120(10):2723-2748.

Lufkin T, Lohnes D, Mark M, Dierich A, Gorry P, Gaub MP, LeMeur M, Chambon P: High postnatal lethality and testis degeneration in retinoic acid receptor alpha mutant mice. *P Natl Acad Sci USA* 1993, 90(15):7225-7229.

Luo GB, Ivics Z, Izsvak Z, Bradley A: Chromosomal transposition of a Tc1/mariner-like element in mouse embryonic stem cells. *P Natl Acad Sci USA* 1998, 95(18):10769-10773.

Luo J, Sucov HM, Bader JA, Evans RM, Giguere V: Compound mutants for retinoic acid receptor (RAR) beta and RAR alpha 1 reveal developmental functions for multiple RAR beta isoforms. *Mechanisms of development* 1996, 55(1):33-44.

Macfarlan TS, Gifford WD, Driscoll S, Lettieri K, Rowe HM, Bonanomi D, Firth A, Singer O, Trono D, Pfaff SL: Embryonic stem cell potency fluctuates with endogenous retrovirus activity. *Nature* 2012.

Maekawa M, Yamaguchi K, Nakamura T, Shibukawa R, Kodanaka I, Ichisaka T, Kawamura Y, Mochizuki H, Goshima N, Yamanaka S: Direct reprogramming of somatic cells is promoted by maternal transcription factor Glis1. *Nature* 2011, 474(7350):225-229.

Maherali N, Sridharan R, Xie W, Utikal J, Eminli S, Arnold K, Stadtfeld M, Yachechko R, Tchieu J, Jaenisch R et al: Directly reprogrammed fibroblasts show global epigenetic remodeling and widespread tissue contribution. *Cell stem cell* 2007, 1(1):55-70.

Maki N, Suetsugu-Maki R, Sano S, Nakamura K, Nishimura O, Tarui H, Del Rio-Tsonis K, Ohsumi K, Agata K, Tsonis PA: Oocyte-type linker histone B4 is required for transdifferentiation of somatic cells in vivo. *Faseb J* 2010, 24(9):3462-3467.

Mann KM, Ward JM, Yew CC, Kovochich A, Dawson DW, Black MA, Brett BT, Sheetz TE, Dupuy AJ, Chang DK et al: Sleeping Beauty mutagenesis reveals

cooperating mutations and pathways in pancreatic adenocarcinoma. *P Natl Acad Sci USA* 2012.

Mansour AA, Gafni O, Weinberger L, Zviran A, Ayyash M, Rais Y, Krupalnik V, Zerbib M, Amann-Zalcenstein D, Maza I et al: The H3K27 demethylase Utx regulates somatic and germ cell epigenetic reprogramming. *Nature* 2012, 488(7411):409-413.

Mansour AA, Gafni O, Weinberger L, Zviran A, Ayyash M, Rais Y, Krupalnik V, Zerbib M, Amann-Zalcenstein D, Maza I et al: The H3K27 demethylase Utx regulates somatic and germ cell epigenetic reprogramming. *Nature* 2012, 488(7411):409-413.

Marion RM, Strati K, Li H, Murga M, Blanco R, Ortega S, Fernandez-Capetillo O, Serrano M, Blasco MA: A p53-mediated DNA damage response limits reprogramming to ensure iPS cell genomic integrity. *Nature* 2009, 460(7259):1149-1153.

Mark M, Ghyselinck NB, Chambon P: Function of retinoic acid receptors during embryonic development. *Nuclear receptor signaling* 2009, 7:e002.

Marks H, Kalkan T, Menafra R, Denissov S, Jones K, Hofemeister H, Nichols J, Kranz A, Stewart AF, Smith A et al: The transcriptional and epigenomic foundations of ground state pluripotency. *Cell* 2012, 149(3):590-604.

Marson A, Foreman R, Chevalier B, Bilodeau S, Kahn M, Young RA, Jaenisch R: Wnt signaling promotes reprogramming of somatic cells to pluripotency. *Cell stem cell* 2008, 3(2):132-135.

Marson A, Levine SS, Cole MF, Frampton GM, Brambrink T, Johnstone S, Guenther MG, Johnston WK, Wernig M, Newman J et al: Connecting microRNA genes to the core transcriptional regulatory circuitry of embryonic stem cells. *Cell* 2008, 134(3):521-533.

Marston SB, Lehman W: Caldesmon Is a Ca²⁺-Regulatory Component of Native Smooth-Muscle Thin-Filaments. *Biochem J* 1985, 231(3):517-522.

Martin GR: Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *P Natl Acad Sci USA* 1981, 78(12):7634-7638.

Martin GR, Evans MJ: The morphology and growth of a pluripotent teratocarcinoma cell line and its derivatives in tissue culture. *Cell* 1974, 2(3):163-172.

Masaki H, Ishikawa T, Takahashi S, Okumura M, Sakai N, Haga M, Kominami K, Migita H, McDonald F, Shimada F et al: Heterogeneity of pluripotent marker gene expression in colonies generated in human iPS cell induction culture. *Stem Cell Res* 2008, 1(2):105-115.

Masia S, Alvarez S, de Lera AR, Baretino D: Rapid, nongenomic actions of retinoic acid on phosphatidylinositol-3-kinase signaling pathway mediated by the retinoic acid receptor. *Mol Endocrinol* 2007, 21(10):2391-2402.

Masui S, Nakatake Y, Toyooka Y, Shimosato D, Yagi R, Takahashi K, Okochi H, Okuda A, Matoba R, Sharov AA et al: Pluripotency governed by Sox2 via regulation of Oct3/4 expression in mouse embryonic stem cells. *Nat Cell Biol* 2007, 9(6):625-635.

Matkovits T, Christakos S: Ligand occupancy is not required for vitamin D receptor and retinoid receptor-mediated transcriptional activation. *Mol Endocrinol* 1995, 9(2):232-242.

Matsui Y, Toksoz D, Nishikawa S, Nishikawa SI, Williams D, Zsebo K, Hogan BLM: Effect of Steel Factor and Leukemia Inhibitory Factor on Murine Primordial Germ-Cells in Culture. *Nature* 1991, 353(6346):750-752.

Matsui Y, Zsebo K, Hogan BLM: Derivation of Pluripotential Embryonic Stem-Cells from Murine Primordial Germ-Cells in Culture. *Cell* 1992, 70(5):841-847.

Matsui Y, Zsebo K, Hogan BLM: Derivation of Pluripotential Embryonic Stem-Cells from Murine Primordial Germ-Cells in Culture. *Cell* 1992, 70(5):841-847.

Mayshar Y, Ben-David U, Lavon N, Biancotti JC, Yakir B, Clark AT, Plath K, Lowry WE, Benvenisty N: Identification and classification of chromosomal aberrations in human induced pluripotent stem cells. *Cell stem cell* 2010, 7(4):521-531.

McClintock B: The origin and behavior of mutable loci in maize. *P Natl Acad Sci USA* 1950, 36(6):344-355.

McGrath J, Solter D: Nuclear transplantation in the mouse embryo by microsurgery and cell fusion. *Science* 1983, 220(4603):1300-1302.

McMahon AP, Bradley A: The Wnt-1 (int-1) proto-oncogene is required for development of a large region of the mouse brain. *Cell* 1990, 62(6):1073-1085.

McNamara S, Wang H, Hanna N, Miller WH, Jr.: Topoisomerase IIbeta negatively modulates retinoic acid receptor alpha function: a novel mechanism of retinoic acid resistance. *Mol Cell Biol* 2008, 28(6):2066-2077.

Medvedev A, Yan ZH, Hirose T, Giguere V, Jetten AM: Cloning of a cDNA encoding the murine orphan receptor RZR/ROR gamma and characterization of its response element. *Gene* 1996, 181(1-2):199-206.

Meeker TC, Nagarajan L, ar-Rushdi A, Croce CM: Cloning and characterization of the human PIM-1 gene: a putative oncogene related to the protein kinases. *Journal of cellular biochemistry* 1987, 35(2):105-112.

Meir YJJ, Weirauch MT, Yang HS, Chung PC, Yu RK, Wu SCY: Genome-wide target profiling of piggyBac and Tol2 in HEK 293: pros and cons for gene discovery and gene therapy. *Bmc Biotechnol* 2011, 11.

Meissner A, Mikkelsen TS, Gu H, Wernig M, Hanna J, Sivachenko A, Zhang X, Bernstein BE, Nusbaum C, Jaffe DB et al: Genome-scale DNA methylation maps of pluripotent and differentiated cells. *Nature* 2008, 454(7205):766-770.

Meissner A, Wernig M, Jaenisch R: Direct reprogramming of genetically unmodified fibroblasts into pluripotent stem cells. *Nat Biotechnol* 2007, 25(10):1177-1181.

Melnikov A, Murugan A, Zhang X, Tesileanu T, Wang L, Rogov P, Feizi S, Gnirke A, Callan CG, Jr., Kinney JB et al: Systematic dissection and optimization of inducible enhancers in human cells using a massively parallel reporter assay. *Nat Biotechnol* 2012, 30(3):271-277.

Mendelsohn C, Lohnes D, Decimo D, Lufkin T, LeMeur M, Chambon P, Mark M: Function of the retinoic acid receptors (RARs) during development (II). Multiple abnormalities at various stages of organogenesis in RAR double mutants. *Development* 1994, 120(10):2749-2771.

Mermod N, O'Neill EA, Kelly TJ, Tjian R: The proline-rich transcriptional activator of CTF/NF-I is distinct from the replication and DNA binding domain. *Cell* 1989, 58(4):741-753.

Metallo CM, Ji L, de Pablo JJ, Palecek SP: Retinoic acid and bone morphogenetic protein signaling synergize to efficiently direct epithelial differentiation of human embryonic stem cells. *Stem Cells* 2008, 26(2):372-380.

Metzker ML: Sequencing technologies - the next generation. *Nat Rev Genet* 2010, 11(1):31-46.

Meyer T, Kneissel M, Mariani J, Fournier B: In vitro and in vivo evidence for orphan nuclear receptor RORalpha function in bone metabolism. *P Natl Acad Sci USA* 2000, 97(16):9197-9202.

Mikkelsen S, Berne B, Staberg B, Vahlquist A: Potentiating effect of dietary vitamin A on photocarcinogenesis in hairless mice. *Carcinogenesis* 1998, 19(4):663-666.

Mikkelsen TS, Hanna J, Zhang X, Ku M, Wernig M, Schorderet P, Bernstein BE, Jaenisch R, Lander ES, Meissner A: Dissecting direct reprogramming through integrative genomic analysis. *Nature* 2008, 454(7200):49-55.

Minucci S, Botquin V, Yeom YI, Dey A, Sylvester I, Zand DJ, Ohbo K, Ozato K, Scholer HR: Retinoic acid-mediated down-regulation of Oct3/4 coincides with the loss of promoter occupancy in vivo. *The EMBO journal* 1996, 15(4):888-899.

Miyoshi N, Ishii H, Nagano H, Haraguchi N, Dewi DL, Kano Y, Nishikawa S, Tanemura M, Mimori K, Tanaka F et al: Reprogramming of mouse and human cells to pluripotency using mature microRNAs. *Cell stem cell* 2011, 8(6):633-638.

Mohan M, Malayer JR, Geisert RD, Morgan GL: Expression of retinol-binding protein messenger RNA and retinoic acid receptors in preattachment bovine embryos. *Molecular reproduction and development* 2001, 60(3):289-296.

Mollersen L, Paulsen JE, Olstorn HB, Knutsen HK, Alexander J: Dietary retinoic acid supplementation stimulates intestinal tumour formation and growth in multiple intestinal neoplasia (Min)/+ mice. *Carcinogenesis* 2004, 25(1):149-153.

Montserrat N, de Onate L, Garreta E, Gonzalez F, Adamo A, Eguizabal C, Hafner S, Vassena R, Belmonte JC: Generation of feeder free pig induced pluripotent stem cells without Pou5f1. *Cell transplantation* 2011.

Moretti A, Bellin M, Welling A, Jung CB, Lam JT, Bott-Flugel L, Dorn T, Goedel A, Hohnke C, Hofmann F et al: Patient-specific induced pluripotent stem-cell models for long-QT syndrome. *The New England journal of medicine* 2011, 363(15):1397-1409.

Morishita K, Parker DS, Mucenski ML, Jenkins NA, Copeland NG, Ihle JN: Retroviral activation of a novel gene encoding a zinc finger protein in IL-3-dependent myeloid leukemia cell lines. *Cell* 1988, 54(6):831-840.

Morriss-Kay GM, Murphy P, Hill RE, Davidson DR: Effects of retinoic acid excess on expression of Hox-2.9 and Krox-20 and on morphological segmentation in the hindbrain of mouse embryos. *The EMBO journal* 1991, 10(10):2985-2995.

Mucenski ML, Taylor BA, Ihle JN, Hartley JW, Morse HC, 3rd, Jenkins NA, Copeland NG: Identification of a common ecotropic viral integration site, Evi-1, in the DNA of AKXD murine myeloid tumors. *Mol Cell Biol* 1988, 8(1):301-308.

Nagamatsu G, Kosaka T, Kawasumi M, Kinoshita T, Takubo K, Akiyama H, Sudo T, Kobayashi T, Oya M, Suda T: A germ cell-specific gene, Prmt5, works in somatic cell reprogramming. *The Journal of biological chemistry* 2011, 286(12):10641-10648.

Nagy A, Gocza E, Diaz EM, Prideaux VR, Ivanyi E, Markkula M, Rossant J: Embryonic stem cells alone are able to support fetal development in the mouse. *Development* 1990, 110(3):815-821.

Nagy A, Rossant J, Nagy R, Abramow-Newerly W, Roder JC: Derivation of completely cell culture-derived mice from early-passage embryonic stem cells. *P Natl Acad Sci USA* 1993, 90(18):8424-8428.

Nakagawa M, Koyanagi M, Tanabe K, Takahashi K, Ichisaka T, Aoi T, Okita K, Mochiduki Y, Takizawa N, Yamanaka S: Generation of induced pluripotent stem cells without Myc from mouse and human fibroblasts. *Nat Biotechnol* 2008, 26(1):101-106.

Nakagawa S, Watanabe M, Inoue Y: Prominent expression of nuclear hormone receptor ROR alpha in Purkinje cells from early development. *Neuroscience research* 1997, 28(2):177-184.

Nakagawa Y, O'Leary DD: Dynamic patterned expression of orphan nuclear receptor genes RORalpha and RORbeta in developing mouse forebrain. *Developmental neuroscience* 2003, 25(2-4):234-244.

Narlikar GJ, Fan HY, Kingston RE: Cooperation between complexes that regulate chromatin structure and transcription. *Cell* 2002, 108(4):475-487.

Narsinh KH, Sun N, Sanchez-Freire V, Lee AS, Almeida P, Hu SJ, Jan T, Wilson KD, Leong D, Rosenberg J et al: Single cell transcriptional profiling reveals heterogeneity of human induced pluripotent stem cells. *J Clin Invest* 2011, 121(3):1217-1221.

Neel BG, Hayward WS, Robinson HL, Fang J, Astrin SM: Avian leukosis virus-induced tumors have common proviral integration sites and synthesize discrete new RNAs: oncogenesis by promoter insertion. *Cell* 1981, 23(2):323-334.

Nichols J, Chambers I, Taga T, Smith A: Physiological rationale for responsiveness of mouse embryonic stem cells to gp130 cytokines. *Development* 2001, 128(12):2333-2339.

Nichols J, Jones K, Phillips JM, Newland SA, Roode M, Mansfield W, Smith A, Cooke A: Validated germline-competent embryonic stem cell lines from nonobese diabetic mice. *Nature medicine* 2009, 15(7):814-818.

Nichols J, Silva J, Roode M, Smith A: Suppression of Erk signalling promotes ground state pluripotency in the mouse embryo. *Development* 2009, 136(19):3215-3222.

Nichols J, Smith A: Naive and primed pluripotent states. *Cell stem cell* 2009, 4(6):487-492.

Nichols J, Zevnik B, Anastassiadis K, Niwa H, Klewe-Nebenius D, Chambers I, Scholer H, Smith A: Formation of pluripotent stem cells in the mammalian embryo depends on the POU transcription factor Oct4. *Cell* 1998, 95(3):379-391.

Nielsen R, Pedersen TA, Hagenbeek D, Moulos P, Siersbaek R, Megens E, Denissov S, Borgesen M, Francoijs KJ, Mandrup S et al: Genome-wide profiling of PPARgamma:RXR and RNA polymerase II occupancy reveals temporal activation of distinct metabolic pathways and changes in RXR dimer composition during adipogenesis. *Gene Dev* 2008, 22(21):2953-2967.

Niwa H, Burdon T, Chambers I, Smith A: Self-renewal of pluripotent embryonic stem cells is mediated via activation of STAT3. *Gene Dev* 1998, 12(13):2048-2060.

Niwa H, Miyazaki J, Smith AG: Quantitative expression of Oct-3/4 defines differentiation, dedifferentiation or self-renewal of ES cells. *Nat Genet* 2000, 24(4):372-376.

Niwa H, Yamamura K, Miyazaki J: Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* 1991, 108(2):193-199.

Noggle S, Fung HL, Gore A, Martinez H, Satriani KC, Prosser R, Oum K, Paull D, Druckenmiller S, Freeby M et al: Human oocytes reprogram somatic cells to a pluripotent state. *Nature* 2011, 478(7367):70-75.

Nordhoff V, Hubner K, Bauer A, Orlova I, Malapetsa A, Scholer HR: Comparative analysis of human, bovine, and murine Oct-4 upstream promoter sequences. *Mamm Genome* 2001, 12(4):309-317.

Nori S, Okada Y, Yasuda A, Tsuji O, Takahashi Y, Kobayashi Y, Fujiyoshi K, Koike M, Uchiyama Y, Ikeda E et al: Grafted human-induced pluripotent stem-cell-derived neurospheres promote motor functional recovery after spinal cord injury in mice. *P Natl Acad Sci USA* 2011, 108(40):16825-16830.

Notarianni E, Galli C, Laurie S, Moor RM, Evans MJ: Derivation of pluripotent, embryonic cell lines from the pig and sheep. *Journal of reproduction and fertility* 1991, 43:255-260.

Notarianni E, Laurie S, Moor RM, Evans MJ: Maintenance and differentiation in culture of pluripotential embryonic cell lines from pig blastocysts. *Journal of reproduction and fertility* 1990, 41:51-56.

Novy RE, Lin JLC, Lin JJC: Characterization of Cdna Clones Encoding a Human Fibroblast Caldesmon Isoform and Analysis of Caldesmon Expression in Normal and Transformed-Cells. *J Biol Chem* 1991, 266(25):16917-16924.

O'Brien RN, Shen Z, Tachikawa K, Lee PA, Briggs SP: Quantitative proteome analysis of pluripotent cells by iTRAQ mass tagging reveals post-transcriptional regulation of proteins required for ES cell self-renewal. *Molecular & cellular proteomics : MCP* 2010, 9(10):2238-2251.

Ohi Y, Qin H, Hong C, Blouin L, Polo JM, Guo T, Qi Z, Downey SL, Manos PD, Rossi DJ et al: Incomplete DNA methylation underlies a transcriptional memory of somatic cells in human iPS cells. *Nat Cell Biol* 2011, 13(5):541-549.

Okada Y, Shimazaki T, Sobue G, Okano H: Retinoic-acid-concentration-dependent acquisition of neural cell identity during in vitro differentiation of mouse embryonic stem cells. *Developmental biology* 2004, 275(1):124-142.

Okamoto K, Okazawa H, Okuda A, Sakai M, Muramatsu M, Hamada H: A novel octamer binding transcription factor is differentially expressed in mouse embryonic cells. *Cell* 1990, 60(3):461-472.

Okazawa H, Okamoto K, Ishino F, Ishino-Kaneko T, Takeda S, Toyoda Y, Muramatsu M, Hamada H: The oct3 gene, a gene for an embryonic transcription factor, is controlled by a retinoic acid repressible enhancer. *The EMBO journal* 1991, 10(10):2997-3005.

Okita K, Ichisaka T, Yamanaka S: Generation of germline-competent induced pluripotent stem cells. *Nature* 2007, 448(7151):313-317.

Okita K, Nakagawa M, Hyenjong H, Ichisaka T, Yamanaka S: Generation of mouse induced pluripotent stem cells without viral vectors. *Science* 2008, 322(5903):949-953.

O'Leary T, Heindryckx B, Lierman S, van Bruggen D, Goeman JJ, Vandewoestyne M, Deforce D, Lopes SMCD, De Sutter P: Tracking the progression of the human inner cell mass during embryonic stem cell derivation. *Nat Biotechnol* 2012, 30(3):278-+.

Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, Keogh JP, Meyskens FL, Valanis B, Williams JH et al: Effects of a combination of beta carotene and vitamin A on lung cancer and cardiovascular disease. *New Engl J Med* 1996, 334(18):1150-1155.

Pain B, Clark ME, Shen M, Nakazawa H, Sakurai M, Samarut J, Etches RJ: Long-term in vitro culture and characterisation of avian embryonic stem cells with multiple morphogenetic potentialities. *Development* 1996, 122(8):2339-2348.

Palmieri SL, Peter W, Hess H, Scholer HR: Oct-4 transcription factor is differentially expressed in the mouse embryo during establishment of the first two extraembryonic cell lineages involved in implantation. *Developmental biology* 1994, 166(1):259-267.

Panagopoulos I, Moller E, Collin A, Mertens F: The POU5F1P1 pseudogene encodes a putative protein similar to POU5F1 isoform 1. *Oncol Rep* 2008, 20(5):1029-1033.

Pander CH: Beiträge zur Entwicklungsgeschichte des Hühnchens im Eye: Würzburg; 1817.

Papalopulu N, Clarke JD, Bradley L, Wilkinson D, Krumlauf R, Holder N: Retinoic acid causes abnormal development and segmental patterning of the anterior hindbrain in *Xenopus* embryos. *Development* 1991, 113(4):1145-1158.

Papamichos SI, Kotoula V, Tarlatzis BC, Agorastos T, Papazisis K, Lambropoulos AF: OCT4B1 isoform: the novel OCT4 alternative spliced variant as a putative marker of stemness. *Mol Hum Reprod* 2009, 15(5):269-270.

Pardo M, Lang B, Yu L, Prosser H, Bradley A, Babu MM, Choudhary J: An expanded Oct4 interaction network: implications for stem cell biology, development, and disease. *Cell stem cell* 2010, 6(4):382-395.

Park IH, Arora N, Huo H, Maherali N, Ahfeldt T, Shimamura A, Lensch MW, Cowan C, Hochedlinger K, Daley GQ: Disease-specific induced pluripotent stem cells. *Cell* 2008, 134(5):877-886.

Patwardhan RP, Hiatt JB, Witten DM, Kim MJ, Smith RP, May D, Lee C, Andrie JM, Lee SI, Cooper GM et al: Massively parallel functional dissection of mammalian enhancers in vivo. *Nat Biotechnol* 2012, 30(3):265-270.

Pavlath GK, Blau HM: Expression of muscle genes in heterokaryons depends on gene dosage. *The Journal of cell biology* 1986, 102(1):124-130.

Pease S, Braghetta P, Gearing D, Grail D, Williams RL: Isolation of embryonic stem (ES) cells in media supplemented with recombinant leukemia inhibitory factor (LIF). *Developmental biology* 1990, 141(2):344-352.

Pereira CF, Terranova R, Ryan NK, Santos J, Morris KJ, Cui W, Merkenschlager M, Fisher AG: Heterokaryon-based reprogramming of human B lymphocytes for pluripotency requires Oct4 but not Sox2. *PLoS genetics* 2008, 4(9):e1000170.

Pesce M, Wang X, Wolgemuth DJ, Scholer H: Differential expression of the Oct-4 transcription factor during mouse germ cell differentiation. *Mechanisms of development* 1998, 71(1-2):89-98.

Petkovich M, Brand NJ, Krust A, Chambon P: A human retinoic acid receptor which belongs to the family of nuclear receptors. *Nature* 1987, 330(6147):444-450.

Petroski MD, Deshaies RJ: Function and regulation of Cullin-RING ubiquitin ligases. *Nat Rev Mol Cell Bio* 2005, 6(1):9-20.

Phanstiel DH, Brumbaugh J, Wenger CD, Tian S, Probasco MD, Bailey DJ, Swaney DL, Tervo MA, Bolin JM, Ruotti V et al: Proteomic and phosphoproteomic comparison of human ES and iPS cells. *Nature methods* 2011, 8(10):821-827.

Picard E, Seguin C, Monhoven N, Rochette-Egly C, Siat J, Borrelly J, Martinet Y, Martinet N, Vignaud JM: Expression of retinoid receptor genes and proteins in non-small-cell lung cancer. *Journal of the National Cancer Institute* 1999, 91(12):1059-1066.

Pikarsky E, Sharir H, Ben-Shushan E, Bergman Y: Retinoic acid represses Oct-3/4 gene expression through several retinoic acid-responsive elements located in the promoter-enhancer region. *Mol Cell Biol* 1994, 14(2):1026-1038.

Polo JM, Liu S, Figueroa ME, Kulalert W, Eminli S, Tan KY, Apostolou E, Stadtfeld M, Li Y, Shioda T et al: Cell type of origin influences the molecular and functional properties of mouse induced pluripotent stem cells. *Nat Biotechnol* 2010, 28(8):848-855.

Prasad SM, Czepiel M, Cetinkaya C, Smigielska K, Weli SC, Lysdahl H, Gabrielsen A, Petersen K, Ehlers N, Fink T et al: Continuous hypoxic culturing maintains activation of Notch and allows long-term propagation of human embryonic stem cells without spontaneous differentiation. *Cell proliferation* 2009, 42(1):63-74.

Prigione A, Fauler B, Lurz R, Lehrach H, Adjaye J: The Senescence-Related Mitochondrial/Oxidative Stress Pathway is Repressed in Human Induced Pluripotent Stem Cells. *Stem Cells* 2010, 28(4):721-733.

Rad R, Rad L, Wang W, Cadinanos J, Vassiliou G, Rice S, Campos LS, Yusa K, Banerjee R, Li MA et al: PiggyBac transposon mutagenesis: a tool for cancer gene discovery in mice. *Science* 2010, 330(6007):1104-1107.

Rahl PB, Lin CY, Seila AC, Flynn RA, McCuine S, Burge CB, Sharp PA, Young RA: c-Myc regulates transcriptional pause release. *Cell* 2010, 141(3):432-445.

Ramirez-Solis R, Davis AC, Bradley A: Gene targeting in embryonic stem cells. *Methods in enzymology* 1993, 225:855-878.

Rashid ST, Corbineau S, Hannan N, Marciniak SJ, Miranda E, Alexander G, Huang-Doran I, Griffin J, Ahrlund-Richter L, Skepper J et al: Modeling inherited metabolic disorders of the liver using human induced pluripotent stem cells. *The Journal of clinical investigation* 2010, 120(9):3127-3136.

Raya A, Rodriguez-Piza I, Guenechea G, Vassena R, Navarro S, Barrero MJ, Consiglio A, Castella M, Rio P, Sleep E et al: Disease-corrected haematopoietic progenitors from Fanconi anaemia induced pluripotent stem cells. *Nature* 2009, 460(7251):53-59.

Redmer T, Diecke S, Grigoryan T, Quiroga-Negreira A, Birchmeier W, Besser D: E-cadherin is crucial for embryonic stem cell pluripotency and can replace OCT4 during somatic cell reprogramming. *EMBO reports* 2011, 12(7):720-726.

Rhinn M, Dolle P: Retinoic acid signalling during development. *Development* 2012, 139(5):843-858.

Rochette-Egly C, Adam S, Rossignol M, Egly JM, Chambon P: Stimulation of RAR alpha activation function AF-1 through binding to the general transcription factor TFIID and phosphorylation by CDK7. *Cell* 1997, 90(1):97-107.

Rochette-Egly C, Germain P: Dynamic and combinatorial control of gene expression by nuclear retinoic acid receptors (RARs). *Nuclear receptor signaling* 2009, 7:e005.

Rodda DJ, Chew JL, Lim LH, Loh YH, Wang B, Ng HH, Robson P: Transcriptional regulation of nanog by OCT4 and SOX2. *The Journal of biological chemistry* 2005, 280(26):24731-24737.

Roh TY, Cuddapah S, Zhao K: Active chromatin domains are defined by acetylation islands revealed by genome-wide mapping. *Gene Dev* 2005, 19(5):542-552.

Romano RA, Li H, Tummala R, Maul R, Sinha S: Identification of Basonuclin2, a DNA-binding zinc-finger protein expressed in germ tissues and skin keratinocytes. *Genomics* 2004, 83(5):821-833.

Roode M, Blair K, Snell P, Elder K, Marchant S, Smith A, Nichols J: Human hypoblast formation is not dependent on FGF signalling. *Developmental biology* 2012, 361(2):358-363.

Rosenthal MD, Wishnow RM, Sato GH: In vitro growth and differentiation of clonal populations of multipotential mouse cells derived from a transplantable testicular teratocarcinoma. *Journal of the National Cancer Institute* 1970, 44(5):1001-1014.

Rosner MH, Vigano MA, Ozato K, Timmons PM, Poirier F, Rigby PW, Staudt LM: A POU-domain transcription factor in early stem cells and germ cells of the mammalian embryo. *Nature* 1990, 345(6277):686-692.

Ross DT, Scherf U, Eisen MB, Perou CM, Rees C, Spellman P, Iyer V, Jeffrey SS, Van de Rijn M, Waltham M et al: Systematic variation in gene expression patterns in human cancer cell lines. *Nat Genet* 2000, 24(3):227-235.

Ross-Innes CS, Stark R, Holmes KA, Schmidt D, Spyrou C, Russell R, Massie CE, Vowler SL, Eldridge M, Carroll JS: Cooperative interaction between retinoic acid receptor-alpha and estrogen receptor in breast cancer. *Gene Dev* 2010, 24(2):171-182.

Roux W: Beitrage zur Entwicklungsmechanik des Embryo. *Virchows Arch Pathol Anat Physiol Klin Med* 1888, 144:113-153.

Rowland BD, Bernards R, Peeper DS: The KLF4 tumour suppressor is a transcriptional repressor of p53 that acts as a context-dependent oncogene. *Nat Cell Biol* 2005, 7(11):1074-1082.

Ruberte E, Dolle P, Chambon P, Morriss-Kay G: Retinoic acid receptors and cellular retinoid binding proteins. II. Their differential pattern of transcription during early morphogenesis in mouse embryos. *Development* 1991, 111(1):45-60.

Rubin GM, Kidwell MG, Bingham PM: The molecular basis of P-M hybrid dysgenesis: the nature of induced mutations. *Cell* 1982, 29(3):987-994.

Rushlow KE, Lautenberger JA, Reddy EP, Souza LM, Baluda MA, Chirikjian JG, Papas TS: Nucleotide sequence analysis of the long terminal repeat of avian myeloblastosis virus and adjacent host sequences. *Journal of virology* 1982, 42(3):840-846.

Saito S, Strelchenko N, Niemann H: Bovine Embryonic Stem Cell-Like Cell-Lines Cultured over Several Passages. *Roux Arch Dev Biol* 1992, 201(3):134-141.

Samavarchi-Tehrani P, Golipour A, David L, Sung HK, Beyer TA, Datti A, Woltjen K, Nagy A, Wrana JL: Functional genomics reveals a BMP-driven mesenchymal-to-epithelial transition in the initiation of somatic cell reprogramming. *Cell stem cell* 2010, 7(1):64-77.

Sanger F, Air GM, Barrell BG, Brown NL, Coulson AR, Fiddes CA, Hutchison CA, Slocombe PM, Smith M: Nucleotide sequence of bacteriophage phi X174 DNA. *Nature* 1977, 265(5596):687-695.

Saxe JP, Tomilin A, Scholer HR, Plath K, Huang J: Post-Translational Regulation of Oct4 Transcriptional Activity. *PloS one* 2009, 4(2).

Schaeren-Wiemers N, Andre E, Kapfhammer JP, Becker-Andre M: The expression pattern of the orphan nuclear receptor RORbeta in the developing and adult rat nervous system suggests a role in the processing of sensory information and in circadian rhythm. *The European journal of neuroscience* 1997, 9(12):2687-2701.

Schnetz MP, Handoko L, Akhtar-Zaidi B, Bartels CF, Pereira CF, Fisher AG, Adams DJ, Flicek P, Crawford GE, Laframboise T et al: CHD7 targets active gene enhancer elements to modulate ES cell-specific gene expression. *PLoS genetics* 2010, 6(7):e1001023.

Scholer HR, Dressler GR, Balling R, Rohdewohld H, Gruss P: Oct-4: a germline-specific transcription factor mapping to the mouse t-complex. *The EMBO journal* 1990, 9(7):2185-2195.

Scholer HR, Hatzopoulos AK, Balling R, Suzuki N, Gruss P: A family of octamer-specific proteins present during mouse embryogenesis: evidence for germline-specific expression of an Oct factor. *The EMBO journal* 1989, 8(9):2543-2550.

Scholer HR, Ruppert S, Suzuki N, Chowdhury K, Gruss P: New type of POU domain in germ line-specific protein Oct-4. *Nature* 1990, 344(6265):435-439.

Schoorlemmer J, van Puijenbroek A, van Den Eijnden M, Jonk L, Pals C, Kruijjer W: Characterization of a negative retinoic acid response element in the murine Oct4 promoter. *Mol Cell Biol* 1994, 14(2):1122-1136.

Schuldiner M, Eiges R, Eden A, Yanuka O, Itskovitz-Eldor J, Goldstein RS, Benvenisty N: Induced neuronal differentiation of human embryonic stem cells. *Brain research* 2001, 913(2):201-205.

Selten G, Cuypers HT, Boelens W, Robanus-Maandag E, Verbeek J, Domen J, van Beveren C, Berns A: The primary structure of the putative oncogene pim-1 shows extensive homology with protein kinases. *Cell* 1986, 46(4):603-611.

Sharon E, Kalma Y, Sharp A, Raveh-Sadka T, Levo M, Zeevi D, Keren L, Yakhini Z, Weinberger A, Segal E: Inferring gene regulatory logic from high-throughput measurements of thousands of systematically designed promoters. *Nat Biotechnol* 2012.

Shi Y, Do JT, Despons C, Hahm HS, Scholer HR, Ding S: A combined chemical and genetic approach for the generation of induced pluripotent stem cells. *Cell stem cell* 2008, 2(6):525-528.

Shi Y, Kirwan P, Smith J, Maclean G, Orkin SH, Livesey FJ: A Human Stem Cell Model of Early Alzheimer's Disease Pathology in Down Syndrome. *Science translational medicine* 2012.

Shiotsugu J, Katsuyama Y, Arima K, Baxter A, Koide T, Song J, Chandraratna RA, Blumberg B: Multiple points of interaction between retinoic acid and FGF signaling during embryonic axis formation. *Development* 2004, 131(11):2653-2667.

Si J, Mueller L, Collins SJ: CaMKII regulates retinoic acid receptor transcriptional activity and the differentiation of myeloid leukemia cells. *The Journal of clinical investigation* 2007, 117(5):1412-1421.

Silva J, Barrandon O, Nichols J, Kawaguchi J, Theunissen TW, Smith A: Promotion of reprogramming to ground state pluripotency by signal inhibition. *PLoS biology* 2008, 6(10):e253.

Silva J, Chambers I, Pollard S, Smith A: Nanog promotes transfer of pluripotency after cell fusion. *Nature* 2006, 441(7096):997-1001.

Silva J, Nichols J, Theunissen TW, Guo G, van Oosten AL, Barrandon O, Wray J, Yamanaka S, Chambers I, Smith A: Nanog Is the Gateway to the Pluripotent Ground State. *Cell* 2009, 138(4):722-737.

Simandi Z, Balint BL, Poliska S, Ruhl R, Nagy L: Activation of retinoic acid receptor signaling coordinates lineage commitment of spontaneously differentiating mouse embryonic stem cells in embryoid bodies. *FEBS letters* 2010, 584(14):3123-3130.

Singh H, Sen R, Baltimore D, Sharp PA: A nuclear factor that binds to a conserved sequence motif in transcriptional control elements of immunoglobulin genes. *Nature* 1986, 319(6049):154-158.

Singh N, George A, Sharma R, Singla SK, Palta P, Manik R, Chauhan MS, Singh D: Characterization of POU5F1 (OCT4) gene and its promoter in buffalo ESC-like cells identifies multiple transcription start sites and expression of four pseudogenes. *Gene* 2012, 491(2):165-172.

Singhal N, Graumann J, Wu G, Arauzo-Bravo MJ, Han DW, Greber B, Gentile L, Mann M, Scholer HR: Chromatin-Remodeling Components of the BAF Complex Facilitate Reprogramming. *Cell* 2010, 141(6):943-955.

Sive HL, Draper BW, Harland RM, Weintraub H: Identification of a retinoic acid-sensitive period during primary axis formation in *Xenopus laevis*. *Gene Dev* 1990, 4(6):932-942.

Smith AG, Heath JK, Donaldson DD, Wong GG, Moreau J, Stahl M, Rogers D: Inhibition of pluripotential embryonic stem cell differentiation by purified polypeptides. *Nature* 1988, 336(6200):688-690.

Smith AG, Hooper ML: Buffalo rat liver cells produce a diffusible activity which inhibits the differentiation of murine embryonal carcinoma and embryonic stem cells. *Developmental biology* 1987, 121(1):1-9.

Smithies O, Gregg RG, Boggs SS, Koralewski MA, Kucherlapati RS: Insertion of DNA sequences into the human chromosomal beta-globin locus by homologous recombination. *Nature* 1985, 317(6034):230-234.

Sobue K, Tanaka T, Kanda K, Ashino N, Kakiuchi S: Purification and Characterization of Caldesmon77 - a Calmodulin-Binding Protein That Interacts with Actin-Filaments from Bovine Adrenal-Medulla. *P Natl Acad Sci USA* 1985, 82(15):5025-5029.

Solt LA, Kumar N, Nuhant P, Wang YJ, Lauer JL, Liu J, Istrate MA, Kamenecka TM, Roush WR, Vidovic D et al: Suppression of T(H)17 differentiation and autoimmunity by a synthetic ROR ligand. *Nature* 2011, 472(7344):491-U547.

Solter D, Skreb N, Damjanov I: Extrauterine growth of mouse egg-cylinders results in malignant teratoma. *Nature* 1970, 227(5257):503-504.

Sommer CA, Stadtfeld M, Murphy GJ, Hochedlinger K, Kotton DN, Mostoslavsky G: Induced pluripotent stem cell generation using a single lentiviral stem cell cassette. *Stem Cells* 2009, 27(3):543-549.

Song HL, Ramus SJ, Tyrer J, Bolton KL, Gentry-Maharaj A, Wozniak E, Anton-Culver H, Chang-Claude J, Cramer DW, DiCioccio R et al: A genome-wide association study identifies a new ovarian cancer susceptibility locus on 9p22.2. *Nat Genet* 2009, 41(9):996-U960.

Song K, Nam YJ, Luo X, Qi X, Tan W, Huang GN, Acharya A, Smith CL, Tallquist MD, Neilson EG et al: Heart repair by reprogramming non-myocytes with cardiac transcription factors. *Nature* 2012, 485(7400):599-604.

Souza LM, Komaromy MC, Baluda MA: Identification of a proviral genome associated with avian myeloblastic leukemia. *P Natl Acad Sci USA* 1980, 77(5):3004-3008.

Spemann H: Die Entwicklung seitlicher und dorso-ventraler Keimhalften bei verzogter Kernversorgung. *Z Wiss Zool* 1928, 132:105–134

Spits C, Mateizel I, Geens M, Mertzaniidou A, Staessen C, Vandesselde Y, Van der Elst J, Liebaers I, Sermon K: Recurrent chromosomal abnormalities in human embryonic stem cells. *Nat Biotechnol* 2008, 26(12):1361-1363.

Sridharan R, Tchieu J, Mason MJ, Yachechko R, Kuoy E, Horvath S, Zhou Q, Plath K: Role of the murine reprogramming factors in the induction of pluripotency. *Cell* 2009, 136(2):364-377.

Stadtfeld M, Apostolou E, Akutsu H, Fukuda A, Follett P, Natesan S, Kono T, Shioda T, Hochedlinger K: Aberrant silencing of imprinted genes on chromosome 12qF1 in mouse induced pluripotent stem cells. *Nature* 2010, 465(7295):175-181.

Stadtfeld M, Nagaya M, Utikal J, Weir G, Hochedlinger K: Induced pluripotent stem cells generated without viral integration. *Science* 2008, 322(5903):945-949.

Starr TK, Allaei R, Silverstein KA, Staggs RA, Sarver AL, Bergemann TL, Gupta M, O'Sullivan MG, Matise I, Dupuy AJ et al: A transposon-based genetic screen in mice identifies genes altered in colorectal cancer. *Science* 2009, 323(5922):1747-1750.

Staudt LM, Singh H, Sen R, Wirth T, Sharp PA, Baltimore D: A lymphoid-specific protein binding to the octamer motif of immunoglobulin genes. *Nature* 1986, 323(6089):640-643.

Stevens LC: The development of transplantable teratocarcinomas from intratesticular grafts of pre- and postimplantation mouse embryos. *Developmental biology* 1970, 21(3):364-382.

Stewart CL, Kaspar P, Brunet LJ, Bhatt H, Gadi I, Kontgen F, Abbondanzo SJ: Blastocyst implantation depends on maternal expression of leukaemia inhibitory factor. *Nature* 1992, 359(6390):76-79.

Stice SL, Strelchenko NS, Keefer CL, Matthews L: Pluripotent bovine embryonic cell lines direct embryonic development following nuclear transfer. *Biol Reprod* 1996, 54(1):100-110.

Stojkovic M, Stojkovic P, Leary C, Hall VJ, Armstrong L, Herbert M, Nesbitt M, Lako M, Murdoch A: Derivation of a human blastocyst after heterologous nuclear transfer to donated oocytes. *Reproductive biomedicine online* 2005, 11(2):226-231.

Strachan T, Read AP. In: *Human molecular genetics*. 2nd edn. New York; 1999.

Strickland S, Mahdavi V: The induction of differentiation in teratocarcinoma stem cells by retinoic acid. *Cell* 1978, 15(2):393-403.

Su Q, Prosser HM, Campos LS, Ortiz M, Nakamura T, Warren M, Dupuy AJ, Jenkins NA, Copeland NG, Bradley A et al: A DNA transposon-based approach to validate oncogenic mutations in the mouse. *P Natl Acad Sci USA* 2008, 105(50):19904-19909.

Subbarayan V, Kastner P, Mark M, Dierich A, Gorry P, Chambon P: Limited specificity and large overlap of the functions of the mouse RAR gamma 1 and RAR gamma 2 isoforms. *Mechanisms of development* 1997, 66(1-2):131-142.

Subramanyam D, Lamouille S, Judson RL, Liu JY, Bucay N, Derynck R, Blelloch R: Multiple targets of miR-302 and miR-372 promote reprogramming of human fibroblasts to induced pluripotent stem cells. *Nat Biotechnol* 2011, 29(5):443-448.

Sukoyan MA, Vatolin SY, Golubitsa AN, Zhelezova AI, Semenova LA, Serov OL: Embryonic stem cells derived from morulae, inner cell mass, and blastocysts of mink: comparisons of their pluripotencies. *Molecular reproduction and development* 1993, 36(2):148-158.

Sundqvist J, Falconer H, Seddighzadeh M, Vodolazkaia A, Fassbender A, Kyama C, Bokor A, Stephansson O, Gemzell-Danielsson K, D'Hooghe TM: Ovarian cancer-associated polymorphisms in the BNC2 gene among women with endometriosis. *Hum Reprod* 2011, 26(8):2253-2257.

Suzuki T, Minehata K, Akagi K, Jenkins NA, Copeland NG: Tumor suppressor gene identification using retroviral insertional mutagenesis in Blm-deficient mice. *The EMBO journal* 2006, 25(14):3422-3431.

Sylvester I, Scholer HR: Regulation of the Oct-4 gene by nuclear receptors. *Nucleic Acids Res* 1994, 22(6):901-911.

Szabo E, Rampalli S, Risueno RM, Schnerch A, Mitchell R, Fiebig-Comyn A, Levadoux-Martin M, Bhatia M: Direct conversion of human fibroblasts to multilineage blood progenitors. *Nature* 2010, 468(7323):521-526.

Szymczak AL, Workman CJ, Wang Y, Vignali KM, Dilioglou S, Vanin EF, Vignali DAA: Correction of multi-gene deficiency in vivo using a single 'self-cleaving' 2A peptide-based retroviral vector (vol 22, pg 589, 2004). *Nat Biotechnol* 2004, 22(12):1590-1590.

Tada M, Takahama Y, Abe K, Nakatsuji N, Tada T: Nuclear reprogramming of somatic cells by in vitro hybridization with ES cells. *Curr Biol* 2001, 11(19):1553-1558.

Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S: Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007, 131(5):861-872.

Takahashi K, Yamanaka S: Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006, 126(4):663-676.

Takeda J, Seino S, Bell GI: Human Oct3 Gene Family - Cdna Sequences, Alternative Splicing, Gene Organization, Chromosomal Location, and Expression at Low-Levels in Adult Tissues. *Nucleic Acids Res* 1992, 20(17):4613-4620.

Tanaka J, Watanabe T, Nakamura N, Sobue K: Morphological and Biochemical Analyses of Contractile Proteins (Actin, Myosin, Caldesmon and Tropomyosin) in Normal and Transformed-Cells. *J Cell Sci* 1993, 104:595-606.

Taranger CK, Noer A, Sorensen AL, Hakelien AM, Boquest AC, Collas P: Induction of dedifferentiation, genomewide transcriptional programming, and epigenetic reprogramming by extracts of carcinoma and embryonic stem cells. *Molecular biology of the cell* 2005, 16(12):5719-5735.

Teranishi T, Tanaka M, Kimoto S, Ono Y, Miyakoshi K, Kono T, Yoshimura Y: Rapid replacement of somatic linker histones with the oocyte-specific linker histone H1foo in nuclear transfer. *Developmental biology* 2004, 266(1):76-86.

Tesar PJ, Chenoweth JG, Brook FA, Davies TJ, Evans EP, Mack DL, Gardner RL, McKay RD: New cell lines from mouse epiblast share defining features with human embryonic stem cells. *Nature* 2007, 448(7150):196-199.

Thaller C, Eichele G: Identification and spatial distribution of retinoids in the developing chick limb bud. *Nature* 1987, 327(6123):625-628.

Thomas KR, Capecchi MR: Site-directed mutagenesis by gene targeting in mouse embryo-derived stem cells. *Cell* 1987, 51(3):503-512.

Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, Jones JM: Embryonic stem cell lines derived from human blastocysts. *Science* 1998, 282(5391):1145-1147.

Thomson JA, Kalishman J, Golos TG, Durning M, Harris CP, Becker RA, Hearn JP: Isolation of a primate embryonic stem cell line. *P Natl Acad Sci USA* 1995, 92(17):7844-7848.

Thomson JA, Kalishman J, Golos TG, Durning M, Harris CP, Hearn JP: Pluripotent cell lines derived from common marmoset (*Callithrix jacchus*) blastocysts. *Biol Reprod* 1996, 55(2):254-259.

Thorel F, Nepote V, Avril I, Kohno K, Desgraz R, Chera S, Herrera PL: Conversion of adult pancreatic alpha-cells to beta-cells after extreme beta-cell loss. *Nature* 2010, 464(7292):1149-1154.

Tiemann U, Sgodda M, Warlich E, Ballmaier M, Scholer HR, Schambach A, Cantz T: Optimal Reprogramming Factor Stoichiometry Increases Colony Numbers and Affects Molecular Characteristics of Murine Induced Pluripotent Stem Cells. *Cytom Part A* 2011, 79A(6):426-435.

Tomioka M, Nishimoto M, Miyagi S, Katayanagi T, Fukui N, Niwa H, Muramatsu M, Okuda A: Identification of Sox-2 regulatory region which is under the control of Oct-3/4-Sox-2 complex. *Nucleic Acids Res* 2002, 30(14):3202-3213.

Tontonoz P, Hu E, Spiegelman BM: Stimulation of adipogenesis in fibroblasts by PPAR gamma 2, a lipid-activated transcription factor. *Cell* 1994, 79(7):1147-1156.

Tsai SY, Clavel C, Kim S, Ang YS, Grisanti L, Lee DF, Kelley K, Rendl M: Oct4 and klf4 reprogram dermal papilla cells into induced pluripotent stem cells. *Stem Cells* 2010, 28(2):221-228.

Tseng H, Green H: Basonuclin: a keratinocyte protein with multiple paired zinc fingers. *P Natl Acad Sci USA* 1992, 89(21):10311-10315.

Tsonis PA, Del Rio-Tsonis K: Lens and retina regeneration: transdifferentiation, stem cells and clinical applications. *Experimental eye research* 2004, 78(2):161-172.

Tsubota S, Ashburner M, Schedl P: P-Element-Induced Control Mutations at the R Gene of *Drosophila-Melanogaster*. *Mol Cell Biol* 1985, 5(10):2567-2574.

Tsubota S, Schedl P: Hybrid Dysgenesis-Induced Revertants of Insertions at the 5' End of the Rudimentary Gene in *Drosophila-Melanogaster* - Transposon-Induced Control Mutations. *Genetics* 1986, 114(1):165-182.

Urbach A, Bar-Nur O, Daley GQ, Benvenisty N: Differential modeling of fragile X syndrome by human embryonic stem cells and induced pluripotent stem cells. *Cell stem cell* 2010, 6(5):407-411.

Utikal J, Polo JM, Stadtfeld M, Maherali N, Kulalert W, Walsh RM, Khalil A, Rheinwald JG, Hochedlinger K: Immortalization eliminates a roadblock during cellular reprogramming into iPS cells. *Nature* 2009, 460(7259):1145-1148.

Vallier L, Alexander M, Pedersen RA: Activin/Nodal and FGF pathways cooperate to maintain pluripotency of human embryonic stem cells. *J Cell Sci* 2005, 118(Pt 19):4495-4509.

van den Berg DL, Snoek T, Mullin NP, Yates A, Bezstarosti K, Demmers J, Chambers I, Poot RA: An Oct4-centered protein interaction network in embryonic stem cells. *Cell stem cell* 2010, 6(4):369-381.

van der Wees J, Schilthuis JG, Koster CH, Diesveld-Schipper H, Folkers GE, van der Saag PT, Dawson MI, Shudo K, van der Burg B, Durston AJ: Inhibition of retinoic acid receptor-mediated signalling alters positional identity in the developing hindbrain. *Development* 1998, 125(3):545-556.

van Eijk MJT, van Rooijen MA, Modina S, Scesi L, Folkers G, van Tol HTA, Bevers MM, Fisher SR, Lewin HA, Rakacolli D et al: Molecular cloning, genetic mapping, and developmental expression of bovine POU5F1. *Biol Reprod* 1999, 60(5):1093-1103.

VandenDriessche T, Ivics Z, Izsvak Z, Chuah MK: Emerging potential of transposons for gene therapy and generation of induced pluripotent stem cells. *Blood* 2009, 114(8):1461-1468.

Vander Heiden MG, Cantley LC, Thompson CB: Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 2009, 324(5930):1029-1033.

Vanhoutteghem A, Bouche C, Maciejewski-Duval A, Herve F, Djian P: Basonuclins and disco: Orthologous zinc finger proteins essential for development in vertebrates and arthropods. *Biochimie* 2011, 93(2):127-133.

Vanhoutteghem A, Djian P: Basonuclin 2: an extremely conserved homolog of the zinc finger protein basonuclin. P Natl Acad Sci USA 2004, 101(10):3468-3473.

Vanhoutteghem A, Djian P: Basonuclins 1 and 2, whose genes share a common origin, are proteins with widely different properties and functions. P Natl Acad Sci USA 2006, 103(33):12423-12428.

Vanhoutteghem A, Djian P: The human basonuclin 2 gene has the potential to generate nearly 90,000 mRNA isoforms encoding over 2000 different proteins. Genomics 2007, 89(1):44-58.

Vanhoutteghem A, Maciejewski-Duval A, Bouche C, Delhomme B, Herve F, Daubigney F, Soubigou G, Araki M, Araki K, Yamamura K et al: Basonuclin 2 has a function in the multiplication of embryonic craniofacial mesenchymal cells and is orthologous to disco proteins. P Natl Acad Sci USA 2009, 106(34):14432-14437.

Varum S, Rodrigues AS, Moura MB, Momcilovic O, Easley CA, Ramalho-Santos J, Van Houten B, Schatten G: Energy Metabolism in Human Pluripotent Stem Cells and Their Differentiated Counterparts. PloS one 2011, 6(6).

Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Sudhof TC, Wernig M: Direct conversion of fibroblasts to functional neurons by defined factors. Nature 2010, 463(7284):1035-1041.

Vivian JL, Chen Y, Yee D, Schneider E, Magnuson T: An allelic series of mutations in Smad2 and Smad4 identified in a genotype-based screen of N-ethyl-N-nitrosourea-mutagenized mouse embryonic stem cells. P Natl Acad Sci USA 2002, 99(24):15542-15547.

von Baer KE: *De ovi mammalium et hominis genesi: Sumptibus L. Vossii*; 1827.

von Baer KE: *Über Entwicklungsgeschichte der Thier. Beobachtung und Reflexion: Königsberg, Bei den Gebrüder Borntäger*; 1828.

Wang G, Zhang H, Zhao Y, Li J, Cai J, Wang P, Meng S, Feng J, Miao C, Ding M et al: Noggin and bFGF cooperate to maintain the pluripotency of human embryonic stem cells in the absence of feeder layers. *Biochem Biophys Res Commun* 2005, 330(3):934-942.

Wang J, Rao S, Chu J, Shen X, Levasseur DN, Theunissen TW, Orkin SH: A protein interaction network for pluripotency of embryonic stem cells. *Nature* 2006, 444(7117):364-368.

Wang T, Chen K, Zeng X, Yang J, Wu Y, Shi X, Qin B, Zeng L, Esteban MA, Pan G et al: The histone demethylases *jhdmla/1b* enhance somatic cell reprogramming in a vitamin-C-dependent manner. *Cell Stem Cell* 2011, 9(6):575-587.

Wang W, Bradley A, Huang Y: A piggyBac transposon-based genome-wide library of insertionally mutated *Blm*-deficient murine ES cells. *Genome Research* 2009, 19(4):667-673.

Wang W, Hale C, Goulding D, Haslam SM, Tissot B, Lindsay C, Michell S, Titball R, Yu J, Toribio AL et al: Mannosidase 2, alpha 1 deficiency is associated with ricin resistance in embryonic stem (ES) cells. *PloS one* 2011, 6(8):e22993.

Wang W, Lin C, Lu D, Ning Z, Cox T, Melvin D, Wang X, Bradley A, Liu P: Chromosomal transposition of PiggyBac in mouse embryonic stem cells. *PNAS* 2008, 105(27):9290-9295.

Wang W, Yang J, Liu H, Lu D, Chen X, Zenonos Z, Campos LS, Rad R, Guo G, Zhang S et al: Rapid and efficient reprogramming of somatic cells to induced pluripotent stem cells by retinoic acid receptor gamma and liver receptor homolog 1. *PNAS* 2011, 108(45):18283-18288.

Wang YJ, Kumar N, Nuhant P, Cameron MD, Istrate MA, Roush WR, Griffin PR, Burris TP: Identification of SR1078, a Synthetic Agonist for the Orphan Nuclear Receptors ROR alpha and ROR gamma. *ACS Chem Biol* 2010, 5(11):1029-1034.

Warkany J, Schraffenberger E: Congenital malformations induced in rats by maternal vitamin A deficiency; defects of the eye. *Archives of ophthalmology* 1946, 35:150-169.

Warren L, Manos PD, Ahfeldt T, Loh YH, Li H, Lau F, Ebina W, Mandal PK, Smith ZD, Meissner A et al: Highly efficient reprogramming to pluripotency and directed differentiation of human cells with synthetic modified mRNA. *Cell stem cell* 2010, 7(5):618-630.

Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, An P et al: Initial sequencing and comparative analysis of the mouse genome. *Nature* 2002, 420(6915):520-562.

Watson JD, Crick FH: Molecular structure of nucleic acids; a structure for deoxyribose nucleic acid. *Nature* 1953, 171(4356):737-738.

Wei F, Scholer HR, Atchison ML: Sumoylation of Oct4 enhances its stability, DNA binding, and transactivation. *The Journal of biological chemistry* 2007, 282(29):21551-21560.

Weintraub H, Tapscott SJ, Davis RL, Thayer MJ, Adam MA, Lassar AB, Miller AD: Activation of muscle-specific genes in pigment, nerve, fat, liver, and fibroblast cell lines by forced expression of MyoD. *P Natl Acad Sci USA* 1989, 86(14):5434-5438.

Weismann A, Parker WN, Rãnnfeldt H: *The germ-plasm: a theory of heredity*: Scribner's; 1893.

Welstead GG, Creighton MP, Bilodeau S, Cheng AW, Markoulaki S, Young RA, Jaenisch R: X-linked H3K27me3 demethylase Utx is required for embryonic development in a sex-specific manner. *P Natl Acad Sci USA* 2012, 109(32):13004-13009.

White JC, Shankar VN, Highland M, Epstein ML, DeLuca HF, Clagett-Dame M: Defects in embryonic hindbrain development and fetal resorption resulting from vitamin A deficiency in the rat are prevented by feeding pharmacological levels of all-trans-retinoic acid. *P Natl Acad Sci USA* 1998, 95(23):13459-13464.

Whyte WA, Bilodeau S, Orlando DA, Hoke HA, Frampton GM, Foster CT, Cowley SM, Young RA: Enhancer decommissioning by LSD1 during embryonic stem cell differentiation. *Nature* 2012, 482(7384):221-225.

Widschwendter M, Berger J, Daxenbichler G, Muller-Holzner E, Widschwendter A, Mayr A, Marth C, Zeimet AG: Loss of retinoic acid receptor beta expression in breast cancer and morphologically normal adjacent tissue but not in the normal breast tissue distant from the cancer. *Cancer research* 1997, 57(19):4158-4161.

Willadsen SM: Nuclear transplantation in sheep embryos. *Nature* 1986, 320(6057):63-65.

Williams RL, Hilton DJ, Pease S, Willson TA, Stewart CL, Gearing DP, Wagner EF, Metcalf D, Nicola NA, Gough NM: Myeloid leukaemia inhibitory factor maintains the developmental potential of embryonic stem cells. *Nature* 1988, 336(6200):684-687.

Wilmot I, Schnieke AE, McWhir J, Kind AJ, Campbell KH: Viable offspring derived from fetal and adult mammalian cells. *Nature* 1997, 385(6619):810-813.

Wilson JG, Roth CB, Warkany J: An analysis of the syndrome of malformations induced by maternal vitamin A deficiency. Effects of restoration of vitamin A at various times during gestation. *The American journal of anatomy* 1953, 92(2):189-217.

Wilson JG, Warkany J: Malformations in the genito-urinary tract induced by maternal vitamin A deficiency in the rat. *The American journal of anatomy* 1948, 83(3):357-407.

Wilson MH, Coates CJ, George AL: PiggyBac transposon-mediated gene transfer in human cells. *Mol Ther* 2007, 15(1):139-145.

Wilson PA, Hemmati-Brivanlou A: Induction of epidermis and inhibition of neural fate by Bmp-4. *Nature* 1995, 376(6538):331-333.

Witte ON, Rosenberg N, Paskind M, Shields A, Baltimore D: Identification of an Abelson murine leukemia virus-encoded protein present in transformed fibroblast and lymphoid cells. *P Natl Acad Sci USA* 1978, 75(5):2488-2492.

Wolff CF: *Theoria generationis: Typis et sumtu Io . Christ. Hendel; 1774.*

Woltjen K, Michael IP, Mohseni P, Desai R, Mileikovsky M, Hamalainen R, Cowling R, Wang W, Liu P, Gertsenstein M et al: piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. *Nature* 2009, 458(7239):766-770.

Wu H, Zhang Y: Mechanisms and functions of Tet protein-mediated 5-methylcytosine oxidation. *Gene Dev* 2011, 25(23):2436-2452.

Wu SCY, Meir YJJ, Coates CJ, Handler AM, Pelczar P, Moisyadi S, Kaminski JM: piggyBac is a flexible and highly active transposon as compared to Sleeping Beauty, Tol2 and Mos1 in mammalian cells. *P Natl Acad Sci USA* 2006, 103(41):15008-15013.

Xie H, Ye M, Feng R, Graf T: Stepwise reprogramming of B cells into macrophages. *Cell* 2004, 117(5):663-676.

Xiong G, Wang C, Evers BM, Zhou BP, Xu R: RORalpha Suppresses Breast Tumor Invasion by Inducing SEMA3F Expression. *Cancer research* 2012, 72(7):1728-1739.

Xu C, Rosler E, Jiang J, Lebkowski JS, Gold JD, O'Sullivan C, Delavan-Boorsma K, Mok M, Bronstein A, Carpenter MK: Basic fibroblast growth factor supports undifferentiated human embryonic stem cell growth without conditioned medium. *Stem Cells* 2005, 23(3):315-323.

Xu H, Wang W, Li C, Yu H, Yang A, Wang B, Jin Y: WWP2 promotes degradation of transcription factor OCT4 in human embryonic stem cells. *Cell research* 2009, 19(5):561-573.

Xu RH, Chen X, Li DS, Li R, Addicks GC, Glennon C, Zwaka TP, Thomson JA: BMP4 initiates human embryonic stem cell differentiation to trophoblast. *Nat Biotechnol* 2002, 20(12):1261-1264.

Xu XC, Liu X, Tahara E, Lippman SM, Lotan R: Expression and up-regulation of retinoic acid receptor-beta is associated with retinoid sensitivity and colony formation in esophageal cancer cell lines. *Cancer research* 1999, 59(10):2477-2483.

Xu XC, Sneige N, Liu X, Nandagiri R, Lee JJ, Lukmanji F, Hortobagyi G, Lippman SM, Dhingra K, Lotan R: Progressive decrease in nuclear retinoic acid receptor beta messenger RNA level during breast carcinogenesis. *Cancer research* 1997, 57(22):4992-4996.

Yamaguchi S, Kimura H, Tada M, Nakatsuji N, Tada T: Nanog expression in mouse germ cell development. *Gene expression patterns : GEP* 2005, 5(5):639-646.

Yamashita A, Takada T, Narita J, Yamamoto G, Torii R: Osteoblastic differentiation of monkey embryonic stem cells in vitro. *Cloning and stem cells* 2005, 7(4):232-237.

Yang H, Shi L, Wang BA, Liang D, Zhong C, Liu W, Nie Y, Liu J, Zhao J, Gao X et al: Generation of genetically modified mice by oocyte injection of androgenetic haploid embryonic stem cells. *Cell* 2012, 149(3):605-617.

Yant SR, Wu XL, Huang Y, Garrison B, Burgess SM, Kay MA: High-resolution genome-wide mapping of transposon integration in mammals. *Mol Cell Biol* 2005, 25(6):2085-2094.

Yao JY, Zhang L, Zhang X, He ZY, Ma Y, Hui LJ, Wang X, Hu YP: H3K27 Trimethylation Is an Early Epigenetic Event of p16(INK4a) Silencing for Regaining Tumorigenesis in Fusion Reprogrammed Hepatoma Cells. *J Biol Chem* 2010, 285(24):18828-18837.

Yeo S, Jeong S, Kim J, Han JS, Han YM, Kang YK: Characterization of DNA methylation change in stem cell marker genes during differentiation of human embryonic stem cells. *Biochem Bioph Res Co* 2007, 359(3):536-542.

Yeom YI, Fuhrmann G, Ovitt CE, Brehm A, Ohbo K, Gross M, Hubner K, Scholer HR: Germline regulatory element of Oct-4 specific for the totipotent cycle of embryonal cells. *Development* 1996, 122(3):881-894.

Yeom YI, Ha HS, Balling R, Scholer HR, Artzt K: Structure, expression and chromosomal location of the Oct-4 gene. *Mechanisms of development* 1991, 35(3):171-179.

Ying QL, Nichols J, Chambers I, Smith A: BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 2003, 115(3):281-292.

Ying QL, Nichols J, Evans EP, Smith AG: Changing potency by spontaneous fusion. *Nature* 2002, 416(6880):545-548.

Ying QL, Wray J, Nichols J, Battle-Morera L, Doble B, Woodgett J, Cohen P, Smith A: The ground state of embryonic stem cell self-renewal. *Nature* 2008, 453(7194):519-523.

Yoshida K, Chambers I, Nichols J, Smith A, Saito M, Yasukawa K, Shoyab M, Taga T, Kishimoto T: Maintenance of the pluripotential phenotype of embryonic stem cells through direct activation of gp130 signalling pathways. *Mechanisms of development* 1994, 45(2):163-171.

Yoshida Y, Takahashi K, Okita K, Ichisaka T, Yamanaka S: Hypoxia enhances the generation of induced pluripotent stem cells. *Cell stem cell* 2009, 5(3):237-241.

You JS, Kelly TK, De Carvalho DD, Taberlay PC, Liang GN, Jones PA: OCT4 establishes and maintains nucleosome-depleted regions that provide additional layers of epigenetic regulation of its target genes. *P Natl Acad Sci USA* 2011, 108(35):14497-14502.

You L, Chang D, Du HZ, Zhao YP: Genome-wide screen identifies PVT1 as a regulator of Gemcitabine sensitivity in human pancreatic cancer cells. *Biochem Bioph Res Co* 2011, 407(1):1-6.

Yu J, Hu K, Smuga-Otto K, Tian S, Stewart R, Slukvin, II, Thomson JA: Human induced pluripotent stem cells free of vector and transgene sequences. *Science* 2009, 324(5928):797-801.

Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, Nie J, Jonsdottir GA, Ruotti V, Stewart R et al: Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007, 318(5858):1917-1920.

Yu VC, Delsert C, Andersen B, Holloway JM, Devary OV, Naar AM, Kim SY, Boutin JM, Glass CK, Rosenfeld MG: RXR beta: a coregulator that enhances binding of retinoic acid, thyroid hormone, and vitamin D receptors to their cognate response elements. *Cell* 1991, 67(6):1251-1266.

Yusa K, Horie K, Kondoh G, Kouno M, Maeda Y, Kinoshita T, Takeda J: Genome-wide phenotype analysis in ES cells by regulated disruption of Bloom's syndrome gene. *Nature* 2004, 429(6994):896-899.

Yusa K, Rad R, Takeda J, Bradley A: Generation of transgene-free induced pluripotent mouse stem cells by the piggyBac transposon. *Nature methods* 2009, 6(5):363-369.

Yusa K, Rashid ST, Strick-Marchand H, Varela I, Liu PQ, Paschon DE, Miranda E, Ordonez A, Hannan NR, Rouhani FJ et al: Targeted gene correction of alpha1-antitrypsin deficiency in induced pluripotent stem cells. *Nature* 2011, 478(7369):391-394.

Zambrowicz BP, Friedrich GA, Buxton EC, Lilleberg SL, Person C, Sands AT: Disruption and sequence identification of 2,000 genes in mouse embryonic stem cells. *Nature* 1998, 392(6676):608-611.

Zambrowicz BP, Imamoto A, Fiering S, Herzenberg LA, Kerr WG, Soriano P: Disruption of overlapping transcripts in the ROSA beta geo 26 gene trap strain leads to widespread expression of beta-galactosidase in mouse embryos and hematopoietic cells. *P Natl Acad Sci USA* 1997, 94(8):3789-3794.

Zelent A, Krust A, Petkovich M, Kastner P, Chambon P: Cloning of murine alpha and beta retinoic acid receptors and a novel receptor gamma predominantly expressed in skin. *Nature* 1989, 339(6227):714-717.

Zelent A, Mendelsohn C, Kastner P, Krust A, Garnier JM, Ruffenach F, Leroy P, Chambon P: Differentially expressed isoforms of the mouse retinoic acid receptor beta generated by usage of two promoters and alternative splicing. *The EMBO journal* 1991, 10(1):71-81.

Zhang J, Lian Q, Zhu G, Zhou F, Sui L, Tan C, Mutalif RA, Navasankari R, Zhang Y, Tse HF et al: A human iPSC model of Hutchinson Gilford Progeria reveals vascular smooth muscle and mesenchymal stem cell defects. *Cell stem cell* 2011, 8(1):31-45.

Zhang P, Andrianakos R, Yang Y, Liu C, Lu W: Kruppel-like factor 4 (Klf4) prevents embryonic stem (ES) cell differentiation by regulating Nanog gene expression. *The Journal of biological chemistry* 2010, 285(12):9180-9189.

Zhang XK, Hoffmann B, Tran PB, Graupner G, Pfahl M: Retinoid X receptor is an auxiliary protein for thyroid hormone and retinoic acid receptors. *Nature* 1992, 355(6359):441-446.

Zhang Z, Liao B, Xu M, Jin Y: Post-translational modification of POU domain transcription factor Oct-4 by SUMO-1. *Faseb J* 2007, 21(12):3042-3051.

Zhao T, Zhang ZN, Rong Z, Xu Y: Immunogenicity of induced pluripotent stem cells. *Nature* 2011, 474(7350):212-215.

Zhao Y, Yin X, Qin H, Zhu F, Liu H, Yang W, Zhang Q, Xiang C, Hou P, Song Z et al: Two supporting factors greatly improve the efficiency of human iPSC generation. *Cell stem cell* 2008, 3(5):475-479.

Zheng PP, Severijnen LA, van der Weiden M, Willemsen R, Kros JM: A crucial role of caldesmon in vascular development in vivo. *Cardiovasc Res* 2009, 81(2):362-369.

Zheng PP, Severijnen LA, Willemsen R, Kros JM: Caldesmon is essential for cardiac morphogenesis and function: In vivo study using a zebrafish model. *Biochem Biophys Res Commun* 2009, 378(1):37-40.

Zhou H, Wu S, Joo JY, Zhu S, Han DW, Lin T, Trauger S, Bien G, Yao S, Zhu Y et al: Generation of induced pluripotent stem cells using recombinant proteins. *Cell stem cell* 2009, 4(5):381-384.

Zhou Q, Brown J, Kanarek A, Rajagopal J, Melton DA: In vivo reprogramming of adult pancreatic exocrine cells to beta-cells. *Nature* 2008, 455(7213):627-632.