

References

- Aanstad, P. and Whitaker, M.** (1999). Predictability of dorso-ventral asymmetry in the cleavage stage zebrafish embryo: an analysis using lithium sensitivity as a dorso-ventral marker. *Mech Dev* **88**, 33-41.
- Aligianis, I. A., Johnson, C. A., Gissen, P., Chen, D., Hampshire, D., Hoffmann, K., Maina, E. N., Morgan, N. V., Tee, L., Morton, J. et al.** (2005). Mutations of the catalytic subunit of RAB3GAP cause Warburg Micro syndrome. *Nat Genet* **37**, 221-3.
- Aligianis, I. A., Morgan, N. V., Mione, M., Johnson, C. A., Rosser, E., Hennekam, R. C., Adams, G., Trembath, R. C., Pilz, D. T., Stoodley, N. et al.** (2006). Mutation in Rab3 GTPase-activating protein (RAB3GAP) noncatalytic subunit in a kindred with Martsolf syndrome. *Am J Hum Genet* **78**, 702-7.
- Allan, B. B., Moyer, B. D. and Balch, W. E.** (2000). Rab1 recruitment of p115 into a cis-SNARE complex: programming budding COPII vesicles for fusion. *Science* **289**, 444-8.
- Alvarez, C., Garcia-Mata, R., Brandon, E. and Sztul, E.** (2003). COPI recruitment is modulated by a Rab1b-dependent mechanism. *Mol Biol Cell* **14**, 2116-27.
- Amatruda, J. F. and Zon, L. I.** (1999). Dissecting hematopoiesis and disease using the zebrafish. *Dev Biol* **216**, 1-15.
- Amaya, E., Stein, P. A., Musci, T. J. and Kirschner, M. W.** (1993). FGF signalling in the early specification of mesoderm in Xenopus. *Development* **118**, 477-87.
- Anant, J. S., Desnoyers, L., Machius, M., Demeler, B., Hansen, J. C., Westover, K. D., Deisenhofer, J. and Seabra, M. C.** (1998). Mechanism of Rab geranylgeranylation: formation of the catalytic ternary complex. *Biochemistry* **37**, 12559-68.
- Anders, R. A., Dore, J. J., Jr., Arline, S. L., Garamszegi, N. and Leof, E. B.** (1998). Differential requirement for type I and type II transforming growth factor beta receptor kinase activity in ligand-mediated receptor endocytosis. *J Biol Chem* **273**, 23118-25.
- Armstrong, J.** (2000). How do Rab proteins function in membrane traffic? *The International Journal of Biochemistry and Cell Biology* **32**, 303-307.
- Ayala, J., Olofsson, B., Touchot, N., Zahraoui, A., Tavitian, A. and Prochiantz, A.** (1989). Developmental and regional expression of three new members of the ras-gene family in the mouse brain. *J Neurosci Res* **22**, 384-9.
- Babb, S. G. and Marrs, J. A.** (2004). E-cadherin regulates cell movements and tissue formation in early zebrafish embryos. *Dev Dyn* **230**, 263-77.
- Bafico, A., Liu, G., Yaniv, A., Gazit, A. and Aaronson, S. A.** (2001). Novel mechanism of Wnt signalling inhibition mediated by Dickkopf-1 interaction with LRP6/Arrow. *Nat Cell Biol* **3**, 683-6.
- Bahadoran, P., Aberdam, E., Mantoux, F., Busca, R., Bille, K., Yalman, N., de Saint-Basile, G., Casaroli-Marano, R., Ortonne, J. P. and Ballotti, R.** (2001). Rab27a: A key to melanosome transport in human melanocytes. *J Cell Biol* **152**, 843-50.
- Batista, A., Millan, J., Mittelbrunn, M., Sanchez-Madrid, F. and Alonso, M. A.** (2004). Recruitment of transferrin receptor to immunological synapse in response to TCR engagement. *J Immunol* **172**, 6709-14.

- Bauer, H., Lele, Z., Rauch, G. J., Geisler, R. and Hammerschmidt, M.** (2001). The type I serine/threonine kinase receptor Alk8/Lost-a-fin is required for Bmp2b/7 signal transduction during dorsoventral patterning of the zebrafish embryo. *Development* **128**, 849-58.
- Beard, M., Satoh, A., Shorter, J., and Warren, G.** (2005). A Cryptic Rab1-binding Site in the p115 Tethering Protein. *J. Biol. Chem.* **280**: 25840-25848.
- Behra, M., Cousin, X., Bertrand, C., Vonesch, J. L., Biellmann, D., Chatonnet, A. and Strahle, U.** (2002). Acetylcholinesterase is required for neuronal and muscular development in the zebrafish embryo. *Nat Neurosci* **5**, 111-8.
- Bejsovec, A. and Wieschaus, E.** (1995). Signaling activities of the Drosophila wingless gene are separately mutable and appear to be transduced at the cell surface. *Genetics* **139**, 309-20.
- Belenkaya, T. Y., Han, C., Yan, D., Opoka, R. J., Khodoun, M., Liu, H. and Lin, X.** (2004). Drosophila Dpp morphogen movement is independent of dynamin-mediated endocytosis but regulated by the glypcan members of heparan sulfate proteoglycans. *Cell* **119**, 231-44.
- Bellipanni, G., Varga, M., Maegawa, S., Imai, Y., Kelly, C., Myers, A. P., Chu, F., Talbot, W. S. and Weinberg, E. S.** (2006). Essential and opposing roles of zebrafish beta-catenins in the formation of dorsal axial structures and neurectoderm. *Development* **133**, 1299-309.
- Bellot, F., Crumley, G., Kaplow, J. M., Schlessinger, J., Jaye, M. and Dionne, C. A.** (1991). Ligand-induced transphosphorylation between different FGF receptors. *Embo J* **10**, 2849-54.
- Benli, M., Doring, F., Robinson, D. G., Yang, X. and Gallwitz, D.** (1996). Two GTPase isoforms, Ypt31p and Ypt32p, are essential for Golgi function in yeast. *Embo J* **15**, 6460-75.
- Bennett, M. K., Calakos, N. and Scheller, R. H.** (1992). Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* **257**, 255-9.
- Bernards, A. and Settleman, J.** (2004). GAP control: regulating the regulators of small GTPases. *Trends Cell Biol* **14**, 377-85.
- Betchaku, T. and Trinkaus, J. P.** (1986). Programmed endocytosis during epiboly of Fundulus heteroclitus. *Am. Zool* **26**, 193-199.
- Bischof, J. and Driever, W.** (2004). Regulation of hhex expression in the yolk syncytial layer, the potential Nieuwkoop center homolog in zebrafish. *Dev Biol* **276**, 552-62.
- Bishop, A. L. and Hall, A.** (2000). Rho GTPases and their effector proteins. *Biochem J* **348 Pt 2**, 241-55.
- Bjornson, C. R., Griffin, K. J., Farr, G. H., 3rd, Terashima, A., Himeda, C., Kikuchi, Y. and Kimelman, D.** (2005). Eomesodermin is a localized maternal determinant required for endoderm induction in zebrafish. *Dev Cell* **9**, 523-33.
- Blader, P., Rastegar, S., Fischer, N. and Strahle, U.** (1997). Cleavage of the BMP-4 antagonist chordin by zebrafish tolloid. *Science* **278**, 1937-40.
- Blitzer, J. T. and Nusse, R.** (2006). A critical role for endocytosis in Wnt signaling. *BMC Cell Biol* **7**, 28.
- Bronner-Fraser, M., Wolf, J. J. and Murray, B. A.** (1992). Effects of antibodies against N-cadherin and N-CAM on the cranial neural crest and neural tube. *Dev Biol* **153**, 291-301.
- Brott, B. K. and Sokol, S. Y.** (2002). Regulation of Wnt/LRP signaling by distinct domains of Dickkopf proteins. *Mol Cell Biol* **22**, 6100-10.

- Bruce, A. E., Howley, C., Zhou, Y., Vickers, S. L., Silver, L. M., King, M. L. and Ho, R. K.** (2003). The maternally expressed zebrafish T-box gene eomesodermin regulates organizer formation. *Development* **130**, 5503-17.
- Brymora, A., Valova, V. A., Larsen, M. R., Roufogalis, B. D. and Robinson, P. J.** (2001). The brain exocyst complex interacts with RalA in a GTP-dependent manner: identification of a novel mammalian Sec3 gene and a second Sec15 gene. *J Biol Chem* **276**, 29792-7.
- Bucci, C., Lutcke, A., Steele-Mortimer, O., Olkkonen, V., Dupree, P., Chiariello, M., Bruni, C., Simons, K. and Zerial, M.** (1995). Co-operative regulation of endocytosis by three RAB5 isoforms. *FEBS Letters* **366**, 65-71.
- Bucci, C., Parton, R. G., Mather, I. H., Stunnenberg, H., Simons, K., Hoflack, B. and Zerial, M.** (1992). The small GTPase rab5 functions as a regulatory factor in the early endocytic pathway. *Cell* **70**, 715-28.
- Bush, J., Franek, K., Daniel, J., Spiegelman, G. B., Weeks, G. and Cardelli, J.** (1993). Cloning and characterization of five novel Dictyostelium discoideum rab-related genes. *Gene* **136**, 55-60.
- Buznikov, G. A. and Ignat'eva, G. M.** (1958). [Hatching enzymes.]. *Usp Sovrem Biol* **46**, 337-56.
- Campos, I.** (2004). Genes Controlling Development: Roles for Rabs and Glis. In *Anatomy*, (ed., pp. 209. London: National Institute for Medical Research and University Collage London.
- Camus, C., Hermann-Le Denmat, S. and Jacquet, M.** (1995). Identification of guanine exchange factor key residues involved in exchange activity and Ras interaction. *Oncogene* **11**, 951-9.
- Cao, X. and Barlowe, C.** (2000). Asymmetric requirements for a Rab GTPase and SNARE proteins in fusion of COPII vesicles with acceptor membranes. *J Cell Biol* **149**, 55-66.
- Casey, P. J. and Seabra, M. C.** (1996). Protein prenyltransferases. *J Biol Chem* **271**, 5289-92.
- Chandrasekhar, A., Warren, J. T., Jr., Takahashi, K., Schauerte, H. E., van Eeden, F. J., Haffter, P. and Kuwada, J. Y.** (1998). Role of sonic hedgehog in branchiomotor neuron induction in zebrafish. *Mech Dev* **76**, 101-15.
- Chavrier, P., Parton, R. G., Hauri, H. P., Simons, K. and Zerial, M.** (1990). Localization of low molecular weight GTP binding proteins to exocytic and endocytic compartments. *Cell* **62**, 317-29.
- Chavrier, P., Simons, K. and Zerial, M.** (1992). The complexity of the Rab and Rho GTP-binding protein subfamilies revealed by a PCR cloning approach. *Gene* **112**, 261-4.
- Chen, S. and Kimelman, D.** (2000). The role of the yolk syncytial layer in germ layer patterning in zebrafish. *Development* **127**, 4681-9.
- Chen, Y. and Scheller, R.** (2001). SNARE-Mediated Membrane Fusion. *Nature Reviews Molecular Cell Biology* **2**, 98-106.
- Chen, Y. and Schier, A. F.** (2001). The zebrafish Nodal signal Squint functions as a morphogen. *Nature* **411**, 607-610.
- Chen, Y. and Struhl, G.** (1996). Dual roles for patched in sequestering and transducing Hedgehog. *Cell* **87**, 553-63.
- Cheney, R. E. and Rodriguez, O. C.** (2001). Cell biology. A switch to release the motor. *Science* **293**, 1263-4.
- Cheng, J. C., Miller, A. L. and Webb, S. E.** (2004). Organization and function of microfilaments during late epiboly in zebrafish embryos. *Dev Dyn* **231**, 313-23.

- Chiariello, M., Bruni, B. and Bucci, C.** (1999). The small GTPases Rab5a, Rab5b and Rab5c are differentially phosphorylated in vitro. *FEBA Letters* **453**, 20-24.
- Child, C. M.** (1915). Individuality in Organisms.
- Child, C. M.** (1941). Patterns and Problems of Development.
- Christoforidis, S., McBride, H. M., Burgoyne, R. D. and Zerial, M.** (1999a). The Rab5 effector EEA1 is a core component of endosome docking. *Nature* **397**, 621-5.
- Christoforidis, S., Miaczynska, M., Ashman, K., Wilm, M., Zhao, L., Yip, S. C., Waterfield, M. D., Backer, J. M. and Zerial, M.** (1999b). Phosphatidylinositol-3-OH kinases are Rab5 effectors. *Nat Cell Biol* **1**, 249-52.
- Christoforidis, S. and Zerial, M.** (2000). Purification and identification of novel Rab effectors using affinity chromatography. *Methods* **20**, 403-10.
- Ciardiello, F., Dono, R., Kim, N., Persico, M. G. and Salomon, D. S.** (1991). Expression of cripto, a novel gene of the epidermal growth factor gene family, leads to in vitro transformation of a normal mouse mammary epithelial cell line. *Cancer Res* **51**, 1051-4.
- Clements, D., Friday, R. V. and Woodland, H. R.** (1999). Mode of action of VegT in mesoderm and endoderm formation. *Development* **126**, 4903-11.
- Clements, D. and Woodland, H. R.** (2003). VegT induces endoderm by a self-limiting mechanism and by changing the competence of cells to respond to TGF-beta signals. *Dev Biol* **258**, 454-63.
- Colicelli, J.** (2004). Human RAS superfamily proteins and related GTPases. *Sci STKE* **2004**, RE13.
- Cong, F., Schweizer, L. and Varmus, H.** (2004). Wnt signals across the plasma membrane to activate the beta-catenin pathway by forming oligomers containing its receptors, Frizzled and LRP. *Development* **131**, 5103-15.
- Connors, S. A., Trout, J., Ekker, M. and Mullins, M. C.** (1999). The role of tolloid/mini fin in dorsoventral pattern formation of the zebrafish embryo. *Development* **126**, 3119-30.
- Cooper, A. A., Gitler, A. D., Cashikar, A., Haynes, C. M., Hill, K. J., Bhullar, B., Liu, K., Xu, K., Strathearn, K. E., Liu, F. et al.** (2006). Alpha-synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science* **313**, 324-8.
- Costa, M. C., Mani, F., Santoro, W., Jr., Espreafico, E. M. and Larson, R. E.** (1999). Brain myosin-V, a calmodulin-carrying myosin, binds to calmodulin-independent protein kinase II and activates its kinase activity. *J Biol Chem* **274**, 15811-9.
- Coutinho, P., Parsons, M. J., Thomas, K. A., Hirst, E. M., Saude, L., Campos, I., Williams, P. H. and Stemple, D. L.** (2004). Differential requirements for COPI transport during vertebrate early development. *Dev Cell* **7**, 547-58.
- Couve-Privat, S., Le Bret, M., Traiffort, E., Queille, S., Coulombe, J., Bouadjar, B., Avril, M. F., Ruat, M., Sarasin, A. and Daya-Grosjean, L.** (2004). Functional analysis of novel sonic hedgehog gene mutations identified in Basal cell carcinomas from xeroderma pigmentosum patients. *Cancer Res* **64**, 3559-65.
- Cremers, F. P., Molloy, C. M., van de Pol, D. J., van den Hurk, J. A., Bach, I., Geurts van Kessel, A. H. and Ropers, H. H.** (1992). An autosomal homologue of the choroideremia gene colocalizes with the Usher syndrome type II locus on the distal part of chromosome 1q. *Hum Mol Genet* **1**, 71-5.
- Cui, W. W., Low, S. E., Hirata, H., Saint-Amant, L., Geisler, R., Hume, R. I. and Kuwada, J. Y.** (2005). The zebrafish shocked gene encodes a glycine

- transporter and is essential for the function of early neural circuits in the CNS. *J Neurosci* **25**, 6610-20.
- Curtis, L. M. and Gluck, S.** (2005). Distribution of Rab GTPases in mouse kidney and comparison with vacuolar H⁺-ATPase. *Nephron Physiol* **100**, p31-42.
- Daggett, D. F., Boyd, C. A., Gautier, P., Bryson-Richardson, R. J., Thisse, C., Thisse, B., Amacher, S. L. and Currie, P. D.** (2004). Developmentally restricted actin-regulatory molecules control morphogenetic cell movements in the zebrafish gastrula. *Curr Biol* **14**, 1632-8.
- de Larco, J. E. and Todaro, G. J.** (1978). Growth factors from murine sarcoma virus-transformed cells. *Proc Natl Acad Sci U S A* **75**, 4001-5.
- Denef, N., Neubuser, D., Perez, L. and Cohen, S. M.** (2000). Hedgehog induces opposite changes in turnover and subcellular localization of patched and smoothened. *Cell* **102**, 521-31.
- Deryck, R. and Miyazono, K.** (2006). The TGF-β Family.
- Dick, A., Hild, M., Bauer, H., Imai, Y., Maifeld, H., Schier, A. F., Talbot, W. S., Bouwmeester, T. and Hammerschmidt, M.** (2000). Essential role of Bmp7 (snailhouse) and its prodomain in dorsoventral patterning of the zebrafish embryo. *Development* **127**, 343-54.
- Dougan, S. T., Warga, R. M., Kane, D. A., Schier, A. F. and Talbot, W. S.** (2003). The role of the zebrafish nodal-related genes squint and cyclops in patterning of mesendoderm. *Development* **130**, 1837-51.
- Draper, B. W., Morcos, P. A. and Kimmel, C. B.** (2001). Inhibition of zebrafish fgf8 pre-mRNA splicing with morpholino oligos: a quantifiable method for gene knockdown. *Genesis* **30**, 154-6.
- Draper, B. W., Stock, D. W. and Kimmel, C. B.** (2003). Zebrafish fgf24 functions with fgf8 to promote posterior mesodermal development. *Development* **130**, 4639-54.
- Dubois, L., Lecourtois, M., Alexandre, C., Hirst, E. and Vincent, J. P.** (2001). Regulated endocytic routing modulates wingless signaling in Drosophila embryos. *Cell* **105**, 613-24.
- Dumas, J. J., Zhu, Z., Connolly, J. L. and Lambright, D. G.** (1999). Structural basis of activation and GTP hydrolysis in Rab proteins. *Structure* **7**, 413-23.
- Dutton, K. A., Pauliny, A., Lopes, S. S., Elworthy, S., Carney, T. J., Rauch, J., Geisler, R., Haffter, P. and Kelsh, R. N.** (2001). Zebrafish colourless encodes sox10 and specifies non-ectomesenchymal neural crest fates. *Development* **128**, 4113-25.
- Ebisawa, T., Fukuchi, M., Murakami, G., Chiba, T., Tanaka, K., Imamura, T. and Miyazono, K.** (2001). Smurfl interacts with transforming growth factor-beta type I receptor through Smad7 and induces receptor degradation. *J Biol Chem* **276**, 12477-80.
- Echard, A., Jollivet, F., Martinez, O., Lacapere, J. J., Rousselet, A., Janoueix-Lerosey, I. and Goud, B.** (1998). Interaction of a Golgi-associated kinesin-like protein with Rab6. *Science* **279**, 580-5.
- Echard, A., Opdam, F. J., de Leeuw, H. J., Jollivet, F., Savelkoul, P., Hendriks, W., Voorberg, J., Goud, B. and Fransen, J. A.** (2000). Alternative splicing of the human Rab6A gene generates two close but functionally different isoforms. *Mol Biol Cell* **11**, 3819-33.
- Eggenschwiler, J., Espinoza and Anderson, K.** (2001). Rab23 is an essential negative regulator of the mouse Sonic hedgehog signalling pathway. *Nature* **412**, 194 - 198.

- Eggenschwiler, J. T. and Anderson, K. V.** (2000). Dorsal and lateral fates in the mouse neural tube require the cell-autonomous activity of the open brain gene. *Dev Biol* **227**, 648-60.
- Eggenschwiler, J. T., Bulgakov, O. V., Qin, J., Li, T. and Anderson, K. V.** (2006). Mouse Rab23 regulates hedgehog signaling from smoothened to Gli proteins. *Dev Biol* **290**, 1-12.
- Eisen, J. S. and Weston, J. A.** (1993). Development of the neural crest in the zebrafish. *Dev Biol* **159**, 50-9.
- Eldar, A., Dorfman, R., Weiss, D., Ashe, H., Shilo, B. Z. and Barkai, N.** (2002). Robustness of the BMP morphogen gradient in Drosophila embryonic patterning. *Nature* **419**, 304-8.
- Elde, N. C., Morgan, G., Winey, M., Sperling, L. and Turkewitz, A. P.** (2005). Elucidation of clathrin-mediated endocytosis in tetrahymena reveals an evolutionarily convergent recruitment of dynamin. *PLoS Genet* **1**, e52.
- Entchev, E. V. and Gonzalez-Gaitan, M. A.** (2002). Morphogen gradient formation and vesicular trafficking. *Traffic* **3**, 98-109.
- Entchev, E. V., Schwabedissen, A. and Gonzalez-Gaitan, M.** (2000). Gradient formation of the TGF-beta homolog Dpp. *Cell* **103**, 981-91.
- Erter, C. E., Wilm, T. P., Basler, N., Wright, C. V. and Solnica-Krezel, L.** (2001). Wnt8 is required in lateral mesendodermal precursors for neural posteriorization in vivo. *Development* **128**, 3571-83.
- Essner, J. J., Amack, J. D., Nyholm, M. K., Harris, E. B. and Yost, H. J.** (2005). Kupffer's vesicle is a ciliated organ of asymmetry in the zebrafish embryo that initiates left-right development of the brain, heart and gut. *Development* **132**, 1247-60.
- Evans, T. M., Ferguson, C., Wainwright, B. J., Parton, R. G. and Wicking, C.** (2003). Rab23, a negative regulator of hedgehog signaling, localizes to the plasma membrane and the endocytic pathway. *Traffic* **4**, 869-84.
- Fainsod, A., Deissler, K., Yelin, R., Marom, K., Epstein, M., Pillemer, G., Steinbeisser, H. and Blum, M.** (1997). The dorsalizing and neural inducing gene follistatin is an antagonist of BMP-4. *Mech Dev* **63**, 39-50.
- Fekany, K., Yamanaka, Y., Leung, T., Sirotnik, H. I., Topczewski, J., Gates, M. A., Hibi, M., Renucci, A., Stemple, D., Radbill, A. et al.** (1999). The zebrafish bozozok locus encodes Dharma, a homeodomain protein essential for induction of gastrula organizer and dorsoanterior embryonic structures. *Development* **126**, 1427-38.
- Feldman, B., Gates, M., Egan, E., Dougan, S., Rennebeck, G., Sirotnik, H., Schier, A. and Talbot, W.** (1998). Zebrafish organizer development and germ-layer formation require nodal-related signals. *Nature* **395**, 181-185.
- Finger, F. P., Hughes, T. E. and Novick, P.** (1998). Sec3p is a spatial landmark for polarized secretion in budding yeast. *Cell* **92**, 559-71.
- Fischer von Mollard, G., Mignery, G. A., Baumert, M., Perin, M. S., Hanson, T. J., Burger, P. M., Jahn, R. and Sudhof, T. C.** (1990). rab3 is a small GTP-binding protein exclusively localized to synaptic vesicles. *Proc Natl Acad Sci U S A* **87**, 1988-92.
- Fischer von Mollard, G., Stahl, B., Khokhlatchev, A., Sudhof, T. and R, J.** (1994). Rab3C is a synaptic vesicle protein that dissociates from synaptic vesicles after stimulation of exocytosis. *J. Biol. Chem.* **269**, 10971-10974.

- Fujii, R., Yamashita, S., Hibi, M. and Hirano, T.** (2000). Asymmetric p38 activation in zebrafish: its possible role in symmetric and synchronous cleavage. *J Cell Biol* **150**, 1335-48.
- Fujimoto, L. M., Roth, R., Heuser, J. E. and Schmid, S. L.** (2000). Actin assembly plays a variable, but not obligatory role in receptor-mediated endocytosis in mammalian cells. *Traffic* **1**, 161-71.
- Furthauer, M., Van Celst, J., Thisse, C. and Thisse, B.** (2004). Fgf signalling controls the dorsoventral patterning of the zebrafish embryo. *Development* **131**, 2853-64.
- Gallwitz, D., Donath, C. and Sander, C.** (1983). A yeast gene encoding a protein homologous to the human c-has/bas proto-oncogene product. *Nature* **306**, 704-7.
- Gardiner, M. R., Daggett, D. F., Zon, L. I. and Perkins, A. C.** (2005). Zebrafish KLF4 is essential for anterior mesendoderm/pre-polster differentiation and hatching. *Dev Dyn* **234**, 992-6.
- Garrett, M. D., Kabcenell, A. K., Zahner, J. E., Kaibuchi, K., Sasaki, T., Takai, Y., Cheney, C. M. and Novick, P. J.** (1993). Interaction of Sec4 with GDI proteins from bovine brain, *Drosophila melanogaster* and *Saccharomyces cerevisiae*. Conservation of GDI membrane dissociation activity. *FEBS Lett* **331**, 233-8.
- Gelbart, W. M.** (1989). The decapentaplegic gene: a TGF-beta homologue controlling pattern formation in *Drosophila*. *Development* **107 Suppl**, 65-74.
- Geng, X., Xiao, L., Lin, G. F., Hu, R., Wang, J. H., Rupp, R. A. and Ding, X.** (2003). Lef/Tcf-dependent Wnt/beta-catenin signaling during Xenopus axis specification. *FEBS Lett* **547**, 1-6.
- Geppert, M., Goda, Y., Stevens, C. F. and Sudhof, T. C.** (1997). The small GTP-binding protein Rab3A regulates a late step in synaptic vesicle fusion. *Nature* **387**, 810-4.
- Geyer, M., Schweins, T., Herrmann, C., Prisner, T., Wittinghofer, A. and Kalbitzer, H. R.** (1996). Conformational transitions in p21ras and in its complexes with the effector protein Raf-RBD and the GTPase activating protein GAP. *Biochemistry* **35**, 10308-20.
- Gibbs, D., Azarian, S. M., Lillo, C., Kitamoto, J., Klomp, A. E., Steel, K. P., Libby, R. T. and Williams, D. S.** (2004). Role of myosin VIIa and Rab27a in the motility and localization of RPE melanosomes. *J Cell Sci* **117**, 6473-83.
- Gilbert, S. F.** (1997). CHAPTER 3 :: "REDISCOVERY" OF THE MORPHOGENETIC FIELD :: THE "RE-DISCOVERY" OF MORPHOGENIC FIELDS. In *Developmental Biology (DevBio.com)*, (ed).
- Gimlich, R. L. and Gerhart, J. C.** (1984). Early cellular interactions promote embryonic axis formation in *Xenopus laevis*. *Dev Biol* **104**, 117-30.
- Goering, L. M., Hoshijima, K., Hug, B., Bisgrove, B., Kispert, A. and Grunwald, D. J.** (2003). An interacting network of T-box genes directs gene expression and fate in the zebrafish mesoderm. *Proc Natl Acad Sci U S A* **100**, 9410-5.
- Gonzalez-Gaitan, M., Capdevila, M. P. and Garcia-Bellido, A.** (1994). Cell proliferation patterns in the wing imaginal disc of *Drosophila*. *Mech Dev* **46**, 183-200.
- Gonzalez-Sancho, J. M., Aguilera, O., Garcia, J. M., Pendas-Franco, N., Pena, C., Cal, S., Garcia de Herreros, A., Bonilla, F. and Munoz, A.** (2005). The Wnt antagonist DICKKOPF-1 gene is a downstream target of beta-catenin/TCF and is downregulated in human colon cancer. *Oncogene* **24**, 1098-103.

- Gonzalez, F., Swales, L., Bejsovec, A., Skaer, H. and Martinez Arias, A.** (1991). Secretion and movement of wingless protein in the epidermis of the Drosophila embryo. *Mech Dev* **35**, 43-54.
- Gore, A. and Sampath, K.** (2002). Localization of transcripts of the Zebrafish morphogen Squint is dependent on egg activation and the microtubule cytoskeleton. *Mechanisms in Development* **112**, 153-156.
- Gore, A. V., Maegawa, S., Cheong, A., Gilligan, P. C., Weinberg, E. S. and Sampath, K.** (2005). The zebrafish dorsal axis is apparent at the four-cell stage. *Nature* **438**, 1030-5.
- Gorvel, J. P., Chavrier, P., Zerial, M. and Gruenberg, J.** (1991). rab5 controls early endosome fusion in vitro. *Cell* **64**, 915-25.
- Graff, J. M., Thies, R. S., Song, J. J., Celeste, A. J. and Melton, D. A.** (1994). Studies with a Xenopus BMP receptor suggest that ventral mesoderm-inducing signals override dorsal signals in vivo. *Cell* **79**, 169-79.
- Granato, M., van Eeden, F. J., Schach, U., Trowe, T., Brand, M., Furutani-Seiki, M., Haffter, P., Hammerschmidt, M., Heisenberg, C. P., Jiang, Y. J. et al.** (1996). Genes controlling and mediating locomotion behavior of the zebrafish embryo and larva. *Development* **123**, 399-413.
- Grant, K. A., Raible, D. W. and Piotrowski, T.** (2005). Regulation of latent sensory hair cell precursors by glia in the zebrafish lateral line. *Neuron* **45**, 69-80.
- Greco, V., Hannus, M. and Eaton, S.** (2001). Argosomes: a potential vehicle for the spread of morphogens through epithelia. *Cell* **106**, 633-45.
- Griffin, K. and Kimelman, D.** (2003). Interplay between FGF, *one-eyed pinhead*, and T-box transcription factors during zebrafish development. *Developmental Biology* **264**, 456-466.
- Griscelli, C., Durandy, A., Guy-Grand, D., Daguillard, F., Herzog, C. and Prunieras, M.** (1978). A syndrome associating partial albinism and immunodeficiency. *Am J Med* **65**, 691-702.
- Gritsman, K., Talbot, W. and Schier, A.** (2000). Nodal signaling patterns the organizer. *Development* **127**, 921-932.
- Gritsman, K., Zhang, J., Cheng, S., Heckscher, E., Talbot, W. S. and Schier, A. F.** (1999). The EGF-CFC protein one-eyed pinhead is essential for nodal signaling. *Cell* **97**, 121-32.
- Grosshans, B. L., Ortiz, D. and Novick, P.** (2006). Rabs and their effectors: achieving specificity in membrane traffic. *Proc Natl Acad Sci U S A* **103**, 11821-7.
- Gruenberg, J. and Howell, K. E.** (1989). Membrane traffic in endocytosis: insights from cell-free assays. *Annu Rev Cell Biol* **5**, 453-81.
- Guger, K. A. and Gumbiner, B. M.** (1995). beta-Catenin has Wnt-like activity and mimics the Nieuwkoop signaling center in Xenopus dorsal-ventral patterning. *Dev Biol* **172**, 115-25.
- Gunther, T., Struwe, M., Aguzzi, A. and Schughart, K.** (1994). Open brain, a new mouse mutant with severe neural tube defects, shows altered gene expression patterns in the developing spinal cord. *Development* **120**, 3119-30.
- Guo, A., Wang, T., Ng, E. L., Aulia, S., Chong, K. H., Teng, F. Y., Wang, Y. and Tang, B. L.** (2006). Open brain gene product Rab23: expression pattern in the adult mouse brain and functional characterization. *J Neurosci Res* **83**, 1118-27.
- Guo, W., Roth, D., Walch-Solimena, C. and Novick, P.** (1999). The exocyst is an effector for Sec4p, targeting secretory vesicles to sites of exocytosis. *Embo J* **18**, 1071-80.

- Gurdon, J. B., Fairman, S., Mohun, T. J. and Brennan, S.** (1985). Activation of muscle-specific actin genes in *Xenopus* development by an induction between animal and vegetal cells of a blastula. *Cell* **41**, 913-22.
- Gurdon, J. B., Harger, P., Mitchell, A. and Lemaire, P.** (1994). Activin signalling and response to a morphogen gradient. *Nature* **371**, 487-92.
- Gurkan, C., Lapp, H., Alory, C., Su, A. I., Hogenesch, J. B. and Balch, W. E.** (2005). Large-scale profiling of Rab GTPase trafficking networks: the membrane. *Mol Biol Cell* **16**, 3847-64.
- Haas, A. K., Fuchs, E., Kopajtich, R. and Barr, F. A.** (2005). A GTPase-activating protein controls Rab5 function in endocytic trafficking. *Nat Cell Biol* **7**, 887-93.
- Halloran, M. C. and Berndt, J. D.** (2003). Current progress in neural crest cell motility and migration and future prospects for the zebrafish model system. *Dev Dyn* **228**, 497-513.
- Hammer, J. and Wu, X.** (2002). Rabs grab motors: defining the connections between Rab GTPases and motor proteins. *Current Opinion in Cell Biology* **14**, 69-75.
- Hammerschmidt, M., Pelegri, F., Mullins, M. C., Kane, D. A., Brand, M., van Eeden, F. J., Furutani-Seiki, M., Granato, M., Haffter, P., Heisenberg, C. P. et al.** (1996a). Mutations affecting morphogenesis during gastrulation and tail formation in the zebrafish, *Danio rerio*. *Development* **123**, 143-51.
- Hammerschmidt, M., Serbedzija, G. N. and McMahon, A. P.** (1996b). Genetic analysis of dorsoventral pattern formation in the zebrafish: requirement of a BMP-like ventralizing activity and its dorsal repressor. *Genes Dev* **10**, 2452-61.
- Hashimoto, H., Itoh, M., Yamanaka, Y., Yamashita, S., Shimizu, T., Solnica-Krezel, L., Hibi, M. and Hirano, T.** (2000). Zebrafish Dkk1 functions in forebrain specification and axial mesendoderm formation. *Dev Biol* **217**, 138-52.
- Hatta, K., Kimmel, C., Ho, R. and Walker, C.** (1991). The cyclops mutation blocks specification of the floor plate of the zebrafish central nervous system. *Nature* **350**, 339-341.
- Haubruck, H., Engelke, U., Mertins, P. and Gallwitz, D.** (1990). Structural and functional analysis of *ypt2*, an essential ras-related gene in the fission yeast *Schizosaccharomyces pombe* encoding a Sec4 protein homologue. *Embo J* **9**, 1957-62.
- Haubruck, H., Prange, R., Vorgias, C. and Gallwitz, D.** (1989). The ras-related mouse *ypt1* protein can functionally replace the *YPT1* gene product in yeast. *Embo J* **8**, 1427-32.
- Hawley, S. H., Wunnenberg-Stapleton, K., Hashimoto, C., Laurent, M. N., Watabe, T., Blumberg, B. W. and Cho, K. W.** (1995). Disruption of BMP signals in embryonic *Xenopus* ectoderm leads to direct neural induction. *Genes Dev* **9**, 2923-35.
- Haycraft, C. J., Banisz, B., Aydin-Son, Y., Zhang, Q., Michaud, E. J. and Yoder, B. K.** (2005). Gli2 and Gli3 localize to cilia and require the intraflagellar transport protein polaris for processing and function. *PLoS Genet* **1**, e53.
- Heisenberg, C. P., Tada, M., Rauch, G. J., Saude, L., Concha, M. L., Geisler, R., Stemple, D. L., Smith, J. C. and Wilson, S. W.** (2000). Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. *Nature* **405**, 76-81.
- Heldin, C. H., Miyazono, K. and ten Dijke, P.** (1997). TGF-beta signalling from cell membrane to nucleus through SMAD proteins. *Nature* **390**, 465-71.

- Hikasa, H., Shibata, M., Hiratani, I. and Taira, M.** (2002). The Xenopus receptor tyrosine kinase Xror2 modulates morphogenetic movements of the axial mesoderm and neuroectoderm via Wnt signaling. *Development* **129**, 5227-39.
- Hild, M., Dick, A., Rauch, G. J., Meier, A., Bouwmeester, T., Haffter, P. and Hammerschmidt, M.** (1999). The smad5 mutation somitabun blocks Bmp2b signaling during early dorsoventral patterning of the zebrafish embryo. *Development* **126**, 2149-59.
- Hill, E., Clarke, M. and Barr, F. A.** (2000). The Rab6-binding kinesin, Rab6-KIFL, is required for cytokinesis. *Embo J* **19**, 5711-9.
- Hinck, L., Nelson, W. J. and Papkoff, J.** (1994). Wnt-1 modulates cell-cell adhesion in mammalian cells by stabilizing beta-catenin binding to the cell adhesion protein cadherin. *J Cell Biol* **124**, 729-41.
- Holloway, B.** (2006 unpublished data). Betty boop, the Zebrafish Homologue of MAPKAPK2, Regulates Epiboly. *7th International meeting on Zebrafish development and genetics*.
- Horowitz, J. M., Mikuckis, G. M. and Longshore, M. A.** (1980). The response of single melanophores to extracellular and intracellular iontophoretic injection of melanocyte-stimulating hormone. *Endocrinology* **106**, 770-7.
- Houart, C., Westerfield, M. and Wilson, S. W.** (1998). A small population of anterior cells patterns the forebrain during zebrafish gastrulation. *Nature* **391**, 788-92.
- Houlden, H., King, R. H., Muddle, J. R., Warner, T. T., Reilly, M. M., Orrell, R. W. and Ginsberg, L.** (2004). A novel RAB7 mutation associated with ulceromutilating neuropathy. *Ann Neurol* **56**, 586-90.
- Hsiung, F., Ramirez-Weber, F. A., Iwaki, D. D. and Kornberg, T. B.** (2005). Dependence of Drosophila wing imaginal disc cytonemes on Decapentaplegic. *Nature* **437**, 560-3.
- Hsu, H. J., Liang, M. R., Chen, C. T. and Chung, B. C.** (2006). Pregnenolone stabilizes microtubules and promotes zebrafish embryonic cell movement. *Nature* **439**, 480-3.
- Imai, Y., Gates, M. A., Melby, A. E., Kimelman, D., Schier, A. F. and Talbot, W. S.** (2001). The homeobox genes vox and vent are redundant repressors of dorsal fates in zebrafish. *Development* **128**, 2407-20.
- Inohaya, K., Yasumasu, S., Ishimaru, M., Ohyama, A., Iuchi, I. and Yamagami, K.** (1995). Temporal and spatial patterns of gene expression for the hatching enzyme in the teleost embryo, Oryzias latipes. *Dev Biol* **171**, 374-85.
- Inohaya, K., Yasumasu, S., Yasumasu, I., Iuchi, I. and Yamagami, K.** (1999). Analysis of the origin and development of hatching gland cells by transplantation of the embryonic shield in the fish, Oryzias latipes. *Dev Growth Differ* **41**, 557-66.
- Jesuthasan, S. and Stahle, U.** (1997). Dynamic microtubules and specification of the zebrafish embryonic axis. *Curr Biol* **7**, 31-42.
- Johnston, P. A., Archer, B. T., 3rd, Robinson, K., Mignery, G. A., Jahn, R. and Sudhof, T. C.** (1991). rab3A attachment to the synaptic vesicle membrane mediated by a conserved polyisoprenylated carboxy-terminal sequence. *Neuron* **7**, 101-9.
- Kane, D. A., Hammerschmidt, M., Mullins, M. C., Maischein, H. M., Brand, M., van Eeden, F. J., Furutani-Seiki, M., Granato, M., Haffter, P., Heisenberg, C. P. et al.** (1996). The zebrafish epiboly mutants. *Development* **123**, 47-55.
- Kane, D.A., Maischein, H.M., Brand, M., van Eeden, F.J., Furutani-Seiki, M., Granato, M., Haffter, P., Hammerschmidt, M., Heisenberg-, C.P., Jiang, Y.J.,**

- Kelsh, R.N., Mullins, M.C., Odenthal, J., Warga, R.M., and Nüsslein-Volhard, C.** (1996b). The zebrafish early arrest mutants. *Development* **123**, 57-66.
- Kane, D. A. and Kimmel, C. B.** (1993). The zebrafish midblastula transition. *Development* **119**, 447-56.
- Kane, D. A., McFarland, K. N. and Warga, R. M.** (2005). Mutations in half baked/E-cadherin block cell behaviors that are necessary for teleost epiboly. *Development* **132**, 1105-16.
- Kapfhamer, D., Valladares, O., Sun, Y., Nolan, P., Rux, J., Arnold, S., Veasey, S. and Bucan, M.** (2002). Mutations in Rab3a alter circadian period and homeostatic response to sleep loss in the mouse. *Nature Genetics* **32**, 290-295.
- Karcher, R. L., Roland, J. T., Zappacosta, F., Huddleston, M. J., Annan, R. S., Carr, S. A. and Gelfand, V. I.** (2001). Cell cycle regulation of myosin-V by calcium/calmodulin-dependent protein kinase II. *Science* **293**, 1317-20.
- Kawahara, A. and Dawid, I. B.** (2000). Expression of the Kruppel-like zinc finger gene biklf during zebrafish development. *Mech Dev* **97**, 173-6.
- Kawahara, A. and Dawid, I. B.** (2001). Critical role of biklf in erythroid cell differentiation in zebrafish. *Curr Biol* **11**, 1353-7.
- Kawahara, A., Wilm, T., Solnica-Krezel, L. and Dawid, I. B.** (2000a). Antagonistic role of vega1 and bozozok/dharma homeobox genes in organizer formation. *Proc Natl Acad Sci U S A* **97**, 12121-6.
- Kawahara, A., Wilm, T., Solnica-Krezel, L. and Dawid, I. B.** (2000b). Functional interaction of vega2 and goosecoid homeobox genes in zebrafish. *Genesis* **28**, 58-67.
- Kee, Y., Yoo, J. S., Hazuka, C. D., Peterson, K. E., Hsu, S. C. and Scheller, R. H.** (1997). Subunit structure of the mammalian exocyst complex. *Proc Natl Acad Sci U S A* **94**, 14438-43.
- Keller, R. E. and Trinkaus, J. P.** (1987). Rearrangement of enveloping layer cells without disruption of the epithelial permeability barrier as a factor in Fundulus epiboly. *Dev Biol* **120**, 12-24.
- Kelly, C., Chin, A. J., Leatherman, J. L., Kozlowski, D. J. and Weinberg, E. S.** (2000). Maternally controlled (beta)-catenin-mediated signaling is required for organizer formation in the zebrafish. *Development* **127**, 3899-911.
- Kelly, G. M., Erezyilmaz, D. F. and Moon, R. T.** (1995). Induction of a secondary embryonic axis in zebrafish occurs following the overexpression of beta-catenin. *Mech Dev* **53**, 261-73.
- Kelsh, R. N., Brand, M., Jiang, Y. J., Heisenberg, C. P., Lin, S., Haffter, P., Odenthal, J., Mullins, M. C., van Eeden, F. J., Furutani-Seiki, M. et al.** (1996). Zebrafish pigmentation mutations and the processes of neural crest development. *Development* **123**, 369-89.
- Kelsh, R. N. and Eisen, J. S.** (2000). The zebrafish colourless gene regulates development of non-ectomesenchymal neural crest derivatives. *Development* **127**, 515-25.
- Kerszberg, M. and Wolpert, L.** (1998). Mechanisms for positional signalling by morphogen transport: a theoretical study. *J Theor Biol* **191**, 103-14.
- Khosravi-Far, R., Lutz, R. J., Cox, A. D., Conroy, L., Bourne, J. R., Sinensky, M., Balch, W. E., Buss, J. E. and Der, C. J.** (1991). Isoprenoid modification of rab proteins terminating in CC or CXC motifs. *Proc Natl Acad Sci U S A* **88**, 6264-8.
- Kierszenbaum, A.** (2000). Fusion of membranes during the acrosome reaction: a tale of two SNAREs. *Molecular Reproduction and Development* **57**, 309-10.

- Kimelman, D. and Kirschner, M.** (1987). Synergistic induction of mesoderm by FGF and TGF-beta and the identification of an mRNA coding for FGF in the early Xenopus embryo. *Cell* **51**, 869-77.
- Kimmel, C., Ballard, W., Kimmel, S., Ullmann, B. and Schilling, T.** (1995). Stages of Embryonic Development of the Zebrafish. *Developmental Dynamics* **203**, 253-310.
- Kimmel, C. B., Kane, D. A., Walker, C., Warga, R. M. and Rothman, M. B.** (1989). A mutation that changes cell movement and cell fate in the zebrafish embryo. *Nature* **337**, 358-62.
- Kimmel, C. B. and Law, R. D.** (1985). Cell lineage of zebrafish blastomeres. II. Formation of the yolk syncytial layer. *Dev Biol* **108**, 86-93.
- Kimmel, C. B., Warga, R. M. and Schilling, T. F.** (1990). Origin and organization of the zebrafish fate map. *Development* **108**, 581-94.
- Kinsella, B. T. and Maltese, W. A.** (1991). rab GTP-binding proteins implicated in vesicular transport are isoprenylated in vitro at cysteines within a novel carboxyl-terminal motif. *J Biol Chem* **266**, 8540-4.
- Knecht, A. K., Good, P. J., Dawid, I. B. and Harland, R. M.** (1995). Dorsal-ventral patterning and differentiation of noggin-induced neural tissue in the absence of mesoderm. *Development* **121**, 1927-35.
- Kodjabachian, L., Dawid, I. B. and Toyama, R.** (1999). Gastrulation in zebrafish: what mutants teach us. *Dev Biol* **213**, 231-45.
- Koenig, B. B., Cook, J. S., Wolsing, D. H., Ting, J., Tiesman, J. P., Correa, P. E., Olson, C. A., Pecquet, A. L., Ventura, F., Grant, R. A. et al.** (1994). Characterization and cloning of a receptor for BMP-2 and BMP-4 from NIH 3T3 cells. *Mol Cell Biol* **14**, 5961-74.
- Kohler, K., Louvard, D. and Zahraoui, A.** (2004). Rab13 regulates PKA signaling during tight junction assembly. *J Cell Biol* **165**, 175-80.
- Kohler, K. and Zahraoui, A.** (2005). Tight junction: a co-ordinator of cell signalling and membrane trafficking. *Biol Cell* **97**, 659-65.
- Koos, D. S. and Ho, R. K.** (1999). The nieuwkoid/dharma homeobox gene is essential for bmp2b repression in the zebrafish pregastrula. *Dev Biol* **215**, 190-207.
- Kornbluth, S., Paulson, K. E. and Hanafusa, H.** (1988). Novel tyrosine kinase identified by phosphotyrosine antibody screening of cDNA libraries. *Mol Cell Biol* **8**, 5541-4.
- Lakadamyali, M., Rust, M. J. and Zhuang, X.** (2006). Ligands for clathrin-mediated endocytosis are differentially sorted into distinct populations of early endosomes. *Cell* **124**, 997-1009.
- Lamaze, C., Fujimoto, L. M., Yin, H. L. and Schmid, S. L.** (1997). The actin cytoskeleton is required for receptor-mediated endocytosis in mammalian cells. *J Biol Chem* **272**, 20332-5.
- Lamb, T. M. and Harland, R. M.** (1995). Fibroblast growth factor is a direct neural inducer, which combined with noggin generates anterior-posterior neural pattern. *Development* **121**, 3627-36.
- Lambert, J., Vancoillie, G. and Naeyaert, J. M.** (1999). Molecular motors and their role in pigmentation. *Cell Mol Biol (Noisy-le-grand)* **45**, 905-18.
- Lanzetti, L., Palamidessi, A., Areces, L., Scita, G. and Di Fiore, P. P.** (2004). Rab5 is a signalling GTPase involved in actin remodelling by receptor tyrosine kinases. *Nature* **429**, 309-14.

- Lanzetti, L., Rybin, V., Malabarba, M. G., Christoforidis, S., Scita, G., Zerial, M. and Di Fiore, P. P.** (2000). The Eps8 protein coordinates EGF receptor signalling through Rac and trafficking through Rab5. *Nature* **408**, 374-7.
- Lawrence, P.** (2004). A Wigglesworth classic: how cells make patterns. *J Exp Biol* **207**, 192-3.
- Lawrence, P. A.** (2001). Morphogens: how big is the big picture? *Nat Cell Biol* **3**, E151-4.
- Lazar, T., Gotte, M. and Gallwitz, D.** (1997). Vesicular transport: how many Ypt/Rab-GTPases make a eukaryotic cell? *TIBS* **22**, 468-472.
- Le Galic, L. and Fort, P.** (1997). Structure of the human ARHG locus encoding the Rho/Rac-like RhoG GTPase. *Genomics* **42**, 157-60.
- Le, P. U. and Nabi, I. R.** (2003). Distinct caveolae-mediated endocytic pathways target the Golgi apparatus and the endoplasmic reticulum. *J Cell Sci* **116**, 1059-71.
- Le Roy, C. and Wrana, J. L.** (2005). Clathrin- and non-clathrin-mediated endocytic regulation of cell signalling. *Nat Rev Mol Cell Biol* **6**, 112-26.
- Lecuit, T., Brook, W. J., Ng, M., Calleja, M., Sun, H. and Cohen, S. M.** (1996). Two distinct mechanisms for long-range patterning by Decapentaplegic in the *Drosophila* wing. *Nature* **381**, 387-93.
- Lekven, A. C., Thorpe, C. J., Waxman, J. S. and Moon, R. T.** (2001). Zebrafish wnt8 encodes two wnt8 proteins on a bicistronic transcript and is required for mesoderm and neurectoderm patterning. *Dev Cell* **1**, 103-14.
- Lele, Z., Nowak, M. and Hammerschmidt, M.** (2001). Zebrafish admp is required to restrict the size of the organizer and to promote posterior and ventral development. *Dev Dyn* **222**, 681-7.
- Leung, K. F., Baron, R. and Seabra, M. C.** (2006). Thematic review series: lipid posttranslational modifications. geranylgeranylation of Rab GTPases. *J Lipid Res* **47**, 467-75.
- Leung, T., Soll, I., Arnold, S. J., Kemler, R. and Driever, W.** (2003). Direct binding of Lef1 to sites in the boz promoter may mediate pre-midblastula-transition activation of boz expression. *Dev Dyn* **228**, 424-32.
- Levi, V., Serpinskaya, A. S., Gratton, E. and Gelfand, V.** (2006). Organelle transport along microtubules in *Xenopus melanophores*: evidence for cooperation between multiple motors. *Biophys J* **90**, 318-27.
- Levina, S. and Gordon, R.** (1983). Methionine enkephalin-induced changes in pigmentation of zebrafish (Cyprinidae, *Brachydanio rerio*) and related species and varieties, measured videodensitometrically. I. Zebrafish. *Gen Comp Endocrinol* **51**, 370-7.
- Li, G. and Stahl, P. D.** (1993). Structure-function relationship of the small GTPase rab5. *J Biol Chem* **268**, 24475-80.
- Li, L., Mao, J., Sun, L., Liu, W. and Wu, D.** (2002). Second cysteine-rich domain of Dickkopf-2 activates canonical Wnt signaling pathway via LRP-6 independently of dishevelled. *J Biol Chem* **277**, 5977-81.
- Lin, H. Y., Wang, X. F., Ng-Eaton, E., Weinberg, R. A. and Lodish, H. F.** (1992). Expression cloning of the TGF-beta type II receptor, a functional transmembrane serine/threonine kinase. *Cell* **68**, 775-85.
- Lin, K., Wang, S., Julius, M. A., Kitajewski, J., Moos, M., Jr. and Luyten, F. P.** (1997). The cysteine-rich frizzled domain of Frzb-1 is required and sufficient for modulation of Wnt signaling. *Proc Natl Acad Sci U S A* **94**, 11196-200.

- Lippe, R., Horiuchi, H., Runge, A. and Zerial, M.** (2001a). Expression, purification, and characterization of Rab5 effector complex, rabaptin-5/rabex-5. *Methods Enzymol* **329**, 132-45.
- Lippe, R., Miaczynska, M., Rybin, V., Runge, A. and Zerial, M.** (2001b). Functional synergy between Rab5 effector Rabaptin-5 and exchange factor Rabex-5 when physically associated in a complex. *Mol Biol Cell* **12**, 2219-28.
- Lock, J. G. and Stow, J. L.** (2005). Rab11 in recycling endosomes regulates the sorting and basolateral transport of E-cadherin. *Mol Biol Cell* **16**, 1744-55.
- Loftus, S. K., Larson, D. M., Baxter, L. L., Antonellis, A., Chen, Y., Wu, X., Jiang, Y., Bittner, M., Hammer, J. A., 3rd and Pavan, W. J.** (2002). Mutation of melanosome protein RAB38 in chocolate mice. *Proc Natl Acad Sci U S A* **99**, 4471-6.
- Logan, D. W., Burn, S. F. and Jackson, I. J.** (2006). Regulation of pigmentation in zebrafish melanophores. *Pigment Cell Res* **19**, 206-13.
- Long, S., Ahmad, N. and Rebagliati, M.** (2003). The zebrafish nodal-related gene southpaw is required for visceral and diencephalic left-right asymmetry. *Development* **130**, 2303-16.
- Lopez-Schier, H. and Hudspeth, A. J.** (2005). Supernumerary neuromasts in the posterior lateral line of zebrafish lacking peripheral glia. *Proc Natl Acad Sci U S A* **102**, 1496-501.
- Lutcke, A., Jansson, S., Parton, R. G., Chavrier, P., Valencia, A., Huber, L. A., Lehtonen, E. and Zerial, M.** (1993). Rab17, a novel small GTPase, is specific for epithelial cells and is induced during cell polarization. *J Cell Biol* **121**, 553-64.
- Logan, D. W., Burn, S. F., and Jackson I. J.** (2006). Regulation of pigmentation in zebrafish melanophores. *Pigment Cell Res* **19**, 206-13.
- Lutcke, A., Parton, R. G., Murphy, C., Olkkonen, V. M., Dupree, P., Valencia, A., Simons, K. and Zerial, M.** (1994). Cloning and subcellular localization of novel rab proteins reveals polarized and cell type-specific expression. *J Cell Sci* **107** (Pt 12), 3437-48.
- Maciver, S.** Myosin V, vol. 2006 (ed).
- Mann, R. K. and Beachy, P. A.** (2000). Cholesterol modification of proteins. *Biochim Biophys Acta* **1529**, 188-202.
- Marois, E., Mahmoud, A. and Eaton, S.** (2006). The endocytic pathway and formation of the Wingless morphogen gradient. *Development* **133**, 307-17.
- Martin, E.** (1998). Concise Colour Medical Dictionary, (ed. Oxford: Oxford University Press.
- Martyn, U. and Schulte-Merker, S.** (2003). The ventralized ogon mutant phenotype is caused by a mutation in the zebrafish homologue of Sizzled, a secreted Frizzled-related protein. *Dev Biol* **260**, 58-67.
- Matesic, L. E., Yip, R., Reuss, A. E., Swing, D. A., O'Sullivan, T. N., Fletcher, C. F., Copeland, N. G. and Jenkins, N. A.** (2001). Mutations in Mlph, encoding a member of the Rab effector family, cause the melanosome transport defects observed in leaden mice. *Proc Natl Acad Sci U S A* **98**, 10238-43.
- Mathieu, J., Griffin, K., Herbomel, P., Dickmeis, T., Strahle, U., Kimelman, D., Rosa, F. M. and Peyrieras, N.** (2004). Nodal and Fgf pathways interact through a positive regulatory loop and synergize to maintain mesodermal cell populations. *Development* **131**, 629-41.
- May, S. R., Ashique, A. M., Karlen, M., Wang, B., Shen, Y., Zarbalis, K., Reiter, J., Ericson, J. and Peterson, A. S.** (2005). Loss of the retrograde motor for IFT

- disrupts localization of Smo to cilia and prevents the expression of both activator and repressor functions of Gli. *Dev Biol* **287**, 378-89.
- McBride, H. M., Rybin, V., Murphy, C., Giner, A., Teasdale, R. and Zerial, M.** (1999). Oligomeric complexes link Rab5 effectors with NSF and drive membrane fusion via interactions between EEA1 and syntaxin 13. *Cell* **98**, 377-86.
- McDowell, N., Gurdon, J. B. and Grainger, D. J.** (2001). Formation of a functional morphogen gradient by a passive process in tissue from the early Xenopus embryo. *Int J Dev Biol* **45**, 199-207.
- McFarland, K. N., Warga, R. M. and Kane, D. A.** (2005). Genetic locus half baked is necessary for morphogenesis of the ectoderm. *Dev Dyn* **233**, 390-406.
- McLauchlan, H., Newell, J., Morrice, N., Osborne, A., West, M. and Smythe, E.** (1998). A novel role for Rab5-GDI in ligand sequestration into clathrin-coated pits. *Curr Biol* **8**, 34-45.
- Melby, A. E., Beach, C., Mullins, M. and Kimelman, D.** (2000). Patterning the early zebrafish by the opposing actions of bozozok and vox/vent. *Dev Biol* **224**, 275-85.
- Melton, D. A.** (1987). Translocation of a localized maternal mRNA to the vegetal pole of Xenopus oocytes. *Nature* **328**, 80-2.
- Menasche, G., Ho, C. H., Sanal, O., Feldmann, J., Tezcan, I., Ersoy, F., Houdusse, A., Fischer, A. and de Saint Basile, G.** (2003). Griscelli syndrome restricted to hypopigmentation results from a melanophilin defect (GS3) or a MYO5A F-exon deletion (GS1). *J Clin Invest* **112**, 450-6.
- Menasche, G., Pastural, E., Feldmann, J., Certain, S., Ersoy, F., Dupuis, S., Wulffraat, N., Bianchi, D., Fischer, A., Le Deist, F. et al.** (2000). Mutations in RAB27A cause Griscelli syndrome associated with haemophagocytic syndrome. *Nat Genet* **25**, 173-6.
- Mercer, J. A., Seperack, P. K., Strobel, M. C., Copeland, N. G. and Jenkins, N. A.** (1991). Novel myosin heavy chain encoded by murine dilute coat colour locus. *Nature* **349**, 709-13.
- Milburn, M. V., Tong, L., deVos, A. M., Brunger, A., Yamaizumi, Z., Nishimura, S. and Kim, S. H.** (1990). Molecular switch for signal transduction: structural differences between active and inactive forms of protooncogenic ras proteins. *Science* **247**, 939-45.
- Miller-Bertoglio, V., Carmany-Rampey, A., Furthauer, M., Gonzalez, E. M., Thisse, C., Thisse, B., Halpern, M. E. and Solnica-Krezel, L.** (1999). Maternal and zygotic activity of the zebrafish ogo locus antagonizes BMP signaling. *Dev Biol* **214**, 72-86.
- Milos, N., Dingle, A. D. and Milos, J. P.** (1983). Dynamics of pigment pattern formation in the zebrafish, *Brachydanio rerio*. III. Effect of anteroposterior location of three-day lateral line melanophores on colonization by the second wave of melanophores. *J Exp Zool* **227**, 81-92.
- Mintzer, K. A., Lee, M. A., Runke, G., Trout, J., Whitman, M. and Mullins, M. C.** (2001). Lost-a-fin encodes a type I BMP receptor, Alk8, acting maternally and zygotically in dorsoventral pattern formation. *Development* **128**, 859-69.
- Mizuno, T., Yamaha, M., Wakahara, A., Kuroiwa, A. and Takeda, H.** (1996). Mesoderm induction in zebrafish. *Nature* **383**, 131-132.
- Montero, J. A., Carvalho, L., Wilsch-Brauninger, M., Kilian, B., Mustafa, C. and Heisenberg, C. P.** (2005). Shield formation at the onset of zebrafish gastrulation. *Development* **132**, 1187-98.

- Moore, I., Schell, J. and Palme, K.** (1995). Subclass-specific sequence motifs identified in Rab GTPases. *Trends Biochem Sci* **20**, 10-2.
- Morgan, A. and Burgoine, R. D.** (1995). Is NSF a fusion protein? *Trends Cell Biol* **5**, 335-9.
- Moskowitz, H. S., Heuser, J., McGraw, T. E. and Ryan, T. A.** (2003). Targeted chemical disruption of clathrin function in living cells. *Mol Biol Cell* **14**, 4437-47.
- Mruk, D. D., Lau, A. S. and Conway, A. M.** (2005). Crosstalk between Rab GTPases and cell junctions. *Contraception* **72**, 280-90.
- Murone, M., Rosenthal, A. and de Sauvage, F. J.** (1999). Sonic hedgehog signaling by the patched-smoothened receptor complex. *Curr Biol* **9**, 76-84.
- Nagashima, K., Torii, S., Yi, Z., Igarashi, M., Okamoto, K., Takeuchi, T. and Izumi, T.** (2002). Melanophilin directly links Rab27a and myosin Va through its distinct coiled-coil regions. *FEBS Lett* **517**, 233-8.
- Nakafuku, M., Nagamine, M., Ohtoshi, A., Tanaka, K., Toh-e, A. and Kaziro, Y.** (1993). Suppression of oncogenic Ras by mutant neurofibromatosis type 1 genes with single amino acid substitutions. *Proc Natl Acad Sci U S A* **90**, 6706-10.
- Nakagawa, S. and Takeichi, M.** (1995). Neural crest cell-cell adhesion controlled by sequential and subpopulation-specific expression of novel cadherins. *Development* **121**, 1321-32.
- Nakagawa, S. and Takeichi, M.** (1998). Neural crest emigration from the neural tube depends on regulated cadherin expression. *Development* **125**, 2963-71.
- Nakao, A., Afrakhte, M., Moren, A., Nakayama, T., Christian, J. L., Heuchel, R., Itoh, S., Kawabata, M., Heldin, N. E., Heldin, C. H. et al.** (1997). Identification of Smad7, a TGFbeta-inducible antagonist of TGF-beta signalling. *Nature* **389**, 631-5.
- Nakato, H., Futch, T. A. and Selleck, S. B.** (1995). The division abnormally delayed (dally) gene: a putative integral membrane proteoglycan required for cell division patterning during postembryonic development of the nervous system in Drosophila. *Development* **121**, 3687-702.
- Nakayama, T., Snyder, M. A., Grewal, S. S., Tsuneizumi, K., Tabata, T. and Christian, J. L.** (1998). Xenopus Smad8 acts downstream of BMP-4 to modulate its activity during vertebrate embryonic patterning. *Development* **125**, 857-67.
- Nasevicius, A. and Ekker, S.** (2000). Effective targeted gene 'knockdown' in zebrafish. *Nature Genetics* **26**, 216-220.
- Nelson, R. W. and Gumbiner, B. M.** (1998). Beta-catenin directly induces expression of the Siamois gene, and can initiate signaling indirectly via a membrane-tethered form. *Ann NY Acad Sci* **857**, 86-98.
- Nicolson, T.** (2005). The genetics of hearing and balance in zebrafish. *Annu Rev Genet* **39**, 9-22.
- Nicolson, T., Rusch, A., Friedrich, R. W., Granato, M., Ruppertsberg, J. P. and Nusslein-Volhard, C.** (1998). Genetic analysis of vertebrate sensory hair cell mechanosensation: the zebrafish circler mutants. *Neuron* **20**, 271-83.
- Nielsen, E., Christoforidis, S., Uttenweiler-Joseph, S., Miaczynska, M., Dewitte, F., Wilim, M., Hoflack, B. and Zerial, M.** (2000). Rabenosyn-5, a novel Rab5 effector, is complexed with hVPS45 and recruited to endosomes through a FYVE finger domain. *J Cell Biol* **151**, 601-12.
- Nielsen, E., Severin, F., Backer, J. M., Hyman, A. A. and Zerial, M.** (1999). Rab5 regulates motility of early endosomes on microtubules. *Nat Cell Biol* **1**, 376-82.
- Nojima, H., Shimizu, T., Kim, C. H., Yabe, T., Bae, Y. K., Muraoka, O., Hirata, T., Chitnis, A., Hirano, T. and Hibi, M.** (2004). Genetic evidence for involvement

- of maternally derived Wnt canonical signaling in dorsal determination in zebrafish. *Mech Dev* **121**, 371-86.
- Nonet, M. L., Staunton, J. E., Kilgard, M. P., Fergestad, T., Hartwieg, E., Horvitz, H. R., Jorgensen, E. M. and Meyer, B. J.** (1997). Caenorhabditis elegans rab-3 mutant synapses exhibit impaired function and are partially depleted of vesicles. *J Neurosci* **17**, 8061-73.
- Nuoffer, C., Davidson, H. W., Matteson, J., Meinkoth, J. and Balch, W. E.** (1994). A GDP-bound of rab1 inhibits protein export from the endoplasmic reticulum and transport between Golgi compartments. *J Cell Biol* **125**, 225-37.
- Oates, A. C., Pratt, S. J., Vail, B., Yan, Y., Ho, R. K., Johnson, S. L., Postlethwait, J. H. and Zon, L. I.** (2001). The zebrafish klf gene family. *Blood* **98**, 1792-801.
- Odenthal, J., Rossnagel, K., Haffter, P., Kelsh, R. N., Vogelsang, E., Brand, M., van Eeden, F. J., Furutani-Seiki, M., Granato, M., Hammerschmidt, M. et al.** (1996). Mutations affecting xanthophore pigmentation in the zebrafish, Danio rerio. *Development* **123**, 391-8.
- Onichtchouk, D., Chen, Y. G., Dosch, R., Gawantka, V., Delius, H., Massague, J. and Niehrs, C.** (1999). Silencing of TGF- β signalling by the pseudoreceptor BAMBI. *Nature* **401**, 480-5.
- Ono, F., Shcherbatko, A., Higashijima, S., Mandel, G. and Brehm, P.** (2002). The Zebrafish motility mutant twitch once reveals new roles for rapsyn in synaptic function. *J Neurosci* **22**, 6491-8.
- Oppenheimer, J.** (1983). Thomas Hunt Morgan as an embryologist: The view from bryn mawr. *Amer. Zool* **23**, 845-854.
- Osanai, K., Takahashi, K., Nakamura, K., Takahashi, M., Ishigaki, M., Sakuma, T., Toga, H., Suzuki, T. and Voelker, D. R.** (2005). Expression and characterization of Rab38, a new member of the Rab small G protein family. *Biol Chem* **386**, 143-53.
- Ossig, R., Laufer, W., Schmitt, H. D. and Gallwitz, D.** (1995). Functionality and specific membrane localization of transport GTPases carrying C-terminal membrane anchors of synaptobrevin-like proteins. *Embo J* **14**, 3645-53.
- Ostermeier, C. and Brunger, A. T.** (1999) Structural basis of Rab effector specificity: crystal structure of the small G protein Rab3A complexed with the effector domain of rabphilin-3A. *Cell* **96**, 363-74
- Oyler, G. A., Higgins, G. A., Hart, R. A., Battenberg, E., Billingsley, M., Bloom, F. E. and Wilson, M. C.** (1989). The identification of a novel synaptosomal-associated protein, SNAP-25, differentially expressed by neuronal subpopulations. *J Cell Biol* **109**, 3039-52.
- Palokangas, H., Ying, M., Vaananen, K. and Saraste, J.** (1998). Retrograde transport from the pre-Golgi intermediate compartment and the Golgi complex is affected by the vacuolar H⁺-ATPase inhibitor bafilomycin A1. *Mol Biol Cell* **9**, 3561-78.
- Pan, J. Y. and Wessling-Resnick, M.** (1998). GEF-mediated GDP/GTP exchange by monomeric GTPases: a regulatory role for Mg²⁺? *Bioessays* **20**, 516-21.
- Papkoff, J., Rubinfeld, B., Schryver, B. and Polakis, P.** (1996). Wnt-1 regulates free pools of catenins and stabilizes APC-catenin complexes. *Mol Cell Biol* **16**, 2128-34.
- Parichy, D. M.** (2003). Pigment patterns: fish in stripes and spots. *Curr Biol* **13**, R947-50.

- Parichy, D. M., Rawls, J. F., Pratt, S. J., Whitfield, T. T. and Johnson, S. L.** (1999). Zebrafish sparse corresponds to an orthologue of c-kit and is required for the morphogenesis of a subpopulation of melanocytes, but is not essential for hematopoiesis or primordial germ cell development. *Development* **126**, 3425-36.
- Parsons, M., Pollard, S., Saude, L., Feldman, B., Coutinho, P., Hirst, E. and Stemple, D.** (2002a). Zebrafish mutants identify an essential role for laminins in notochord formation. *Development* **129**, 3137-3146.
- Parsons, M. J., Campos, I., Hirst, E. M. and Stemple, D. L.** (2002b). Removal of dystroglycan causes severe muscular dystrophy in zebrafish embryos. *Development* **129**, 3505-12.
- Pastural, E., Barrat, F. J., Dufourcq-Lagelouse, R., Certain, S., Sanal, O., Jabado, N., Seger, R., Griscelli, C., Fischer, A. and de Saint Basile, G.** (1997). Griscelli disease maps to chromosome 15q21 and is associated with mutations in the myosin-Va gene. *Nat Genet* **16**, 289-92.
- Pastural, E., Ersoy, F., Yalman, N., Wulffraat, N., Grillo, E., Ozkinay, F., Tezcan, I., Gedikoglu, G., Philippe, N., Fischer, A. et al.** (2000). Two genes are responsible for Griscelli syndrome at the same 15q21 locus. *Genomics* **63**, 299-306.
- Pavlos, N., Xu, J., Papadimitriou, J. and Zheng, M.** (2001). Molecular cloning of the mouse homologue of Rab3c. *Journal of Molecular Endocrinology* **27**, 117-122.
- Pelkmans, L., Fava, E., Grabner, H., Hannus, M., Habermann, B., Krausz, E. and Zerial, M.** (2005). Genome-wide analysis of human kinases in clathrin- and caveolae/raft-mediated endocytosis. *Nature* **436**, 78-86.
- Pellinen, T., Arjonen, A., Vuoriluoto, K., Kallio, K., Fransen, J. A. and Ivaska, J.** (2006). Small GTPase Rab21 regulates cell adhesion and controls endosomal traffic of beta1-integrins. *J Cell Biol* **173**, 767-80.
- Pepinsky, R. B., Zeng, C., Wen, D., Rayhorn, P., Baker, D. P., Williams, K. P., Bixler, S. A., Ambrose, C. M., Garber, E. A., Miatkowski, K. et al.** (1998). Identification of a palmitic acid-modified form of human Sonic hedgehog. *J Biol Chem* **273**, 14037-45.
- Pereira-Leal, J. and Seabra, M.** (2000). The Mammalian Rab Family of Small GTPases: Definition of Family and SubfamilySequence Motifs Suggests a Mechanism for Functional Specificity in the Ras Superfamily. *Journal of Molecular Biology* **301**, 1077-1087.
- Piccolo, S., Agius, E., Leyns, L., Bhattacharyya, S., Grunz, H., Bouwmeester, T. and De Robertis, E. M.** (1999). The head inducer Cerberus is a multifunctional antagonist of Nodal, BMP and Wnt signals. *Nature* **397**, 707-10.
- Piccolo, S., Sasai, Y., Lu, B. and De Robertis, E. M.** (1996). Dorsoventral patterning in Xenopus: inhibition of ventral signals by direct binding of chordin to BMP-4. *Cell* **86**, 589-98.
- Piek, E., Heldin, C. H. and Ten Dijke, P.** (1999). Specificity, diversity, and regulation in TGF-beta superfamily signaling. *Faseb J* **13**, 2105-24.
- Plutner, H., Cox, A. D., Pind, S., Khosravi-Far, R., Bourne, J. R., Schwaninger, R., Der, C. J. and Balch, W. E.** (1991). Rab1b regulates vesicular transport between the endoplasmic reticulum and successive Golgi compartments. *J Cell Biol* **115**, 31-43.
- Polo, S. and Di Fiore, P. P.** (2006). Endocytosis conducts the cell signaling orchestra. *Cell* **124**, 897-900.
- Poulain, M., Furthauer, M., Thisse, B., Thisse, C. and Lepage, T.** (2006). Zebrafish endoderm formation is regulated by combinatorial Nodal, FGF and BMP signalling. *Development* **133**, 2189-200.

- Price, A., Seals, D., Wickner, W. and Ungermann, C.** (2000a). The docking stage of yeast vacuole fusion requires the transfer of proteins from a cis-SNARE complex to a Rab/Ypt protein. *J Cell Biol* **148**, 1231-8.
- Price, A., Wickner, W. and Ungermann, C.** (2000b). Proteins needed for vesicle budding from the Golgi complex are also required for the docking step of homotypic vacuole fusion. *J Cell Biol* **148**, 1223-29.
- Pruyne, D. W., Schott, D. H. and Bretscher, A.** (1998). Tropomyosin-containing actin cables direct the Myo2p-dependent polarized delivery of secretory vesicles in budding yeast. *J Cell Biol* **143**, 1931-45.
- Pyati, U. J., Webb, A. E. and Kimelman, D.** (2005). Transgenic zebrafish reveal stage-specific roles for Bmp signaling in ventral and posterior mesoderm development. *Development* **132**, 2333-43.
- Raible, D. W., Wood, A., Hodsdon, W., Henion, P. D., Weston, J. A. and Eisen, J. S.** (1992). Segregation and early dispersal of neural crest cells in the embryonic zebrafish. *Dev Dyn* **195**, 29-42.
- Ramel, M. C., Buckles, G. R., Baker, K. D. and Lekven, A. C.** (2005). WNT8 and BMP2B co-regulate non-axial mesoderm patterning during zebrafish gastrulation. *Dev Biol* **287**, 237-48.
- Ramel, M. C. and Lekven, A. C.** (2004). Repression of the vertebrate organizer by Wnt8 is mediated by Vent and Vox. *Development* **131**, 3991-4000.
- Ramirez-Weber, F. A. and Kornberg, T. B.** (1999). Cytonemes: cellular processes that project to the principal signaling center in *Drosophila* imaginal discs. *Cell* **97**, 599-607.
- Rebagliati, M., Toyama, R., Haffter, P. and Dawida, I.** (1998). *cyclops* encodes a noda-related factor involved in midline signaling. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 9932-9937.
- Rebagliati, M., Toyama, R., Fricke, C., Haffter, P. and Dawida, I.** (1998). Zebrafish Nodal-Related Genes Are Implicated in Axial Patterning and Establishing Left-Right Asymmetry. *Developmental Biology* **199**, 261-272.
- Reifers, F., Bohli, H., Walsh, E. C., Crossley, P. H., Stainier, D. Y. and Brand, M.** (1998). Fgf8 is mutated in zebrafish acerebellar (ace) mutants and is required for maintenance of midbrain-hindbrain boundary development and somitogenesis. *Development* **125**, 2381-95.
- Reissmann, E., Jornvall, H., Blokzijl, A., Andersson, O., Chang, C., Minchiotti, G., Persico, M. G., Ibáñez, C. F. and Brivanlou, A. H.** (2001). The orphan receptor ALK7 and the Activin receptor ALK4 mediate signaling by Nodal proteins during vertebrate development. *Genes Dev* **15**, 2010-22.
- Rieder, S. E. and Emr, S. D.** (1997). A novel RING finger protein complex essential for a late step in protein transport to the yeast vacuole. *Mol Biol Cell* **8**, 2307-27.
- Riederer, M. A., Soldati, T., Shapiro, A. D., Lin, J. and Pfeffer, S. R.** (1994). Lysosome biogenesis requires Rab9 function and receptor recycling from endosomes to the trans-Golgi network. *J Cell Biol* **125**, 573-82.
- Rietveld, A., Neutz, S., Simons, K. and Eaton, S.** (1999). Association of sterol- and glycosylphosphatidylinositol-linked proteins with *Drosophila* raft lipid microdomains. *J Biol Chem* **274**, 12049-54.
- Rodionov, V. I., Hope, A. J., Svitkina, T. M. and Borisy, G. G.** (1998). Functional coordination of microtubule-based and actin-based motility in melanophores. *Curr Biol* **8**, 165-8.

- Rogers, S. L. and Gelfand, V. I.** (1998). Myosin cooperates with microtubule motors during organelle transport in melanophores. *Curr Biol* **8**, 161-4.
- Rothman, J. E. and Warren, G.** (1994). Implications of the SNARE hypothesis for intracellular membrane topology and dynamics. *Curr Biol* **4**, 220-33.
- Rybin, V., Ullrich, O., Rubino, M., Alexandrov, K., Simon, I., Seabra, M. C., Goody, R. and Zerial, M.** (1996). GTPase activity of Rab5 acts as a timer for endocytic membrane fusion. *Nature* **383**, 266-9.
- Sacher, M., Jiang, Y., Barrowman, J., Scarpa, A., Burston, J., Zhang, L., Schieltz, D., Yates, J. R., 3rd, Abeliovich, H. and Ferro-Novick, S.** (1998). TRAPP, a highly conserved novel complex on the cis-Golgi that mediates vesicle docking and fusion. *Embo J* **17**, 2494-503.
- Sakuma, R., Ohnishi Yi, Y., Meno, C., Fujii, H., Juan, H., Takeuchi, J., Ogura, T., Li, E., Miyazono, K. and Hamada, H.** (2002). Inhibition of Nodal signalling by Lefty mediated through interaction with common receptors and efficient diffusion. *Genes Cells* **7**, 401-12.
- Salminen, A. and Novick, P. J.** (1987). A ras-like protein is required for a post-Golgi event in yeast secretion. *Cell* **49**, 527-38.
- Sampath, K., Rubinstein, A., Cheng, A., Liang, J., Fekany, K., Solnica-Krezel, L., Korzh, V., Halpern, M. and Wright, C.** (1998). Induction of the zebrafish ventral brain and floorplate requires cyclops/nodal signalling. *Nature* **395**, 185-189.
- Sannerud, R., Saraste, J. and Goud, B.** (2003). Retrograde traffic in the biosynthetic-secretory route: pathways and machinery. *Curr Opin Cell Biol* **15**, 438-45.
- Sanson, B., Alexandre, C., Fascetti, N. and Vincent, J. P.** (1999). Engrailed and hedgehog make the range of Wingless asymmetric in Drosophila embryos. *Cell* **98**, 207-16.
- Sasaki, T., Kaibuchi, K., Kabcenell, A. K., Novick, P. J. and Takai, Y.** (1991). A mammalian inhibitory GDP/GTP exchange protein (GDP dissociation inhibitor) for smg p25A is active on the yeast SEC4 protein. *Mol Cell Biol* **11**, 2909-12.
- Schier, A. and Talbot, W.** (2001). Nodal signaling and the zebrafish organizer. *Int.J.Dev.Biol* **45**, 289-297.
- Schier, A. F. and Talbot, W. S.** (2005). Molecular Genetics of Axis Formation in Zebrafish. *Annu Rev Genet*.
- Schimmoller, F., Simon, I. and Pfeffer, S. R.** (1998). Rab GTPases, directors of vesicle docking. *J Biol Chem* **273**, 22161-4.
- Schmid, B., Furthauer, M., Connors, S. A., Trout, J., Thisse, B., Thisse, C. and Mullins, M. C.** (2000). Equivalent genetic roles for bmp7/snailhouse and bmp2b/swirl in dorsoventral pattern formation. *Development* **127**, 957-67.
- Schmidt, A. and Hall, A.** (2002). Guanine nucleotide exchange factors for Rho GTPases: turning on the switch. *Genes Dev* **16**, 1587-609.
- Schmitt, H. D., Puzicha, M. and Gallwitz, D.** (1988). Study of a temperature-sensitive mutant of the ras-related YPT1 gene product in yeast suggests a role in the regulation of intracellular calcium. *Cell* **53**, 635-47.
- Schmitt, H. D., Wagner, P., Pfaff, E. and Gallwitz, D.** (1986). The ras-related YPT1 gene product in yeast: a GTP-binding protein that might be involved in microtubule organization. *Cell* **47**, 401-12.
- Schneider, S., Steinbeisser, H., Warga, R. M. and Hausen, P.** (1996). Beta-catenin translocation into nuclei demarcates the dorsalizing centers in frog and fish embryos. *Mech Dev* **57**, 191-8.

- Scholpp, S. and Brand, M.** (2004). Endocytosis controls spreading and effective signaling range of Fgf8 protein. *Curr Biol* **14**, 1834-41.
- Schott, D., Ho, J., Pruyne, D. and Bretscher, A.** (1999). The COOH-terminal domain of Myo2p, a yeast myosin V, has a direct role in secretory vesicle targeting. *J Cell Biol* **147**, 791-808.
- Schulte-Merker, S., Lee, K. J., McMahon, A. P. and Hammerschmidt, M.** (1997). The zebrafish organizer requires chordino. *Nature* **387**, 862-3.
- Scott, M.** (2001). Signalling and endocytosis: Wnt breaks down on back roads. *Nat Cell Biol* **3**, E185-6.
- Seabra, M., Mules, E. and Hume, A.** (2002). Rab GTPases, intracellular traffic and disease. *Trends In Molecular Medicine* **8**, 23-26.
- Seabra, M. C. and Wasmeier, C.** (2004). Controlling the location and activation of Rab GTPases. *Curr Opin Cell Biol* **16**, 451-7.
- Seals, D. F., Eitzen, G., Margolis, N., Wickner, W. T. and Price, A.** (2000). A Ypt/Rab effector complex containing the Sec1 homolog Vps33p is required for homotypic vacuole fusion. *Proc Natl Acad Sci U S A* **97**, 9402-7.
- Sepich, D. S., Myers, D. C., Short, R., Topczewski, J., Marlow, F. and Solnica-Krezel, L.** (2000). Role of the zebrafish trilobite locus in gastrulation movements of convergence and extension. *Genesis* **27**, 159-73.
- Seto, E. S. and Bellen, H. J.** (2006). Internalization is required for proper Wingless signaling in *Drosophila melanogaster*. *J Cell Biol* **173**, 95-106.
- Settleman, J., Albright, C. F., Foster, L. C. and Weinberg, R. A.** (1992). Association between GTPase activators for Rho and Ras families. *Nature* **359**, 153-4.
- Sheldahl, L. C., Park, M., Malbon, C. C. and Moon, R. T.** (1999). Protein kinase C is differentially stimulated by Wnt and Frizzled homologs in a G-protein-dependent manner. *Curr Biol* **9**, 695-8.
- Shimizu, T., Bae, Y. K., Muraoka, O. and Hibi, M.** (2005a). Interaction of Wnt and caudal-related genes in zebrafish posterior body formation. *Dev Biol* **279**, 125-41.
- Shimizu, T., Yabe, T., Muraoka, O., Yonemura, S., Aramaki, S., Hatta, K., Bae, Y. K., Nojima, H. and Hibi, M.** (2005b). E-cadherin is required for gastrulation cell movements in zebrafish. *Mech Dev* **122**, 747-63.
- Shimizu, T., Yamanaka, Y., Nojima, H., Yabe, T., Hibi, M. and Hirano, T.** (2002). A novel repressor-type homeobox gene, ved, is involved in dharma/bozozok-mediated dorsal organizer formation in zebrafish. *Mech Dev* **118**, 125-38.
- Singer-Kruger, B., Stenmark, H., Dusterhoff, A., Philippse, P., Yoo, J. S., Gallwitz, D. and Zerial, M.** (1994). Role of three rab5-like GTPases, Ypt51p, Ypt52p, and Ypt53p, in the endocytic and vacuolar protein sorting pathways of yeast. *J Cell Biol* **125**, 283-98.
- Singer-Kruger, B., Stenmark, H. and Zerial, M.** (1995). Yeast Ypt51p and mammalian Rab5: counterparts with similar function in the early endocytic pathway *J Cell Sci* **108**, 3509-3521.
- Sivars, U., Aivazian, D. and Pfeffer, S. R.** (2003). Yip3 catalyses the dissociation of endosomal Rab-GDI complexes. *Nature* **425**, 856-9.
- Skold, H. N., Norstrom, E. and Wallin, M.** (2002). Regulatory control of both microtubule- and actin-dependent fish melanosome movement. *Pigment Cell Res* **15**, 357-66.
- Slack, J. M.** (1994). Developmental biology. How to make the gradient. *Nature* **371**, 477-8.

- Smith, W. C. and Harland, R. M.** (1991). Injected Xwnt-8 RNA acts early in Xenopus embryos to promote formation of a vegetal dorsalizing center. *Cell* **67**, 753-65.
- Sollner, C., Burghammer, M., Busch-Nentwich, E., Berger, J., Schwarz, H., Riekel, C. and Nicolson, T.** (2003). Control of crystal size and lattice formation by starmaker in otolith biomineralization. *Science* **302**, 282-6.
- Sollner, T., Whiteheart, S. W., Brunner, M., Erdjument-Bromage, H., Geromanos, S., Tempst, P. and Rothman, J. E.** (1993). SNAP receptors implicated in vesicle targeting and fusion. *Nature* **362**, 318-24.
- Solnica-Krezel, L. and Driever, W.** (1994). Microtubule arrays of the zebrafish yolk cell: organization and function during epiboly. *Development* **120**, 2443-55.
- Solnica-Krezel, L. and Driever, W.** (2001). The role of the homeodomain protein Bozozok in zebrafish axis formation. *Int J Dev Biol* **45**, 299-310.
- Solnica-Krezel, L., Stemple, D. L., Mountcastle-Shah, E., Rangini, Z., Neuhauss, S. C., Malicki, J., Schier, A. F., Stainier, D. Y., Zwartkruis, F., Abdelilah, S. et al.** (1996). Mutations affecting cell fates and cellular rearrangements during gastrulation in zebrafish. *Development* **123**, 67-80.
- Southard-Smith, E. M., Kos, L. and Pavan, W. J.** (1998). Sox10 mutation disrupts neural crest development in Dom Hirschsprung mouse model. *Nat Genet* **18**, 60-4.
- Spemann, H. and Mangold, H.** (1924). On the induction of embryonic structures by implantation of organizers of different species. *Wilhelm Roux Arch Entw Mech* **100**, 599-638.
- Starr, C. J., Kappler, J. A., Chan, D. K., Kollmar, R. and Hudspeth, A. J.** (2004). Mutation of the zebrafish choroideremia gene encoding Rab escort protein 1 devastates hair cells. *Proc Natl Acad Sci U S A* **101**, 2572-7.
- Stenmark, H. and Aasland, R.** (1999). FYVE-finger proteins--effectors of an inositol lipid. *J Cell Sci* **112** (Pt 23), 4175-83.
- Stenmark, H., Aasland, R., Toh, B. H. and D'Arrigo, A.** (1996). Endosomal localization of the autoantigen EEA1 is mediated by a zinc-binding FYVE finger. *J Biol Chem* **271**, 24048-54.
- Stenmark, H., Valencia, A., Martinez, O., Ullrich, O., Goud, B. and Zerial, M.** (1994). Distinct structural elements of rab5 define its functional specificity. *Embo J* **13**, 575-83.
- Stenmark, H., Vitale, G., Ullrich, O. and Zerial, M.** (1995). Rabaptin-5 is a direct effector of the small GTPase Rab5 in endocytic membrane fusion. *Cell* **83**, 423-32.
- Strigini, M. and Cohen, S. M.** (2000). Wingless gradient formation in the Drosophila wing. *Curr Biol* **10**, 293-300.
- Sun, Z., Jin, P., Tian, T., Gu, Y., Chen, Y. G. and Meng, A.** (2006). Activation and roles of ALK4/ALK7-mediated maternal TGFbeta signals in zebrafish embryo. *Biochem Biophys Res Commun* **345**, 694-703.
- Suzuki, A., Thies, R. S., Yamaji, N., Song, J. J., Wozney, J. M., Murakami, K. and Ueno, N.** (1994). A truncated bone morphogenetic protein receptor affects dorsal-ventral patterning in the early Xenopus embryo. *Proc Natl Acad Sci U S A* **91**, 10255-9.
- Suzuki, T., Miyamura, Y., Inagaki, K. and Tomita, Y.** (2003). Characterization of the human RAB38 and RAB7 genes: exclusion of new major pathological loci for Japanese OCA. *J Dermatol Sci* **32**, 131-6.
- Takei, K., McPherson, P. S., Schmid, S. L. and De Camilli, P.** (1995). Tubular membrane invaginations coated by dynamin rings are induced by GTP-gamma S in nerve terminals. *Nature* **374**, 186-90.

- Tall, G. G., Hama, H., DeWald, D. B. and Horazdovsky, B. F.** (1999). The phosphatidylinositol 3-phosphate binding protein Vac1p interacts with a Rab GTPase and a Sec1p homologue to facilitate vesicle-mediated vacuolar protein sorting. *Mol Biol Cell* **10**, 1873-89.
- Tao, Q., Yokota, C., Puck, H., Kofron, M., Birsoy, B., Yan, D., Asashima, M., Wylie, C. C., Lin, X. and Heasman, J.** (2005). Maternal wnt11 activates the canonical wnt signaling pathway required for axis formation in Xenopus embryos. *Cell* **120**, 857-71.
- Telegman, A. and Cohen, S.** (2000). Dpp Gradient Formation in the Drosophila Wing Imaginal Disc. *Cell* **103**, 971-980.
- ten Dijke, P., Yamashita, H., Ichijo, H., Franzen, P., Laiho, M., Miyazono, K. and Heldin, C. H.** (1994). Characterization of type I receptors for transforming growth factor-beta and activin. *Science* **264**, 101-4.
- TerBush, D. R., Maurice, T., Roth, D. and Novick, P.** (1996). The Exocyst is a multiprotein complex required for exocytosis in *Saccharomyces cerevisiae*. *Embo J* **15**, 6483-94.
- Thoma, N. H., Iakovenko, A., Goody, R. S. and Alexandrov, K.** (2001a). Phosphoisoprenoids modulate association of Rab geranylgeranyltransferase with REP-1. *J Biol Chem* **276**, 48637-43.
- Thoma, N. H., Iakovenko, A., Kalinin, A., Waldmann, H., Goody, R. S. and Alexandrov, K.** (2001b). Allosteric regulation of substrate binding and product release in geranylgeranyltransferase type II. *Biochemistry* **40**, 268-74.
- Thoma, N. H., Niculae, A., Goody, R. S. and Alexandrov, K.** (2001c). Double prenylation by RabGGTase can proceed without dissociation of the mono-prenylated intermediate. *J Biol Chem* **276**, 48631-6.
- Tickle, C., Summerbell, D. and Wolpert, L.** (1975). Positional signalling and specification of digits in chick limb morphogenesis. *Nature* **254**, 199-202.
- Tisdale, E. J., Bourne, J. R., Khosravi-Far, R., Der, C. J. and Balch, W. E.** (1992). GTP-binding mutants of rab1 and rab2 are potent inhibitors of vesicular transport from the endoplasmic reticulum to the Golgi complex. *J Cell Biol* **119**, 749-61.
- Topczewski, J., Sepich, D. S., Myers, D. C., Walker, C., Amores, A., Lele, Z., Hammerschmidt, M., Postlethwait, J. and Solnica-Krezel, L.** (2001). The zebrafish glypcan knypek controls cell polarity during gastrulation movements of convergent extension. *Dev Cell* **1**, 251-64.
- Touchot, N., Chardin, P. and Tavitian, A.** (1987). Four additional members of the ras gene superfamily isolated by an oligonucleotide strategy: molecular cloning of YPT-related cDNAs from a rat brain library. *Proc Natl Acad Sci U S A* **84**, 8210-4.
- Trimble, W. S., Cowan, D. M. and Scheller, R. H.** (1988). VAMP-1: a synaptic vesicle-associated integral membrane protein. *Proc Natl Acad Sci U S A* **85**, 4538-42.
- Trinh, L. A., Meyer, D. and Stainier, D. Y.** (2003). The Mix family homeodomain gene bonnie and clyde functions with other components of the Nodal signaling pathway to regulate neural patterning in zebrafish. *Development* **130**, 4989-98.
- Trinkaus, J. P.** (1963). The cellular basis of Fundulus epiboly. Adhesivity of blastula and gastrula cells in culture. *Dev Biol* **7**, 513-32.
- Trinkaus, J. P.** (1984). Mechanism of Fundulus epiboly - a current view. *Amer. Zool* **24**, 673-688.
- Tsikolia, N.** (2006). The role and limits of a gradient based explanation of morphogenesis: a theoretical consideration. *Int J Dev Biol* **50**, 333-40.

- Tsukazaki, T., Chiang, T. A., Davison, A. F., Attisano, L. and Wrana, J. L.** (1998). SARA, a FYVE domain protein that recruits Smad2 to the TGFbeta receptor. *Cell* **95**, 779-91.
- Ullrich, O., Reinsch, S., Urbe, S., Zerial, M. and Parton, R. G.** (1996). Rab11 regulates recycling through the pericentriolar recycling endosome. *J Cell Biol* **135**, 913-24.
- Ulrich, F., Concha, M. L., Heid, P. J., Voss, E., Witzel, S., Roehl, H., Tada, M., Wilson, S. W., Adams, R. J., Soll, D. R. et al.** (2003). Slb/Wnt11 controls hypoblast cell migration and morphogenesis at the onset of zebrafish gastrulation. *Development* **130**, 5375-84.
- Ulrich, F., Krieg, M., Schotz, E. M., Link, V., Castanon, I., Schnabel, V., Taubenberger, A., Mueller, D., Puech, P. H. and Heisenberg, C. P.** (2005). Wnt11 functions in gastrulation by controlling cell cohesion through Rab5c and E-cadherin. *Dev Cell* **9**, 555-64.
- Vagner, S., Galy, B. and Pyronnet, S.** (2001). Irresistible IRES. Attracting the translation machinery to internal ribosome entry sites. *EMBO Rep* **2**, 893-8.
- Valencia, A., Chardin, P., Wittinghofer, A. and Sander, C.** (1991). The ras protein family: evolutionary tree and role of conserved amino acids. *Biochemistry* **30**, 4637-48.
- Verhoeven, K., De Jonghe, P., Coen, K., Verpoorten, N., Auer-Grumbach, M., Kwon, J. M., FitzPatrick, D., Schmedding, E., De Vriendt, E., Jacobs, A. et al.** (2003). Mutations in the small GTP-ase late endosomal protein RAB7 cause Charcot-Marie-Tooth type 2B neuropathy. *Am J Hum Genet* **72**, 722-7.
- Wagner, D. S., Dosch, R., Mintzer, K. A., Wiemelt, A. P. and Mullins, M. C.** (2004). Maternal control of development at the midblastula transition and beyond: mutants from the zebrafish II. *Dev Cell* **6**, 781-90.
- Wagner, D. S. and Mullins, M. C.** (2002). Modulation of BMP activity in dorsal-ventral pattern formation by the chordin and ogoen antagonists. *Dev Biol* **245**, 109-23.
- Wallingford, J. B., Rowning, B. A., Vogeli, K. M., Rothbacher, U., Fraser, S. E. and Harland, R. M.** (2000). Dishevelled controls cell polarity during Xenopus gastrulation. *Nature* **405**, 81-5.
- Wang, W., Sacher, M. and Ferro-Novick, S.** (2000). TRAPP stimulates guanine nucleotide exchange on Ypt1p. *J Cell Biol* **151**, 289-96.
- Wang, Y., Ng, E. L. and Tang, B. L.** (2006). Rab23: what exactly does it traffic? *Traffic* **7**, 746-50.
- Warga, R. M. and Kimmel, C. B.** (1990). Cell movements during epiboly and gastrulation in zebrafish. *Development* **108**, 569-80.
- Weigmann, K. and Cohen, S. M.** (1999). Lineage-tracing cells born in different domains along the PD axis of the developing Drosophila leg. *Development* **126**, 3823-30.
- Wennerberg, K., Rossman, K. L. and Der, C. J.** (2005). The Ras superfamily at a glance. *J Cell Sci* **118**, 843-6.
- Westerfield, M.** (2000). The zebrafish book. A guide for the laboratory use of zebrafish (*Danio rerio*). 4th ed., Univ. of Oregon Press, Eugene.
- Whitman, M.** (2001). Nodal Signaling in Early Vertebrate Embryos: Themes and Variations. *Developmental Cell* **1**, 605-617.
- Wichmann, H., Hengst, L. and Gallwitz, D.** (1992). Endocytosis in yeast: evidence for the involvement of a small GTP-binding protein (Ypt7p). *Cell* **71**, 1131-42.

- Wiederkehr, A., De Craene, J. O., Ferro-Novick, S. and Novick, P.** (2004). Functional specialization within a vesicle tethering complex: bypass of a subset of exocyst deletion mutants by Sec1p or Sec4p. *J Cell Biol* **167**, 875-87.
- Williams, P. H., Hagemann, A., Gonzalez-Gaitan, M. and Smith, J. C.** (2004). Visualizing long-range movement of the morphogen Xnr2 in the Xenopus embryo. *Curr Biol* **14**, 1916-23.
- Willot, V., Mathieu, J., Lu, Y., Schmid, B., Sidi, S., Yan, Y. L., Postlethwait, J. H., Mullins, M., Rosa, F. and Peyreras, N.** (2002). Cooperative action of ADMP- and BMP-mediated pathways in regulating cell fates in the zebrafish gastrula. *Dev Biol* **241**, 59-78.
- Wilson, D. B. and Wilson, M. P.** (1992). Identification and subcellular localization of human rab5b, a new member of the ras-related superfamily of GTPases. *J Clin Invest* **89**, 996-1005.
- Wilson, S. M., Yip, R., Swing, D. A., O'Sullivan, T. N., Zhang, Y., Novak, E. K., Swank, R. T., Russell, L. B., Copeland, N. G. and Jenkins, N. A.** (2000). A mutation in Rab27a causes the vesicle transport defects observed in ashen mice. *Proc Natl Acad Sci U S A* **97**, 7933-8.
- Wolpert, L.** (1996). One hundred years of positional information. *Trends Genet* **12**, 359-64.
- Wu, S. K., Zeng, K., Wilson, I. A. and Balch, W. E.** (1996). Structural insights into the function of the Rab GDI superfamily. *Trends Biochem Sci* **21**, 472-6.
- Wu, X., Bowers, B., Rao, K., Wei, Q. and Hammer, J. A., 3rd.** (1998). Visualization of melanosome dynamics within wild-type and dilute melanocytes suggests a paradigm for myosin V function In vivo. *J Cell Biol* **143**, 1899-918.
- Wu, X., Bowers, B., Wei, Q., Kocher, B. and Hammer, J. A., 3rd.** (1997). Myosin V associates with melanosomes in mouse melanocytes: evidence that myosin V is an organelle motor. *J Cell Sci* **110** (Pt 7), 847-59.
- Wylie, C., Kofron, M., Payne, C., Anderson, R., Hosobuchi, M., Joseph, E. and Heasman, J.** (1996). Maternal beta-catenin establishes a 'dorsal signal' in early Xenopus embryos. *Development* **122**, 2987-96.
- Xu, X. Z., Wes, P. D., Chen, H., Li, H. S., Yu, M., Morgan, S., Liu, Y. and Montell, C.** (1998). Retinal targets for calmodulin include proteins implicated in synaptic transmission. *J Biol Chem* **273**, 31297-307.
- Yabe, T., Shimizu, T., Muraoka, O., Bae, Y. K., Hirata, T., Nojima, H., Kawakami, A., Hirano, T. and Hibi, M.** (2003). Ogon/Secreted Frizzled functions as a negative feedback regulator of Bmp signaling. *Development* **130**, 2705-16.
- Yang-Snyder, J., Miller, J. R., Brown, J. D., Lai, C. J. and Moon, R. T.** (1996). A frizzled homolog functions in a vertebrate Wnt signaling pathway. *Curr Biol* **6**, 1302-6.
- Yang, C. T., Sengelmann, R. D. and Johnson, S. L.** (2004). Larval melanocyte regeneration following laser ablation in zebrafish. *J Invest Dermatol* **123**, 924-9.
- Yisraeli, J. K. and Melton, D. A.** (1988). The material mRNA Vg1 is correctly localized following injection into Xenopus oocytes. *Nature* **336**, 592-5.
- Yucel, G. and Small, S.** (2006). Morphogens: precise outputs from a variable gradient. *Curr Biol* **16**, R29-31.
- Zalik, S. E., Lewandowski, E., Kam, Z. and Geiger, B.** (1999). Cell adhesion and the actin cytoskeleton of the enveloping layer in the zebrafish embryo during epiboly. *Biochem Cell Biol* **77**, 527-42.
- Zerial, M. and McBride, H.** (2001). Rab Proteins as Membrane Organizers. *Nature Reviews Molecular Cell Biology* **2**, 107-119.

- Zhang, J., Houston, D. W., King, M. L., Payne, C., Wylie, C. and Heasman, J.** (1998). The role of maternal VegT in establishing the primary germ layers in *Xenopus* embryos. *Cell* **94**, 515-24.
- Zhang, J. and King, M. L.** (1996). Xenopus VegT RNA is localized to the vegetal cortex during oogenesis and encodes a novel T-box transcription factor involved in mesodermal patterning. *Development* **122**, 4119-29.
- Zhang, Y., Chang, C., Gehling, D. J., Hemmati-Brivanlou, A. and Derynck, R.** (2001). Regulation of Smad degradation and activity by Smurf2, an E3 ubiquitin ligase. *Proc Natl Acad Sci U S A* **98**, 974-9.
- Zhang, Y., Musci, T. and Derynck, R.** (1997). The tumor suppressor Smad4/DPC 4 as a central mediator of Smad function. *Curr Biol* **7**, 270-6.
- Zhou, X., Sasaki, H., Lowe, L., Hogan, B. L. and Kuehn, M. R.** (1993). Nodal is a novel TGF-beta-like gene expressed in the mouse node during gastrulation. *Nature* **361**, 543-7.