

References

- ADLER DA, BRESSLER SL, CHAPMAN VM, PAGE DC and DISTECHE CM. Inactivation of the *Zfx* gene on the mouse X chromosome. *Proc Natl Acad Sci U S A*, 88(11):4592–4595, 1991.
- ADLER DA, RUGARLI EI, LINGENFELTER PA, TSUCHIYA K, POSLINSKI D *et al.* Evidence of evolutionary up-regulation of the single active X chromosome in mammals based on *Clc4* expression levels in *mus spretus* and *mus musculus*. *Proc Natl Acad Sci U S A*, 94(17):9244–9248, 1997.
- ADORJÁN P, DISTLER J, LIPSCHER E, MODEL F, MÜLLER J *et al.* Tumour class prediction and discovery by microarray-based DNA methylation analysis. *Nucleic Acids Res*, 30(5):e21, 2002.
- AGULNIK AI, MITCHELL MJ, MATTEI MG, BORSANI G, AVNER PA, LERNER JL and BISHOP CE. A novel X gene with a widely transcribed Y-linked homologue escapes X-inactivation in mouse and human. *Hum Mol Genet*, 3(6):879–884, 1994.
- ANTEQUERA F and BIRD A. Number of CpG islands and genes in human and mouse. *Proc Natl Acad Sci U S A*, 90(24):11995–11999, 1993.
- ASHWORTH A, RASTAN S, LOVELL-BADGE R and KAY G. X-chromosome inactivation may explain the difference in viability of XO humans and mice. *Nature*, 351(6325):406–408, 1991.
- BAILEY JA, CARREL L, CHAKRAVARTI A and EICHLER EE. Molecular evidence for a relationship between LINE-1 elements and X chromosome inactivation: the Lyon repeat hypothesis. *Proc Natl Acad Sci U S A*, 97(12):6634–6639, 2000.
- BAKER BS, GORMAN M and MARÍN I. Dosage compensation in *Drosophila*. *Annu Rev Genet*, 28:491–521, 1994.
- BARR ML and BERTRAM EG. A morphological distinction between neurones of the male and female, and the behaviour of the nucleolar satellite during accelerated nucleoprotein synthesis. *Nature*, 163(4148):676, 1949.

REFERENCES

- BELL AC, WEST AG and FELSENFELD G. The protein CTCF is required for the enhancer blocking activity of vertebrate insulators. *Cell*, 98(3):387–396, 1999.
- BENJAMIN D, BAKEL IV and CRAIG IW. A novel expression based approach for assessing the inactivation status of human X-linked genes. *Eur J Hum Genet*, 8(2):103–108, 2000.
- BININDA-EMONDS ORP, CARDILLO M, JONES KE, MACPHEE RDE, BECK RMD *et al.* The delayed rise of present-day mammals. *Nature*, 446(7135):507–512, 2007.
- BIRD A, TAGGART M, FROMMER M, MILLER OJ and MACLEOD D. A fraction of the mouse genome that is derived from islands of nonmethylated, CpG-rich DNA. *Cell*, 40(1):91–99, 1985.
- BIRD AP. CpG-rich islands and the function of DNA methylation. *Nature*, 321(6067):209–213, 1986.
- BIRD AP and TAGGART MH. Variable patterns of total DNA and rDNA methylation in animals. *Nucleic Acids Res*, 8(7):1485–1497, 1980.
- BOGGS BA, CONNORS B, SOBEL RE, CHINAULT AC and ALLIS CD. Reduced levels of histone H3 acetylation on the inactive X chromosome in human females. *Chromosoma*, 105(5):303–309, 1996.
- BORSANI G, TONLORENZI R, SIMMLER MC, DANDOLO L, ARNAUD D *et al.* Characterization of a murine gene expressed from the inactive X chromosome. *Nature*, 351(6324):325–329, 1991.
- BROCKDORFF N, ASHWORTH A, KAY GF, MCCABE VM, NORRIS DP, COOPER PJ, SWIFT S and RASTAN S. The product of the mouse Xist gene is a 15 kb inactive X-specific transcript containing no conserved ORF and located in the nucleus. *Cell*, 71(3):515–526, 1992.
- BROWN CJ, BALLABIO A, RUPERT JL, LAFRENIERE RG, GROMPE M, TONLORENZI R and WILLARD HF. A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature*, 349(6304):38–44, 1991.
- BROWN CJ, HENDRICH BD, RUPERT JL, LAFRENIÈRE RG, XING Y, LAWRENCE J and WILLARD HF. The human Xist gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is highly localized within the nucleus. *Cell*, 71(3):527–542, 1992.

REFERENCES

- BROWN CJ and WILLARD HF. The human X-inactivation centre is not required for maintenance of X-chromosome inactivation. *Nature*, 368(6467):154–156, 1994.
- BULL J. *Evolution of sex determining mechanisms*. Benjamin Cummings, Menlo Park, CA, 1983.
- CARREL L, COTTLE AA, GOGLIN KC and WILLARD HF. A first-generation X-inactivation profile of the human X chromosome. *Proc Natl Acad Sci U S A*, 96(25):14440–14444, 1999.
- CARREL L and WILLARD HF. X-inactivation profile reveals extensive variability in X-linked gene expression in females. *Nature*, 434(7031):400–404, 2005.
- CATTANACH BM and ISAACSON JH. Controlling elements in the mouse X chromosome. *Genetics*, 57(2):331–346, 1967.
- CATTANACH BM and WILLIAMS CE. Evidence of non-random X chromosome activity in the mouse. *Genet Res*, 19(3):229–240, 1972.
- CHADWICK LH, PERTZ LM, BROMAN KW, BARTOLOMEI MS and WILLARD HF. Genetic control of X chromosome inactivation in mice: definition of the Xce candidate interval. *Genetics*, 173(4):2103–2110, 2006.
- CLEMSON CM, MCNEIL JA, WILLARD HF and LAWRENCE JB. XIST RNA paints the inactive X chromosome at interphase: evidence for a novel RNA involved in nuclear/chromosome structure. *J Cell Biol*, 132(3):259–275, 1996.
- CLERC P and AVNER P. Role of the region 3' to Xist exon 6 in the counting process of X-chromosome inactivation. *Nat Genet*, 19(3):249–253, 1998.
- CLINE TW and MEYER BJ. Vive la différence: males vs females in flies vs worms. *Annu Rev Genet*, 30:637–702, 1996.
- COMB M and GOODMAN HM. CpG methylation inhibits proenkephalin gene expression and binding of the transcription factor AP-2. *Nucleic Acids Res*, 18(13):3975–3982, 1990.
- COOPER DN, TAGGART MH and BIRD AP. Unmethylated domains in vertebrate DNA. *Nucleic Acids Res*, 11(3):647–658, 1983.
- COOPER DW, EDWARDS C, JAMES E, SHARMAN GB, VANDEBERG JL and GRAVES JAM. Studies on metathrician sex chromosome. VI. A third state of an X-linked gene: partial activity for the paternally derived *pgk-a* allele in cultured fibroblasts of *macropus giganteus* and *m. pangi*. *Aust J Biol Sci*, 30:431–433, 1977.

REFERENCES

- COOPER DW, JOHNSTON PG, WATSON JM and GRAVES JAM. X-inactivation in marsupials and monotremes. *Semin Dev Biol*, 4:117–128, 1993.
- COOPER DW, VANDEBERG JL, SHARMAN GB and POOLE WE. Phosphoglycerate kinase polymorphism in kangaroos provides further evidence for paternal X inactivation. *Nat New Biol*, 230(13):155–157, 1971.
- COSTANZI C and PEHRSON JR. Histone macroH2A1 is concentrated in the inactive X chromosome of female mammals. *Nature*, 393(6685):599–601, 1998.
- COSTELLO JF, FRÜHWALD MC, SMIRAGLIA DJ, RUSH LJ, ROBERTSON GP *et al.* Aberrant CpG-island methylation has non-random and tumour-type-specific patterns. *Nat Genet*, 24(2):132–138, 2000.
- COULONDRE C, MILLER JH, FARABAUGH PJ and GILBERT W. Molecular basis of base substitution hotspots in *escherichia coli*. *Nature*, 274(5673):775–780, 1978.
- CSANKOVSKI G, NAGY A and JAENISCH R. Synergism of Xist RNA, DNA methylation, and histone hypoacetylation in maintaining X chromosome inactivation. *J Cell Biol*, 153(4):773–784, 2001.
- CSANKOVSKI G, PANNING B, BATES B, PEHRSON JR and JAENISCH R. Conditional deletion of Xist disrupts histone macroH2A localization but not maintenance of X inactivation. *Nat Genet*, 22(4):323–324, 1999.
- DAVIDSON RG, NITOWSKY HM and CHILDS B. Demonstration of two populations of cells in the human female heterozygous for glucose-6-phosphate dehydrogenase variants. *Proc Natl Acad Sci U S A*, 50:481–485, 1963.
- DISTECHE CM. Escape from X inactivation in human and mouse. *Trends Genet*, 11(1):17–22, 1995.
- DISTECHE CM, FILIPPOVA GN and TSUCHIYA KD. Escape from X inactivation. *Cytogenet Genome Res*, 99(1-4):36–43, 2002.
- DURET L, EYRE-WALKER A and GALTIER N. A new perspective on isochore evolution. *Gene*, 385:71–74, 2006.
- ECKHARDT F, LEWIN J, CORTESE R, RAKYAN VK, ATTWOOD J *et al.* DNA methylation profiling of human chromosomes 6, 20 and 22. *Nat Genet*, 38(12):1378–1385, 2006.

REFERENCES

- EHRlich M, GAMA-SOSA MA, HUANG LH, MIDGETT RM, KUO KC, McCUNE RA and GEHRKE C. Amount and distribution of 5-methylcytosine in human DNA from different types of tissues of cells. *Nucleic Acids Res*, 10(8):2709–2721, 1982.
- EHRMANN IE, ELLIS PS, MAZEYRAT S, DUTHIE S, BROCKDORFF N *et al.* Characterization of genes encoding translation initiation factor eif-2gamma in mouse and human: sex chromosome localization, escape from X-inactivation and evolution. *Hum Mol Genet*, 7(11):1725–1737, 1998.
- ELLISON J, PASSAGE M, YU LC, YEN P, MOHANDAS TK and SHAPIRO L. Directed isolation of human genes that escape X inactivation. *Somat Cell Mol Genet*, 18(3):259–268, 1992.
- ESTELLER M, CORN PG, BAYLIN SB and HERMAN JG. A gene hypermethylation profile of human cancer. *Cancer Res*, 61(8):3225–3229, 2001.
- FERGUSON-SMITH AC, SASAKI H, CATTANACH BM and SURANI MA. Parental-origin-specific epigenetic modification of the mouse H19 gene. *Nature*, 362(6422):751–755, 1993.
- FIALKOW PJ. X-chromosome inactivation and the *xg* locus. *Am J Hum Genet*, 22(4):460–463, 1970.
- FILIPPOVA GN, CHENG MK, MOORE JM, TRUONG JP, HU YJ, NGUYEN DK, TSUCHIYA KD and DISTECHE CM. Boundaries between chromosomal domains of X inactivation and escape bind CTCF and lack CpG methylation during early development. *Dev Cell*, 8(1):31–42, 2005.
- FISHER EM, BEER-ROMERO P, BROWN LG, RIDLEY A, McNEIL JA *et al.* Homologous ribosomal protein genes on the human X and Y chromosomes: escape from X inactivation and possible implications for Turner syndrome. *Cell*, 63(6):1205–1218, 1990.
- FOSTER JW and GRAVES JA. An *sry*-related sequence on the marsupial X chromosome: implications for the evolution of the mammalian testis-determining gene. *Proc Natl Acad Sci U S A*, 91(5):1927–1931, 1994.
- FRAZER KA, PACTER L, POLIAKOV A, RUBIN EM and DUBCHAK I. VISTA: computational tools for comparative genomics. *Nucleic Acids Res*, 32(Web Server issue):W273–W279, 2004.
- FRIGOLA J, RIBAS M, RISQUES RA and PEINADO MA. Methylome profiling of cancer cells by amplification of inter-methylated sites (AIMS). *Nucleic Acids Res*, 30(7):e28, 2002.

REFERENCES

- FROMMER M, McDONALD LE, MILLAR DS, COLLIS CM, WATT F, GRIGG GW, MOLLOY PL and PAUL CL. A genomic sequencing protocol that yields a positive display of 5-methylcytosine residues in individual DNA strands. *Proc Natl Acad Sci U S A*, 89(5):1827–1831, 1992.
- GARDINER-GARDEN M and FROMMER M. CpG islands in vertebrate genomes. *J Mol Biol*, 196(2):261–282, 1987.
- GARTLER SM and RIGGS AD. Mammalian X-chromosome inactivation. *Annu Rev Genet*, 17:155–190, 1983.
- GENEREUX DP, MINER BE, BERGSTROM CT and LAIRD CD. A population-epigenetic model to infer site-specific methylation rates from double-stranded DNA methylation patterns. *Proc Natl Acad Sci U S A*, 102(16):5802–5807, 2005.
- GITAN RS, SHI H, CHEN CM, YAN PS and HUANG THM. Methylation-specific oligonucleotide microarray: a new potential for high-throughput methylation analysis. *Genome Res*, 12(1):158–164, 2002.
- GONZALGO ML and JONES PA. Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). *Nucleic Acids Res*, 25(12):2529–2531, 1997.
- GONZALGO ML, LIANG G, SPRUCK CH, ZINGG JM, RIDEOUT WM and JONES PA. Identification and characterization of differentially methylated regions of genomic DNA by methylation-sensitive arbitrarily primed pcr. *Cancer Res*, 57(4):594–599, 1997.
- GOODFELLOW P, PYM B, MOHANDAS T and SHAPIRO LJ. The cell surface antigen locus, *mic2x*, escapes X-inactivation. *Am J Hum Genet*, 36(4):777–782, 1984.
- GORMAN JG, DI RE J, TREACY AM and CAHAN A. The application of -Xga antiserum to the question of red cell mosaicism in female heterozygotes. *J Lab Clin Med*, 61:642–649, 1963.
- GOTO Y, GOMEZ M, BROCKDORFF N and FEIL R. Differential patterns of histone methylation and acetylation distinguish active and repressed alleles at X-linked genes. *Cytogenet Genome Res*, 99(1-4):66–74, 2002.
- GRAVES JA. 5-azacytidine-induced re-expression of alleles on the inactive X chromosome in a hybrid mouse cell line. *Exp Cell Res*, 141(1):99–105, 1982.

REFERENCES

- GRAVES JA. The evolution of mammalian sex chromosomes and the origin of sex determining genes. *Philos Trans R Soc Lond B Biol Sci*, 350(1333):305–11; discussion 311–2, 1995.
- GRAVES JA. Mammals that break the rules: genetics of marsupials and monotremes. *Annu Rev Genet*, 30:233–260, 1996.
- GRAVES JA and WATSON JM. Mammalian sex chromosomes: evolution of organization and function. *Chromosoma*, 101(2):63–68, 1991.
- GRAVES JAM. Sex chromosome specialization and degeneration in mammals. *Cell*, 124(5):901–914, 2006.
- GREENFIELD A, CARREL L, PENNISI D, PHILIPPE C, QUADERI N *et al.* The *utx* gene escapes X inactivation in mice and humans. *Hum Mol Genet*, 7(4):737–742, 1998.
- GREGER V, PASSARGE E, HÖPPING W, MESSMER E and HORSTHEMKE B. Epigenetic changes may contribute to the formation and spontaneous regression of retinoblastoma. *Hum Genet*, 83(2):155–158, 1989.
- GRUNAU C, SCHATTEVOY R, MACHE N and ROSENTHAL A. MethTools—a toolbox to visualize and analyze DNA methylation data. *Nucleic Acids Res*, 28(5):1053–1058, 2000.
- GRUNSTEIN M. Histone acetylation in chromatin structure and transcription. *Nature*, 389(6649):349–352, 1997.
- HEARD E, ROUGEULLE C, ARNAUD D, AVNER P, ALLIS CD and SPECTOR DL. Methylation of histone H3 at Lys-9 is an early mark on the X chromosome during X inactivation. *Cell*, 107(6):727–738, 2001.
- HEDGES SB and KUMAR S. Precision of molecular time estimates. *Trends Genet*, 20(5):242–247, 2004.
- HENDRIKS RW, CHEN ZY, HINDS H, SCHUURMAN RK and CRAIG IW. An X chromosome inactivation assay based on differential methylation of a CpG island coupled to a VNTR polymorphism at the 5' end of the monoamine oxidase A gene. *Hum Mol Genet*, 1(3):187–194, 1992.
- HERMAN JG, GRAFF JR, MYÖHÄNEN S, NELKIN BD and BAYLIN SB. Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proc Natl Acad Sci U S A*, 93(18):9821–9826, 1996.

REFERENCES

- HERMAN JG, LATIF F, WENG Y, LERMAN MI, ZBAR B *et al.* Silencing of the VHL tumor-suppressor gene by DNA methylation in renal carcinoma. *Proc Natl Acad Sci U S A*, 91(21):9700–9704, 1994.
- HERZING LB, ROMER JT, HORN JM and ASHWORTH A. Xist has properties of the X-chromosome inactivation centre. *Nature*, 386(6622):272–275, 1997.
- HORE TA, KOINA E, WAKEFIELD MJ and GRAVES JAM. The region homologous to the X-chromosome inactivation centre has been disrupted in marsupial and monotreme mammals. *Chromosome Res*, 15(2):147–161, 2007.
- HORNECKER JL, SAMOLLOW PB, ROBINSON ES, VANDEBERG JL and MCCARREY JR. Meiotic sex chromosome inactivation in the marsupial *monodelphis domestica*. *Genesis*, 45(11):696–708, 2007.
- HUANG TH, LAUX DE, HAMLIN BC, TRAN P, TRAN H and LUBAHN DB. Identification of DNA methylation markers for human breast carcinomas using the methylation-sensitive restriction fingerprinting technique. *Cancer Res*, 57(6):1030–1034, 1997.
- HUANG TH, PERRY MR and LAUX DE. Methylation profiling of CpG islands in human breast cancer cells. *Hum Mol Genet*, 8(3):459–470, 1999.
- HUYNH KD and LEE JT. X-chromosome inactivation: a hypothesis linking ontogeny and phylogeny. *Nat Rev Genet*, 6(5):410–418, 2005.
- JEGALIAN K and PAGE DC. A proposed path by which genes common to mammalian X and Y chromosomes evolve to become X inactivated. *Nature*, 394(6695):776–780, 1998.
- JEPPESEN P and TURNER BM. The inactive X chromosome in female mammals is distinguished by a lack of histone H4 acetylation, a cytogenetic marker for gene expression. *Cell*, 74(2):281–289, 1993.
- JOHNSTON PG, SHARMAN GB, JAMES EA and COOPER DW. Studies on metatherian sex chromosomes. VII. Glucose-6-phosphate dehydrogenase expression in tissues and cultured fibroblasts of kangaroos. *Aust J Biol Sci*, 31:415–424, 1978.
- JONES PA, TAYLOR SM, MOHANDAS T and SHAPIRO LJ. Cell cycle-specific reactivation of an inactive X-chromosome locus by 5-azadeoxycytidine. *Proc Natl Acad Sci U S A*, 79(4):1215–1219, 1982.

REFERENCES

- KASLOW DC and MIGEON BR. DNA methylation stabilizes X chromosome inactivation in eutherians but not in marsupials: evidence for multistep maintenance of mammalian X dosage compensation. *Proc Natl Acad Sci U S A*, 84(17):6210–6214, 1987.
- KAWAKAMI T, OKAMOTO K, OGAWA O and OKADA Y. XIST unmethylated DNA fragments in male-derived plasma as a tumour marker for testicular cancer. *Lancet*, 363(9402):40–42, 2004.
- KAY GF, ASHWORTH A, PENNY GD, DUNLOP M, SWIFT S, BROCKDORFF N and RASTAN S. A candidate spermatogenesis gene on the mouse Y chromosome is homologous to ubiquitin-activating enzyme E1. *Nature*, 354(6353):486–489, 1991.
- KEOHANE AM, O’NEILL LP, BELYAEV ND, LAVENDER JS and TURNER BM. X-inactivation and histone H4 acetylation in embryonic stem cells. *Dev Biol*, 180(2):618–630, 1996.
- KESHET I, YISRAELI J and CEDAR H. Effect of regional DNA methylation on gene expression. *Proc Natl Acad Sci U S A*, 82(9):2560–2564, 1985.
- LABARGA A, VALENTIN F, ANDERSON M and LOPEZ R. Web services at the European bioinformatics institute. *Nucleic Acids Res*, 35(Web Server issue):W6–11, 2007.
- LAHN BT and PAGE DC. Four evolutionary strata on the human X chromosome. *Science*, 286(5441):964–967, 1999.
- LANDER ES, LINTON LM, BIRREN B, NUSBAUM C, ZODY MC *et al.* Initial sequencing and analysis of the human genome. *Nature*, 409(6822):860–921, 2001.
- LARKIN MA, BLACKSHIELDS G, BROWN NP, CHENNA R, MCGETTIGAN PA *et al.* Clustal W and Clustal X version 2.0. *Bioinformatics*, 23(21):2947–2948, 2007.
- LARSEN F, GUNDERSEN G, LOPEZ R and PRYDZ H. CpG islands as gene markers in the human genome. *Genomics*, 13(4):1095–1107, 1992.
- LEE JT and JAENISCH R. Long-range *cis* effects of ectopic X-inactivation centres on a mouse autosome. *Nature*, 386(6622):275–279, 1997.
- LEE JT and LU N. Targeted mutagenesis of Tsix leads to nonrandom X inactivation. *Cell*, 99(1):47–57, 1999.

REFERENCES

- LEWIS JA. Structure and expression of the Chinese hamster thymidine kinase gene. *Mol Cell Biol*, 6(6):1998–2010, 1986.
- LIN H, GUPTA V, VERMILYEA MD, FALCIANI F, LEE JT, O'NEILL LP and TURNER BM. Dosage compensation in the mouse balances up-regulation and silencing of X-linked genes. *PLoS Biol*, 5(12):e326, 2007.
- LINDBLAD-TOH K, WADE CM, MIKKELSEN TS, KARLSSON EK, JAFFE DB *et al.* Genome sequence, comparative analysis and haplotype structure of the domestic dog. *Nature*, 438(7069):803–819, 2005.
- LINGENFELTER PA, ADLER DA, POSLINSKI D, THOMAS S, ELLIOTT RW, CHAPMAN VM and DISTECHE CM. Escape from X inactivation of *smcx* is preceded by silencing during mouse development. *Nat Genet*, 18(3):212–213, 1998.
- LISTER R, O'MALLEY RC, TONTI-FILIPPINI J, GREGORY BD, BERRY CC, MILLAR AH and ECKER JR. Highly integrated single-base resolution maps of the epigenome in Arabidopsis. *Cell*, 133(3):523–536, 2008.
- LOBANENKOV VV, NICOLAS RH, ADLER VV, PATERSON H, KLENOVA EM, POLOTSKAJA AV and GOODWIN GH. A novel sequence-specific DNA binding protein which interacts with three regularly spaced direct repeats of the CCCTC-motif in the 5'-flanking sequence of the chicken *c-myc* gene. *Oncogene*, 5(12):1743–1753, 1990.
- LOCK LF, MELTON DW, CASKEY CT and MARTIN GR. Methylation of the mouse *hprt* gene differs on the active and inactive X chromosomes. *Mol Cell Biol*, 6(3):914–924, 1986.
- LOCK LF, TAKAGI N and MARTIN GR. Methylation of the *hprt* gene on the inactive X occurs after chromosome inactivation. *Cell*, 48(1):39–46, 1987.
- LOEBEL DA and JOHNSTON PG. Analysis of DNase 1 sensitivity and methylation of active and inactive X chromosomes of kangaroos (*macropus robustus*) by *in situ* nick translation. *Chromosoma*, 102(2):81–87, 1993.
- LOEBEL DA and JOHNSTON PG. Methylation analysis of a marsupial X-linked CpG island by bisulfite genomic sequencing. *Genome Res*, 6(2):114–123, 1996.
- LYON MF. Gene action in the X-chromosome of the mouse (*mus musculus l.*). *Nature*, 190:372–373, 1961.
- LYON MF. X-chromosome inactivation: a repeat hypothesis. *Cytogenet Cell Genet*, 80(1-4):133–137, 1998.

REFERENCES

- LYON MF, SEARLE AG, FORD CE and OHNO S. A mouse translocation suppressing sex-linked variegation. *Cytogenetics*, 15:306–323, 1964.
- MCKEON C, OHKUBO H, PASTAN I and DE CROMBRUGGHE B. Unusual methylation pattern of the alpha 2 (I) collagen gene. *Cell*, 29(1):203–210, 1982.
- MERLO A, HERMAN JG, MAO L, LEE DJ, GABRIELSON E, BURGER PC, BAYLIN SB and SIDRANSKY D. 5' CpG island methylation is associated with transcriptional silencing of the tumour suppressor *p16/cdkn2/mts1* in human cancers. *Nat Med*, 1(7):686–692, 1995.
- MERMOUD JE, COSTANZI C, PEHRSON JR and BROCKDORFF N. Histone macroH2A1.2 relocates to the inactive X chromosome after initiation and propagation of X-inactivation. *J Cell Biol*, 147(7):1399–1408, 1999.
- MEYER BJ. Sex in the worm counting and compensating X-chromosome dose. *Trends Genet*, 16(6):247–253, 2000.
- MIGEON BR, DE BEUR SJ and AXELMAN J. Frequent derepression of *g6pd* and *hpert* on the marsupial inactive X chromosome associated with cell proliferation *in vitro*. *Exp Cell Res*, 182(2):597–609, 1989.
- MIGEON BR, CHOWDHURY AK, DUNSTON JA and MCINTOSH I. Identification of TSIX, encoding an RNA antisense to human XIST, reveals differences from its murine counterpart: implications for X inactivation. *Am J Hum Genet*, 69(5):951–960, 2001.
- MIGEON BR, LEE CH, CHOWDHURY AK and CARPENTER H. Species differences in TSIX/Tsix reveal the roles of these genes in X-chromosome inactivation. *Am J Hum Genet*, 71(2):286–293, 2002.
- MIKKELSEN TS, WAKEFIELD MJ, AKEN B, AMEMIYA CT, CHANG JL *et al*. Genome of the marsupial *monodelphis domestica* reveals innovation in non-coding sequences. *Nature*, 447(7141):167–177, 2007.
- MITCHELL MJ, WOODS DR, WILCOX SA, GRAVES JA and BISHOP CE. Marsupial Y chromosome encodes a homologue of the mouse Y-linked candidate spermatogenesis gene *ube1y*. *Nature*, 359(6395):528–531, 1992.
- MOHANDAS T, SPARKES RS, HELLKUHL B, GRZESCHIK KH and SHAPIRO LJ. Expression of an X-linked gene from an inactive human X chromosome in mouse-human hybrid cells: further evidence for the noninactivation of the steroid sulfatase locus in man. *Proc Natl Acad Sci U S A*, 77(11):6759–6763, 1980.

REFERENCES

- MOREY C, ARNAUD D, AVNER P and CLERC P. Tsix-mediated repression of Xist accumulation is not sufficient for normal random X inactivation. *Hum Mol Genet*, 10(13):1403–1411, 2001.
- MUKHOPADHYAY R, YU W, WHITEHEAD J, XU J, LEZCANO M *et al.* The binding sites for the chromatin insulator protein CTCF map to DNA methylation-free domains genome-wide. *Genome Res*, 14(8):1594–1602, 2004.
- MURPHY WJ, PEVZNER PA and O'BRIEN SJ. Mammalian phylogenomics comes of age. *Trends Genet*, 20(12):631–639, 2004.
- NAMEKAWA SH, VANDEBERG JL, MCCARREY JR and LEE JT. Sex chromosome silencing in the marsupial male germ line. *Proc Natl Acad Sci U S A*, 104(23):9730–9735, 2007.
- NGUYEN DK and DISTECHE CM. Dosage compensation of the active X chromosome in mammals. *Nat Genet*, 38(1):47–53, 2006.
- NORDQUIST N and ORELAND L. Monoallelic expression of *maoa* in skin fibroblasts. *Biochem Biophys Res Commun*, 348(2):763–767, 2006.
- OGAWA Y and LEE JT. Xite, X-inactivation intergenic transcription elements that regulate the probability of choice. *Mol Cell*, 11(3):731–743, 2003.
- OHNO S. *Sex chromosomes and sex-linked genes*. Springer-Verlag Berlin, 1967.
- OHNO S and HAUSCHKA TS. Allocyclcy of the X-chromosome in tumors and normal tissues. *Cancer Res*, 20:541–545, 1960.
- OHNO S, KAPLAN WD and KINOSITA R. Formation of the sex chromatin by a single X-chromosome in liver cells of *rattus norvegicus*. *Exp Cell Res*, 18:415–418, 1959.
- OLEK A, OSWALD J and WALTER J. A modified and improved method for bisulphite based cytosine methylation analysis. *Nucleic Acids Res*, 24(24):5064–5066, 1996.
- PARK JG and CHAPMAN VM. CpG island promoter region methylation patterns of the inactive-X-chromosome hypoxanthine phosphoribosyltransferase (*hprt*) gene. *Mol Cell Biol*, 14(12):7975–7983, 1994.
- PENNY GD, KAY GF, SHEARDOWN SA, RASTAN S and BROCKDORFF N. Requirement for Xist in X chromosome inactivation. *Nature*, 379(6561):131–137, 1996.

REFERENCES

- PLATH K, MLYNARCZYK-EVANS S, NUSINOW DA and PANNING B. Xist RNA and the mechanism of X chromosome inactivation. *Annu Rev Genet*, 36:233–278, 2002.
- RASTAN S. Non-random X-chromosome inactivation in mouse X-autosome translocation embryos—location of the inactivation centre. *J Embryol Exp Morphol*, 78:1–22, 1983.
- RASTAN S and ROBERTSON EJ. X-chromosome deletions in embryo-derived (EK) cell lines associated with lack of X-chromosome inactivation. *J Embryol Exp Morphol*, 90:379–388, 1985.
- RENS W, GRÜTZNER F, O'BRIEN PCM, FAIRCLOUGH H, GRAVES JAM and FERGUSON-SMITH MA. Resolution and evolution of the duck-billed platypus karyotype with an X1Y1X2Y2X3Y3X4Y4X5Y5 male sex chromosome constitution. *Proc Natl Acad Sci U S A*, 101(46):16257–16261, 2004.
- RICHARDSON BJ, CZUPPON AB and SHARMAN GB. Inheritance of glucose-6-phosphate dehydrogenase variation in kangaroos. *Nat New Biol*, 230(13):154–155, 1971.
- RIGGS AD. X inactivation, differentiation, and DNA methylation. *Cytogenet Cell Genet*, 14(1):9–25, 1975.
- ROSS MT, GRAFHAM DV, COFFEY AJ, SCHERER S, MCLAY K *et al.* The DNA sequence of the human X chromosome. *Nature*, 434(7031):325–337, 2005.
- RUSSELL GJ, WALKER PM, ELTON RA and SUBAK-SHARPE JH. Doublet frequency analysis of fractionated vertebrate nuclear DNA. *J Mol Biol*, 108(1):1–23, 1976.
- RUSSELL L and MONTGOMERY C. The use of X-autosome translocations in locating the X-chromosome inactivation center. *Genetics*, 52:470–471, 1965.
- RUSSELL LB. Genetics of mammalian sex chromosomes. *Science*, 133:1795–1803, 1961.
- RUSSELL LB. Mammalian X-chromosome action: inactivation limited in spread and region of origin. *Science*, 140:976–978, 1963.
- SAKAI T, TOGUCHIDA J, OHTANI N, YANDELL DW, RAPAPORT JM and DRYJA TP. Allele-specific hypermethylation of the retinoblastoma tumor-suppressor gene. *Am J Hum Genet*, 48(5):880–888, 1991.

REFERENCES

- SAMOLLOW PB. Status and applications of genomic resources for the gray, short-tailed opossum, *monodelphis domestica*, an american marsupial model for comparative biology. *Australian Journal of Zoology*, 54:173–196, 2006.
- SAMOLLOW PB, FORD AL and VANDEBERG JL. X-linked gene expression in the Virginia opossum: differences between the paternally derived *gpd* and *pgk-a* loci. *Genetics*, 115(1):185–195, 1987.
- SAMOLLOW PB, JOHNSTON PG, FORD AL and VANDEBERG JL. X-linked gene expression in metatherian fibroblasts: evidence from the *gpd* and *pgk-a* loci of the virginia opossum and the red-necked wallaby. *Biochem Genet*, 27(5-6):313–320, 1989.
- SCHIEBEL K, WEISS B, WÖHRLE D and RAPPOLD G. A human pseudoautosomal gene, ADP/ATP translocase, escapes X-inactivation whereas a homologue on Xq is subject to X-inactivation. *Nat Genet*, 3(1):82–87, 1993.
- SCHMID CW. Does SINE evolution preclude Alu function? *Nucleic Acids Res*, 26(20):4541–4550, 1998.
- SCHNEIDER-GÄDICKE A, BEER-ROMERO P, BROWN LG, NUSSBAUM R and PAGE DC. *ZFX* has a gene structure similar to *zfy*, the putative human sex determinant, and escapes X inactivation. *Cell*, 57(7):1247–1258, 1989.
- SCHUMACHER A, KAPRANOV P, KAMINSKY Z, FLANAGAN J, ASSADZADEH A *et al.* Microarray-based DNA methylation profiling: technology and applications. *Nucleic Acids Res*, 34(2):528–542, 2006.
- SHAPIRO LJ, MOHANDAS T, WEISS R and ROMEO G. Non-inactivation of an X-chromosome locus in man. *Science*, 204(4398):1224–1226, 1979.
- SHARMAN GB. Late DNA replication in the paternally derived X chromosome of female kangaroos. *Nature*, 230(5291):231–232, 1971.
- SIMMLER MC, CATTANACH BM, RASBERRY C, ROUGEULLE C and AVNER P. Mapping the murine Xce locus with (CA)_n repeats. *Mamm Genome*, 4(9):523–530, 1993.
- SINGER-SAM J, GRANT M, LEBON JM, OKUYAMA K, CHAPMAN V, MONK M and RIGGS AD. Use of a HpaII-polymerase chain reaction assay to study DNA methylation in the *pgk-1* CpG island of mouse embryos at the time of X-chromosome inactivation. *Mol Cell Biol*, 10(9):4987–4989, 1990a.

REFERENCES

- SINGER-SAM J, LEBON JM, TANGUAY RL and RIGGS AD. A quantitative HpaII-PCR assay to measure methylation of DNA from a small number of cells. *Nucleic Acids Res*, 18(3):687, 1990b.
- SMITS G, MUNGALL AJ, GRIFFITHS-JONES S, SMITH P, BEURY D *et al.* Conservation of the H19 noncoding RNA and H19-IGF2 imprinting mechanism in therians. *Nat Genet*, 40(8):971–976, 2008.
- SPENCER JA, SINCLAIR AH, WATSON JM and GRAVES JA. Genes on the short arm of the human X chromosome are not shared with the marsupial X. *Genomics*, 11(2):339–345, 1991a.
- SPENCER JA, WATSON JM and GRAVES JA. The X chromosome of marsupials shares a highly conserved region with eutherians. *Genomics*, 9(4):598–604, 1991b.
- STEIN R, SCIAKY-GALLILI N, RAZIN A and CEDAR H. Pattern of methylation of two genes coding for housekeeping functions. *Proc Natl Acad Sci U S A*, 80(9):2422–2426, 1983.
- STÖGER R, KAJIMURA TM, BROWN WT and LAIRD CD. Epigenetic variation illustrated by DNA methylation patterns of the fragile-X gene *fmr1*. *Hum Mol Genet*, 6(11):1791–1801, 1997.
- STÖGER R, KUBICKA P, LIU CG, KAFRI T, RAZIN A, CEDAR H and BARLOW DP. Maternal-specific methylation of the imprinted mouse *igf2r* locus identifies the expressed locus as carrying the imprinting signal. *Cell*, 73(1):61–71, 1993.
- STIRZAKER C, MILLAR DS, PAUL CL, WARNECKE PM, HARRISON J, VINCENT PC, FROMMER M and CLARK SJ. Extensive DNA methylation spanning the *rb* promoter in retinoblastoma tumors. *Cancer Res*, 57(11):2229–2237, 1997.
- SUZUKI S, ONO R, NARITA T, PASK AJ, SHAW G *et al.* Retrotransposon silencing by DNA methylation can drive mammalian genomic imprinting. *PLoS Genet*, 3(4):e55, 2007.
- SWARTZ MN, TRAUTNER TA and KORNBERG A. Enzymatic synthesis of deoxyribonucleic acid. XI. Further studies on nearest neighbor base sequences in deoxyribonucleic acids. *J Biol Chem*, 237:1961–1967, 1962.
- TAKAGI N and SASAKI M. Preferential inactivation of the paternally derived X chromosome in the extraembryonic membranes of the mouse. *Nature*, 256(5519):640–642, 1975.

REFERENCES

- TAKAI D and JONES PA. Comprehensive analysis of CpG islands in human chromosomes 21 and 22. *Proc Natl Acad Sci U S A*, 99(6):3740–3745, 2002.
- TREMBLAY KD, SAAM JR, INGRAM RS, TILGHMAN SM and BARTOLOMEI MS. A paternal-specific methylation imprint marks the alleles of the mouse *h19* gene. *Nat Genet*, 9(4):407–413, 1995.
- TRIBIOLI C, TAMANINI F, PATROSSO C, MILANESI L, VILLA A *et al.* Methylation and sequence analysis around Eagi sites: identification of 28 new CpG islands in Xq24-Xq28. *Nucleic Acids Res*, 20(4):727–733, 1992.
- TSUCHIYA KD, GREALLY JM, YI Y, NOEL KP, TRUONG JP and DISTECHE CM. Comparative sequence and X-inactivation analyses of a domain of escape in human Xp11.2 and the conserved segment in mouse. *Genome Res*, 14(7):1275–1284, 2004.
- TYKOCINSKI ML and MAX EE. CG dinucleotide clusters in *mhc* genes and in 5' demethylated genes. *Nucleic Acids Res*, 12(10):4385–4396, 1984.
- VANDEBERG J. The gray short-tailed opossum (*monodelphis-domestica*) as a model didelphid species for genetic research. *Australian Journal of Zoology*, 37:235–247, 1989.
- VEYRUNES F, WATERS PD, MIETHKE P, RENS W, MCMILLAN D *et al.* Bird-like sex chromosomes of platypus imply recent origin of mammal sex chromosomes. *Genome Res*, 18(6):965–973, 2008.
- WAKEFIELD MJ, KEOHANE AM, TURNER BM and GRAVES JA. Histone underacetylation is an ancient component of mammalian X chromosome inactivation. *Proc Natl Acad Sci U S A*, 94(18):9665–9668, 1997.
- WALLIS MC, WATERS PD, DELBRIDGE ML, KIRBY PJ, PASK AJ *et al.* Sex determination in platypus and echidna: autosomal location of *sox3* confirms the absence of *sry* from monotremes. *Chromosome Res*, 15(8):949–959, 2007.
- WARNECKE PM, STIRZAKER C, MELKI JR, MILLAR DS, PAUL CL and CLARK SJ. Detection and measurement of pcr bias in quantitative methylation analysis of bisulphite-treated DNA. *Nucleic Acids Res*, 25(21):4422–4426, 1997.
- WARREN WC, HILLIER LW, GRAVES JAM, BIRNEY E, PONTING CP *et al.* Genome analysis of the platypus reveals unique signatures of evolution. *Nature*, 453(7192):175–183, 2008.

REFERENCES

- WATERSTON RH, LINDBLAD-TOH K, BIRNEY E, ROGERS J, ABRIL JF *et al.* Initial sequencing and comparative analysis of the mouse genome. *Nature*, 420(6915):520–562, 2002.
- WEBER M, DAVIES JJ, WITTIG D, OAKELEY EJ, HAASE M, LAM WL and SCHÜBELER D. Chromosome-wide and promoter-specific analyses identify sites of differential DNA methylation in normal and transformed human cells. *Nat Genet*, 37(8):853–862, 2005.
- WELSHONS WJ and RUSSELL LB. The y-chromosome as the bearer of male determining factors in the mouse. *Proc Natl Acad Sci U S A*, 45(4):560–566, 1959.
- WILSON DE and REEDER DM, editors. *Mammal Species of the World: A Taxonomic and Geographic Reference*. JHU Press, 2005.
- WOLF SF, JOLLY DJ, LUNNEN KD, FRIEDMANN T and MIGEON BR. Methylation of the hypoxanthine phosphoribosyltransferase locus on the human X chromosome: implications for X-chromosome inactivation. *Proc Natl Acad Sci U S A*, 81(9):2806–2810, 1984.
- WOODBURNE MO, RICH TH and SPRINGER MS. The evolution of tribospheny and the antiquity of mammalian clades. *Mol Phylogenet Evol*, 28(2):360–385, 2003.
- WU J, SALIDO EC, YEN PH, MOHANDAS TK, HENG HH *et al.* The murine *xe169* gene escapes X-inactivation like its human homologue. *Nat Genet*, 7(4):491–496, 1994.
- XUE F, TIAN XC, DU F, KUBOTA C, TANEJA M *et al.* Aberrant patterns of X chromosome inactivation in bovine clones. *Nat Genet*, 31(2):216–220, 2002.
- YEN PH, PATEL P, CHINAULT AC, MOHANDAS T and SHAPIRO LJ. Differential methylation of hypoxanthine phosphoribosyltransferase genes on active and inactive human X chromosomes. *Proc Natl Acad Sci U S A*, 81(6):1759–1763, 1984.
- YEN ZC, MEYER IM, KARALIC S and BROWN CJ. A cross-species comparison of X-chromosome inactivation in Eutheria. *Genomics*, 90(4):453–463, 2007.