

Bibliography

- Ahlemeyer, B. and Baumgart-Vogt, E.** (2005). Optimized protocols for the simultaneous preparation of primary neuronal cultures of the neocortex, hippocampus and cerebellum from individual newborn (P0.5) C57Bl/6J mice. *J Neurosci Methods* **149**, 110–20.
- Akahoshi, N., Murashima, Y. L., Himi, T., Ishizaki, Y. and Ishii, I.** (2007). Increased expression of the lysosomal protease cathepsin S in hippocampal microglia following kainate-induced seizures. *Neurosci Lett* **429**, 136–41.
- Alexopoulou, A. N., Couchman, J. R. and Whiteford, J. R.** (2008). The CMV early enhancer/chicken beta actin (CAG) promoter can be used to drive transgene expression during the differentiation of murine embryonic stem cells into vascular progenitors. *BMC Cell Biol* **9**, 2.
- Altman, S. A., Randers, L. and Rao, G.** (1993). Comparison of trypan blue dye exclusion and fluorometric assays for mammalian cell viability determinations. *Biotechnol Prog* **9**, 671–4.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J.** (1990). Basic local alignment search tool. *J Mol Biol* **215**, 403–10.
- Alvarez-Saavedra, E. and Horvitz, H. R.** (2010). Many families of *C. elegans* microRNAs are not essential for development or viability. *Curr Biol* **20**, 367–73.
- Andersson, T., Rahman, S., Sansom, S. N., Alsiö, J. M., Kaneda, M., Smith, J., O'Carroll, D., Tarakhovsky, A. and Livesey, F. J.** (2010). Reversible block of mouse neural stem cell differentiation in the absence of dicer and microRNAs. *PLoS ONE* **5**, e13453.
- Aravin, A. A., Hannon, G. J. and Brennecke, J.** (2007). The Piwi-piRNA pathway provides an adaptive defense in the transposon arms race. *Science* **318**, 761–4.

- Arnold, F. J. L., Hofmann, F., Bengtson, C. P., Wittmann, M., Vanhoutte, P. and Bading, H.** (2005). Microelectrode array recordings of cultured hippocampal networks reveal a simple model for transcription and protein synthesis-dependent plasticity. *J Physiol (Lond)* **564**, 3–19.
- Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M. and Sherlock, G.** (2000). Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* **25**, 25–9.
- Azuma-Mukai, A., Oguri, H., Mituyama, T., Qian, Z. R., Asai, K., Siomi, H. and Siomi, M. C.** (2008). Characterization of endogenous human Argonautes and their miRNA partners in RNA silencing. *Proc Natl Acad Sci USA* **105**, 7964–9.
- Babiarz, J. E., Ruby, J. G., Wang, Y., Bartel, D. P. and Blelloch, R.** (2008). Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes & Development* **22**, 2773–85.
- Bading, H., Segal, M. M., Sucher, N. J., Dudek, H., Lipton, S. A. and Greenberg, M. E.** (1995). N-methyl-D-aspartate receptors are critical for mediating the effects of glutamate on intracellular calcium concentration and immediate early gene expression in cultured hippocampal neurons. *Neuroscience* **64**, 653–64.
- Baek, D., Villén, J., Shin, C., Camargo, F. D., Gygi, S. P. and Bartel, D. P.** (2008). The impact of microRNAs on protein output. *Nature* **455**, 64–71.
- Baillat, D. and Shiekhattar, R.** (2009). Functional dissection of the human TNRC6 (GW182-related) family of proteins. *Mol Cell Biol* **29**, 4144–55.
- Bak, M., Silahtaroglu, A., Moller, M., Christensen, M., Rath, M. F., Skryabin, B., Tommerup, N. and Kauppinen, S.** (2008). MicroRNA expression in the adult mouse central nervous system. *RNA* **14**, 432–444.
- Ballas, N., Grunseich, C., Lu, D. D., Speh, J. C. and Mandel, G.** (2005). REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell* **121**, 645–57.
- Ballas, N. and Mandel, G.** (2005). The many faces of REST oversee epigenetic programming of neuronal genes. *Curr Opin Neurobiol* **15**, 500–6.

- Barrett, T., Troup, D. B., Wilhite, S. E., Ledoux, P., Rudnev, D., Evangelista, C., Kim, I. F., Soboleva, A., Tomashevsky, M. and Edgar, R.** (2007). NCBI GEO: mining tens of millions of expression profiles—database and tools update. *Nucleic Acids Research* **35**, D760–5.
- Bartel, D. P.** (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **116**, 281–97.
- Bartel, D. P.** (2009). MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215–33.
- Basyuk, E., Suavet, F., Doglio, A., Bordonné, R. and Bertrand, E.** (2003). Human let-7 stem-loop precursors harbor features of RNase III cleavage products. *Nucleic acids research* **31**, 6593–7.
- Bayés, A., van de Lagemaat, L. N., Collins, M. O., Croning, M. D. R., Whittle, I. R., Choudhary, J. S. and Grant, S. G. N.** (2011). Characterization of the proteome, diseases and evolution of the human postsynaptic density. *Nat Neurosci* **14**, 19–21.
- Behm-Ansmant, I., Rehwinkel, J., Doerks, T., Stark, A., Bork, P. and Izaurralde, E.** (2006). mRNA degradation by miRNAs and GW182 requires both CCR4:NOT deadenylase and DCP1:DCP2 decapping complexes. *Genes & Development* **20**, 1885–98.
- Benjamini, Y. and Hochberg, Y.** (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)* **57**, 289–300.
- Bernstein, E., Caudy, A. A., Hammond, S. M. and Hannon, G. J.** (2001). Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* **409**, 363–6.
- Bernstein, E., Kim, S. Y., Carmell, M. A., Murchison, E. P., Alcorn, H., Li, M. Z., Mills, A. A., Elledge, S. J., Anderson, K. V. and Hannon, G. J.** (2003). Dicer is essential for mouse development. *Nat Genet* **35**, 215–7.
- Bhattacharyya, S. N., Habermacher, R., Martine, U., Closs, E. I. and Filipowicz, W.** (2006). Relief of microRNA-mediated translational repression in human cells subjected to stress. *Cell* **125**, 1111–24.

- Bohnsack, M. T., Czaplinski, K. and Gorlich, D.** (2004). Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA (New York, NY)* **10**, 185–91.
- Borchert, G. M., Lanier, W. and Davidson, B. L.** (2006). RNA polymerase III transcribes human microRNAs. *Nat Struct Mol Biol* **13**, 1097–101.
- Brennecke, J., Stark, A., Russell, R. B. and Cohen, S. M.** (2005). Principles of microRNA-target recognition. *PLoS Biol* **3**, e85.
- Brenner, J. L., Jasiewicz, K. L., Fahley, A. F., Kemp, B. J. and Abbott, A. L.** (2010). Loss of individual microRNAs causes mutant phenotypes in sensitized genetic backgrounds in *C. elegans*. *Curr Biol* **20**, 1321–5.
- Brewer, G. J.** (1997). Isolation and culture of adult rat hippocampal neurons. *J Neurosci Methods* **71**, 143–55.
- Brewer, G. J., Boehler, M. D., Pearson, R. A., Demaris, A. A., Ide, A. N. and Wheeler, B. C.** (2009). Neuron network activity scales exponentially with synapse density. *J. Neural Eng.* **6**, 014001.
- Brewer, G. J. and Torricelli, J. R.** (2007). Isolation and culture of adult neurons and neurospheres. *Nat Protoc* **2**, 1490–8.
- Brewer, G. J., Torricelli, J. R., Evege, E. K. and Price, P. J.** (1993). Optimized survival of hippocampal neurons in B27-supplemented Neurobasal, a new serum-free medium combination. *J Neurosci Res* **35**, 567–76.
- Cai, X., Hagedorn, C. H. and Cullen, B. R.** (2004). Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA (New York, NY)* **10**, 1957–66.
- Cao, X., Pfaff, S. L. and Gage, F. H.** (2007). A functional study of miR-124 in the developing neural tube. *Genes & Development* **21**, 531–6.
- Carthew, R. W. and Sontheimer, E. J.** (2009). Origins and Mechanisms of miRNAs and siRNAs. *Cell* **136**, 642–55.
- Chekulaeva, M., Filipowicz, W. and Parker, R.** (2009). Multiple independent domains of dGW182 function in miRNA-mediated repression in *Drosophila*. *RNA (New York, NY)* **15**, 794–803.

- Chendrimada, T. P., Finn, K. J., Ji, X., Baillat, D., Gregory, R. I., Liebhaber, S. A., Pasquinelli, A. E. and Shiekhattar, R.** (2007). MicroRNA silencing through RISC recruitment of eIF6. *Nature* **447**, 823–8.
- Chendrimada, T. P., Gregory, R. I., Kumaraswamy, E., Norman, J., Cooch, N., Nishikura, K. and Shiekhattar, R.** (2005). TRBP recruits the Dicer complex to Ago2 for microRNA processing and gene silencing. *Nature* **436**, 740–4.
- Cheng, L.-C., Pastrana, E., Tavazoie, M. and Doetsch, F.** (2009). miR-124 regulates adult neurogenesis in the subventricular zone stem cell niche. *Nat Neurosci* **12**, 399–408.
- Chi, S., Zang, J., Mele, A. and Darnell, R.** (2009). Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature* **460**, 479–86.
- Christodoulou, F., Raible, F., Tomer, R., Simakov, O., Trachana, K., Klaus, S., Snijman, H., Hannon, G. J., Bork, P. and Arendt, D.** (2010). Ancient animal microRNAs and the evolution of tissue identity. *Nature* **463**, 1084–8.
- Cifuentes, D., Xue, H., Taylor, D. W., Patnode, H., Mishima, Y., Cheloufi, S., Ma, E., Mane, S., Hannon, G. J., Lawson, N. D., Wolfe, S. A. and Giraldez, A. J.** (2010). A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science* **328**, 1694–8.
- Clark, A. M., Goldstein, L. D., Tevlin, M., Tavaré, S., Shaham, S. and Miska, E. A.** (2010). The microRNA miR-124 controls gene expression in the sensory nervous system of *Caenorhabditis elegans*. *Nucleic Acids Research* **38**, 3780–93.
- Cogoni, C., Irelan, J. T., Schumacher, M., Schmidhauser, T. J., Selker, E. U. and Macino, G.** (1996). Transgene silencing of the al-1 gene in vegetative cells of *Neurospora* is mediated by a cytoplasmic effector and does not depend on DNA-DNA interactions or DNA methylation. *EMBO J* **15**, 3153–63.
- Conaco, C., Otto, S., Han, J.-J. and Mandel, G.** (2006). Reciprocal actions of REST and a microRNA promote neuronal identity. *Proc Natl Acad Sci USA* **103**, 2422–7.
- Connelly, S. and Manley, J. L.** (1988). A functional mRNA polyadenylation signal is required for transcription termination by RNA polymerase II. *Genes & Development* **2**, 440–52.

- Davis, E., Caiment, F., Tordoir, X., Cavaillé, J., Ferguson-Smith, A., Cockett, N., Georges, M. and Charlier, C.** (2005). RNAi-mediated allelic trans-interaction at the imprinted Rtl1/Peg11 locus. *Curr Biol* **15**, 743–9.
- Davis, T. H., Cuellar, T. L., Koch, S. M., Barker, A. J., Harfe, B. D., Mcmanus, M. T. and Ullian, E. M.** (2008). Conditional loss of Dicer disrupts cellular and tissue morphogenesis in the cortex and hippocampus. *J Neurosci* **28**, 4322–30.
- de Carvalho, F., Gheysen, G., Kushnir, S., Montagu, M. V., Inzé, D. and Castresana, C.** (1992). Suppression of beta-1,3-glucanase transgene expression in homozygous plants. *EMBO J* **11**, 2595–602.
- Dichter, M. A.** (1978). Rat cortical neurons in cell culture: culture methods, cell morphology, electrophysiology, and synapse formation. *Brain Res* **149**, 279–93.
- Doench, J. G. and Sharp, P. A.** (2004). Specificity of microRNA target selection in translational repression. *Genes & Development* **18**, 504–11.
- Du, P., Kibbe, W. A. and Lin, S. M.** (2008). lumi: a pipeline for processing Illumina microarray. *Bioinformatics* **24**, 1547–8.
- Elbashir, S. M., Lendeckel, W. and Tuschl, T.** (2001). RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes & Development* **15**, 188–200.
- Elia, L., Quintavalle, M., Zhang, J., Contu, R., Cossu, L., Latronico, M. V. G., Peterson, K. L., Indolfi, C., Catalucci, D., Chen, J., Courtneidge, S. A. and Condorelli, G.** (2009). The knockout of miR-143 and -145 alters smooth muscle cell maintenance and vascular homeostasis in mice: correlates with human disease. *Cell Death Differ* **16**, 1590–8.
- Enright, A. J., John, B., Gaul, U., Tuschl, T., Sander, C. and Marks, D. S.** (2003). MicroRNA targets in Drosophila. *Genome Biol* **5**, R1.
- Eulalio, A., Huntzinger, E. and Izaurralde, E.** (2008). GW182 interaction with Argonaute is essential for miRNA-mediated translational repression and mRNA decay. *Nat Struct Mol Biol* **15**, 346–53.
- Eulalio, A., Huntzinger, E., Nishihara, T., Rehwinkel, J., Fauser, M. and Izaurralde, E.** (2009). Deadenylation is a widespread effect of miRNA regulation. *RNA (New York, NY)* **15**, 21–32.

- Fabian, M. R., Sonenberg, N. and Filipowicz, W.** (2010). Regulation of mRNA translation and stability by microRNAs. *Annu Rev Biochem* **79**, 351–79.
- Falcon, S. and Gentleman, R.** (2007). Using GOSTATS to test gene lists for GO term association. *Bioinformatics* **23**, 257–8.
- Farh, K. K.-H., Grimson, A., Jan, C., Lewis, B. P., Johnston, W. K., Lim, L. P., Burge, C. B. and Bartel, D. P.** (2005). The widespread impact of mammalian MicroRNAs on mRNA repression and evolution. *Science* **310**, 1817–21.
- Felgner, P. L., Gadek, T. R., Holm, M., Roman, R., Chan, H. W., Wenz, M., Northrop, J. P., Ringold, G. M. and Danielsen, M.** (1987). Lipofection: a highly efficient, lipid-mediated DNA-transfection procedure. *Proc Natl Acad Sci USA* **84**, 7413–7.
- Flicek, P., Aken, B. L., Beal, K., Ballester, B., Caccamo, M., Chen, Y., Clarke, L., Coates, G., Cunningham, F., Cutts, T., Down, T., Dyer, S. C., Eyre, T., Fitzgerald, S., Fernandez-Banet, J., Gräf, S., Haider, S., Hammond, M., Holland, R., Howe, K. L., Howe, K., Johnson, N., Jenkinson, A., Kähäri, A., Keefe, D., Kokocinski, F., Kulesha, E., Lawson, D., Longden, I., Megy, K., Meidl, P., Overduin, B., Parker, A., Pritchard, B., Prlic, A., Rice, S., Rios, D., Schuster, M., Sealy, I., Slater, G., Smedley, D., Spudich, G., Trevanian, S., Vilella, A. J., Vogel, J., White, S., Wood, M., Birney, E., Cox, T., Curwen, V., Durbin, R., Fernandez-Suarez, X. M., Herrero, J., Hubbard, T. J. P., Kasprzyk, A., Proctor, G., Smith, J., Ureta-Vidal, A. and Searle, S.** (2008). Ensembl 2008. *Nucleic Acids Res* **36**, D707–14.
- Flynt, A. S., Thatcher, E. J., Burkewitz, K., Li, N., Liu, Y. and Patton, J. G.** (2009). miR-8 microRNAs regulate the response to osmotic stress in zebrafish embryos. *J Cell Biol* **185**, 115–27.
- Freeman, M. R.** (2010). Specification and morphogenesis of astrocytes. *Science* **330**, 774–8.
- Freeman, T. C., Goldovsky, L., Brosch, M., van Dongen, S., Mazière, P., Grock, R. J., Freilich, S., Thornton, J. and Enright, A. J.** (2007). Construction, visualisation, and clustering of transcription networks from microarray expression data. *PLoS Comput Biol* **3**, 2032–42.

- Gao, J., Wang, W.-Y., Mao, Y.-W., Gräff, J., Guan, J.-S., Pan, L., Mak, G., Kim, D., Su, S. C. and Tsai, L.-H.** (2010). A novel pathway regulates memory and plasticity via SIRT1 and miR-134. *Nature* **466**, 1105–9.
- Gautier, L., Cope, L., Bolstad, B. M. and Irizarry, R. A.** (2004). affy—analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* **20**, 307–15.
- Gelder, R. N. V., von Zastrow, M. E., Yool, A., Dement, W. C., Barchas, J. D. and Eberwine, J. H.** (1990). Amplified RNA synthesized from limited quantities of heterogeneous cDNA. *Proc Natl Acad Sci USA* **87**, 1663–7.
- Gentleman, R. C., Carey, V. J., Bates, D. M., Bolstad, B., Dettling, M., Du-
doit, S., Ellis, B., Gautier, L., Ge, Y., Gentry, J., Hornik, K., Hothorn, T.,
Huber, W., Iacus, S., Irizarry, R., Leisch, F., Li, C., Maechler, M., Rossini,
A. J., Sawitzki, G., Smith, C., Smyth, G., Tierney, L., Yang, J. Y. H. and
Zhang, J.** (2004). Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol* **5**, R80.
- Giraldez, A. J., Cinalli, R. M., Glasner, M. E., Enright, A. J., Thomson,
J. M., Baskerville, S., Hammond, S. M., Bartel, D. P. and Schier, A. F.** (2005). MicroRNAs regulate brain morphogenesis in zebrafish. *Science* **308**, 833–8.
- Giraldez, A. J., Mishima, Y., Rihel, J., Grocock, R. J., van Dongen, S.,
Inoue, K., Enright, A. J. and Schier, A. F.** (2006). Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science* **312**, 75–9.
- Götz, M. and Huttner, W.** (2005). The cell biology of neurogenesis. *Nat Rev Mol Cell Biol* **6**, 777–788.
- Griffiths-Jones, S.** (2004). The microRNA Registry. *Nucleic Acids Research* **32**, D109–11.
- Griffiths-Jones, S., Grocock, R. J., van Dongen, S., Bateman, A. and Enright, A. J.** (2006). miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Research* **34**, D140–4.
- Griffiths-Jones, S., Saini, H. K., van Dongen, S. and Enright, A. J.** (2008). miRBase: tools for microRNA genomics. *Nucleic Acids Res* **36**, D154–8.

- Grimson, A., Farh, K. K.-H., Johnston, W. K., Garrett-Engle, P., Lim, L. P. and Bartel, D. P.** (2007). MicroRNA targeting specificity in mammals: determinants beyond seed pairing. *Molecular Cell* **27**, 91–105.
- Grimson, A., Srivastava, M., Fahey, B., Woodcroft, B. J., Chiang, H. R., King, N., Degnan, B. M., Rokhsar, D. S. and Bartel, D. P.** (2008). Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature* **455**, 1193–7.
- Grishok, A., Pasquinelli, A. E., Conte, D., Li, N., Parrish, S., Ha, I., Baillie, D. L., Fire, A., Ruvkun, G. and Mello, C. C.** (2001). Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control *C. elegans* developmental timing. *Cell* **106**, 23–34.
- Grün, D., Wang, Y.-L., Langenberger, D., Gunsalus, K. C. and Rajewsky, N.** (2005). microRNA target predictions across seven *Drosophila* species and comparison to mammalian targets. *PLoS Comput Biol* **1**, e13.
- Gu, S., Jin, L., Zhang, F., Sarnow, P. and Kay, M. A.** (2009). Biological basis for restriction of microRNA targets to the 3' untranslated region in mammalian mRNAs. *Nat Struct Mol Biol* **16**, 144–50.
- Guo, H., Ingolia, N. T., Weissman, J. S. and Bartel, D. P.** (2010). Mammalian microRNAs predominantly act to decrease target mRNA levels. *Nature* **466**, 835–40.
- Guo, S. and Kemphues, K. J.** (1995). par-1, a gene required for establishing polarity in *C. elegans* embryos, encodes a putative Ser/Thr kinase that is asymmetrically distributed. *Cell* **81**, 611–20.
- Hamilton, A. J. and Baulcombe, D. C.** (1999). A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science* **286**, 950–2.
- Hammond, S. M., Bernstein, E., Beach, D. and Hannon, G. J.** (2000). An RNA-directed nuclease mediates post-transcriptional gene silencing in *Drosophila* cells. *Nature* **404**, 293–6.
- Han, J., Lee, Y., Yeom, K.-H., Kim, Y.-K., Jin, H. and Kim, V. N.** (2004). The Drosha-DGCR8 complex in primary microRNA processing. *Genes & Development* **18**, 3016–27.

- Hanina, S. A., Mifsud, W., Down, T. A., Hayashi, K., O'Carroll, D., Lao, K., Miska, E. A. and Surani, M. A.** (2010). Genome-Wide Identification of Targets and Function of Individual MicroRNAs in Mouse Embryonic Stem Cells. *PLoS Genetics* **6**, e1001163.
- Hardingham, G. E., Arnold, F. J. and Bading, H.** (2001). Nuclear calcium signaling controls CREB-mediated gene expression triggered by synaptic activity. *Nat Neurosci* **4**, 261–7.
- Hébert, S. S., Papadopoulou, A. S., Smith, P., Galas, M.-C., Planel, E., Silahtaroglu, A. N., Sergeant, N., Buée, L. and Strooper, B. D.** (2010). Genetic ablation of Dicer in adult forebrain neurons results in abnormal tau hyperphosphorylation and neurodegeneration. *Hum Mol Genet* **19**, 3959–69.
- Hendrickson, D. G., Hogan, D. J., McCullough, H. L., Myers, J. W., Herschlag, D., Ferrell, J. E. and Brown, P. O.** (2009). Concordant regulation of translation and mRNA abundance for hundreds of targets of a human microRNA. *PLoS Biol* **7**, e1000238.
- Henke, J. I., Goergen, D., Zheng, J., Song, Y., Schüttler, C. G., Fehr, C., Jünemann, C. and Niepmann, M.** (2008). microRNA-122 stimulates translation of hepatitis C virus RNA. *EMBO J* **27**, 3300–10.
- Herranz, H. and Cohen, S. M.** (2010). MicroRNAs and gene regulatory networks: managing the impact of noise in biological systems. *Genes & Development* **24**, 1339–44.
- Höck, J. and Meister, G.** (2008). The Argonaute protein family. *Genome Biol* **9**, 210.
- Hornstein, E. and Shomron, N.** (2006). Canalization of development by microRNAs. *Nat Genet* **38 Suppl**, S20–4.
- Huang, J. C., Babak, T., Corson, T. W., Chua, G., Khan, S., Gallie, B. L., Hughes, T. R., Blencowe, B. J., Frey, B. J. and Morris, Q. D.** (2007). Using expression profiling data to identify human microRNA targets. *Nat Meth* **4**, 1045–9.
- Hubbard, T. J. P., Aken, B. L., Ayling, S., Ballester, B., Beal, K., Bragin, E., Brent, S., Chen, Y., Clapham, P., Clarke, L., Coates, G., Fairley, S., Fitzgerald, S., Fernandez-Banet, J., Gordon, L., Graf, S., Haider, S., Hammond, M., Holland, R., Howe, K., Jenkinson, A., Johnson, N., Kahari, A.,**

- Keefe, D., Keenan, S., Kinsella, R., Kokocinski, F., Kulesha, E., Lawson, D., Longden, I., Megy, K., Meidl, P., Overduin, B., Parker, A., Pritchard, B., Rios, D., Schuster, M., Slater, G., Smedley, D., Spooner, W., Spudich, G., Trevanion, S., Vilella, A., Vogel, J., White, S., Wilder, S., Zadissa, A., Birney, E., Cunningham, F., Curwen, V., Durbin, R., Fernandez-Suarez, X. M., Herrero, J., Kasprzyk, A., Proctor, G., Smith, J., Searle, S. and Flliceck, P. (2009). Ensembl 2009. *Nucleic acids research* **37**, D690–7.
- Humphreys, D. T., Westman, B. J., Martin, D. I. K. and Preiss, T. (2005). MicroRNAs control translation initiation by inhibiting eukaryotic initiation factor 4E/cap and poly(A) tail function. *Proc Natl Acad Sci USA* **102**, 16961–6.
- Hutvágner, G., McLachlan, J., Pasquinelli, A. E., Bálint, E., Tuschl, T. and Zamore, P. D. (2001). A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA. *Science* **293**, 834–8.
- Hutvágner, G. and Zamore, P. D. (2002). A microRNA in a multiple-turnover RNAi enzyme complex. *Science* **297**, 2056–60.
- Inui, M., Martello, G. and Piccolo, S. (2010). MicroRNA control of signal transduction. *Nat Rev Mol Cell Biol* **11**, 252–63.
- Irizarry, R. A., Bolstad, B. M., Collin, F., Cope, L. M., Hobbs, B. and Speed, T. P. (2003a). Summaries of Affymetrix GeneChip probe level data. *Nucleic Acids Research* **31**, e15.
- Irizarry, R. A., Hobbs, B., Collin, F., Beazer-Barclay, Y. D., Antonellis, K. J., Scherf, U. and Speed, T. P. (2003b). Exploration, normalization, and summaries of high density oligonucleotide array probe level data. *Biostatistics* **4**, 249–64.
- Joshua-Tor, L. and Hannon, G. J. (2010). Ancestral Roles of Small RNAs: An Ago-Centric Perspective. *Cold Spring Harbor perspectives in biology* **10**, a003772.
- Judson, R. L., Babiarz, J. E., Venere, M. and Blelloch, R. (2009). Embryonic stem cell-specific microRNAs promote induced pluripotency. *Nat Biotechnol* **27**, 459–61.
- Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T., Kawashima, S., Okuda, S., Tokimatsu, T. and Yamanashi, Y. (2000). KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic acids research* **28**, 27–30.

- Kanehisa, M., Araki, M., Goto, S., Hattori, M., Hirakawa, M., Itoh, M., Katayama, T., Kawashima, S., Okuda, S., Tokimatsu, T. and Yamanashi, Y.** (2008). KEGG for linking genomes to life and the environment. *Nucleic acids research* **36**, D480–4.
- Kanellopoulou, C., Muljo, S. A., Kung, A. L., Ganesan, S., Drapkin, R., Jenuwein, T., Livingston, D. M. and Rajewsky, K.** (2005). Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes & Development* **19**, 489–501.
- Karginov, F. V., Conaco, C., Xuan, Z., Schmidt, B. H., Parker, J. S., Mandel, G. and Hannon, G. J.** (2007). A biochemical approach to identifying microRNA targets. *Proc Natl Acad Sci USA* **104**, 19291–6.
- Kennerdell, J. R. and Carthew, R. W.** (1998). Use of dsRNA-mediated genetic interference to demonstrate that frizzled and frizzled 2 act in the wingless pathway. *Cell* **95**, 1017–26.
- Ketting, R. F., Fischer, S. E., Bernstein, E., Sijen, T., Hannon, G. J. and Plasterk, R. H.** (2001). Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes & Development* **15**, 2654–9.
- Khvorova, A., Reynolds, A. and Jayasena, S. D.** (2003). Functional siRNAs and miRNAs exhibit strand bias. *Cell* **115**, 209–16.
- Kim, J., Inoue, K., Ishii, J., Vanti, W. B., Voronov, S. V., Murchison, E., Hannon, G. and Abeliovich, A.** (2007). A MicroRNA feedback circuit in midbrain dopamine neurons. *Science* **317**, 1220–4.
- Kim, V. N., Han, J. and Siomi, M. C.** (2009). Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol* **10**, 126–39.
- Klein, M. E., Lioy, D. T., Ma, L., Impey, S., Mandel, G. and Goodman, R. H.** (2007). Homeostatic regulation of MeCP2 expression by a CREB-induced microRNA. *Nat Neurosci* **10**, 1513–4.
- Knight, S. W. and Bass, B. L.** (2001). A role for the RNase III enzyme DCR-1 in RNA interference and germ line development in *Caenorhabditis elegans*. *Science* **293**, 2269–71.

- Konopka, W., Kiryk, A., Novak, M., Herwerth, M., Parkitna, J. R., Wawrzyńiak, M., Kowarsch, A., Michaluk, P., Dzwonek, J., Arnsperger, T., Wilczynski, G., Merkenschlager, M., Theis, F. J., Köhr, G., Kaczmarek, L. and Schütz, G.** (2010). MicroRNA Loss Enhances Learning and Memory in Mice. *J Neurosci* **30**, 14835–14842.
- Kriegstein, A. R. and Dichter, M. A.** (1983). Morphological classification of rat cortical neurons in cell culture. *J Neurosci* **3**, 1634–47.
- Krützfeldt, J., Rajewsky, N., Braich, R., Rajeev, K. G., Tuschl, T., Manoharan, M. and Stoffel, M.** (2005). Silencing of microRNAs in vivo with ‘antagomirs’. *Nature* **438**, 685–689.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W. and Tuschl, T.** (2001). Identification of novel genes coding for small expressed RNAs. *Science* **294**, 853–8.
- Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W. and Tuschl, T.** (2002). Identification of tissue-specific microRNAs from mouse. *Curr Biol* **12**, 735–9.
- Lai, E. C.** (2002). Micro RNAs are complementary to 3' UTR sequence motifs that mediate negative post-transcriptional regulation. *Nat Genet* **30**, 363–4.
- Lai, E. C., Tam, B. and Rubin, G. M.** (2005). Pervasive regulation of Drosophila Notch target genes by GY-box-, Brd-box-, and K-box-class microRNAs. *Genes & Development* **19**, 1067–80.
- Landgraf, P., Rusu, M., Sheridan, R., Sewer, A., Iovino, N., Aravin, A., Pfeffer, S., Rice, A., Kamphorst, A. O., Landthaler, M., Lin, C., Socci, N. D., Hermida, L., Fulci, V., Chiaretti, S., Foà, R., Schliwka, J., Fuchs, U., Novosel, A., Müller, R.-U., Schermer, B., Bissels, U., Inman, J., Phan, Q., Chien, M., Weir, D. B., Choksi, R., Vita, G. D., Frezzetti, D., Trompeter, H.-I., Hornung, V., Teng, G., Hartmann, G., Palkovits, M., Lauro, R. D., Wernet, P., Macino, G., Rogler, C. E., Nagle, J. W., Ju, J., Papavasiliou, F. N., Benzing, T., Lichter, P., Tam, W., Brownstein, M. J., Bosio, A., Borkhardt, A., Russo, J. J., Sander, C., Zavolan, M. and Tuschl, T.** (2007). A mammalian microRNA expression atlas based on small RNA library sequencing. *Cell* **129**, 1401–14.

- Landthaler, M., Gaidatzis, D., Rothballer, A., Chen, P. Y., Soll, S. J., Dinic, L., Ojo, T., Hafner, M., Zavolan, M. and Tuschl, T.** (2008). Molecular characterization of human Argonaute-containing ribonucleoprotein complexes and their bound target mRNAs. *RNA (New York, NY)* **14**, 2580–96.
- Lattin, J. E., Schroder, K., Su, A. I., Walker, J. R., Zhang, J., Wiltshire, T., Saijo, K., Glass, C. K., Hume, D. A., Kellie, S. and Sweet, M. J.** (2008). Expression analysis of G Protein-Coupled Receptors in mouse macrophages. *Immunome Res* **4**, 5.
- Le, M. T. N., Xie, H., Zhou, B., Chia, P. H., Rizk, P., Um, M., Udolph, G., Yang, H., Lim, B. and Lodish, H. F.** (2009). MicroRNA-125b promotes neuronal differentiation in human cells by repressing multiple targets. *Mol Cell Biol* **29**, 5290–305.
- Lee, M. K., Tuttle, J. B., Rebhun, L. I., Cleveland, D. W. and Frankfurter, A.** (1990). The expression and posttranslational modification of a neuron-specific beta-tubulin isotype during chick embryogenesis. *Cell Motil Cytoskeleton* **17**, 118–32.
- Lee, R. C., Feinbaum, R. L. and Ambros, V.** (1993). The *C. elegans* heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. *Cell* **75**, 843–54.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Rådmark, O., Kim, S. and Kim, V. N.** (2003). The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**, 415–9.
- Lee, Y., Jeon, K., Lee, J.-T., Kim, S. and Kim, V. N.** (2002). MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J* **21**, 4663–70.
- Lee, Y., Kim, M., Han, J., Yeom, K.-H., Lee, S., Baek, S. H. and Kim, V. N.** (2004). MicroRNA genes are transcribed by RNA polymerase II. *EMBO J* **23**, 4051–60.
- Leucht, C., Stigloher, C., Wizenmann, A., Klafke, R., Folchert, A. and Bally-Cuif, L.** (2008). MicroRNA-9 directs late organizer activity of the midbrain-hindbrain boundary. *Nat Neurosci* **11**, 641–8.
- Leung, A. K. L., Calabrese, J. M. and Sharp, P. A.** (2006). Quantitative analysis of Argonaute protein reveals microRNA-dependent localization to stress granules. *Proc Natl Acad Sci USA* **103**, 18125–30.

- Leung, A. K. L. and Sharp, P. A.** (2006). Function and localization of microRNAs in mammalian cells. *Cold Spring Harb Symp Quant Biol* **71**, 29–38.
- Leung, A. K. L. and Sharp, P. A.** (2010). MicroRNA Functions in Stress Responses. *Molecular Cell* **40**, 205–15.
- Leung, A. K. L., Young, A. G., Bhutkar, A., Zheng, G. X., Bosson, A. D., Nielsen, C. B. and Sharp, P. A.** (2011). Genome-wide identification of Ago2 binding sites from mouse embryonic stem cells with and without mature microRNAs. *Nat Struct Mol Biol* **18**, 237–44.
- Leuschner, P. J. F., Ameres, S. L., Kueng, S. and Martinez, J.** (2006). Cleavage of the siRNA passenger strand during RISC assembly in human cells. *EMBO Rep* **7**, 314–20.
- Lewis, A. and Redrup, L.** (2005). Genetic imprinting: conflict at the Callipyge locus. *Curr Biol* **15**, R291–4.
- Lewis, B. P., Burge, C. B. and Bartel, D. P.** (2005). Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* **120**, 15–20.
- Lewis, B. P., hung Shih, I., Jones-Rhoades, M. W., Bartel, D. P. and Burge, C. B.** (2003). Prediction of mammalian microRNA targets. *Cell* **115**, 787–98.
- Lewis, M. A., Quint, E., Glazier, A. M., Fuchs, H., Angelis, M. H. D., Langford, C., van Dongen, S., Abreu-Goodger, C., Piipari, M., Redshaw, N., Dalmay, T., Moreno-Pelayo, M. A., Enright, A. J. and Steel, K. P.** (2009). An ENU-induced mutation of miR-96 associated with progressive hearing loss in mice. *Nat Genet* **41**, 614–8.
- Li, X., Cassidy, J. J., Reinke, C. A., Fischboeck, S. and Carthew, R. W.** (2009). A microRNA imparts robustness against environmental fluctuation during development. *Cell* **137**, 273–82.
- Li, X. and Jin, P.** (2010). Roles of small regulatory RNAs in determining neuronal identity. *Nat Rev Neurosci* **11**, 329–38.
- Licatalosi, D. D., Mele, A., Fak, J. J., Ule, J., Kayikci, M., Chi, S. W., Clark, T. A., Schweitzer, A. C., Blume, J. E., Wang, X., Darnell, J. C. and**

- Darnell, R. B.** (2008). HITS-CLIP yields genome-wide insights into brain alternative RNA processing. *Nature* **456**, 464–9.
- Lim, L. P., Lau, N. C., Garrett-Engle, P., Grimson, A., Schelter, J. M., Castle, J., Bartel, D. P., Linsley, P. S. and Johnson, J. M.** (2005). Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* **433**, 769–73.
- Lim, L. P., Lau, N. C., Weinstein, E. G., Abdelhakim, A., Yekta, S., Rhoades, M. W., Burge, C. B. and Bartel, D. P.** (2003). The microRNAs of *Caenorhabditis elegans*. *Genes & Development* **17**, 991–1008.
- Lin, S. M., Du, P., Huber, W. and Kibbe, W. A.** (2007). Model-based variance-stabilizing transformation for Illumina microarray data. *Nucleic Acids Research* **36**, e11–e11.
- Liu, J., Carmell, M. A., Rivas, F. V., Marsden, C. G., Thomson, J. M., Song, J.-J., Hammond, S. M., Joshua-Tor, L. and Hannon, G. J.** (2004). Argonaute2 is the catalytic engine of mammalian RNAi. *Science* **305**, 1437–41.
- Liu, J., Valencia-Sanchez, M. A., Hannon, G. J. and Parker, R.** (2005). MicroRNA-dependent localization of targeted mRNAs to mammalian P-bodies. *Nat Cell Biol* **7**, 719–23.
- Lotterman, C. D., Kent, O. A. and Mendell, J. T.** (2008). Functional integration of microRNAs into oncogenic and tumor suppressor pathways. *Cell Cycle* **7**, 2493–9.
- Lu, T., Pan, Y., Kao, S.-Y., Li, C., Kohane, I., Chan, J. and Yankner, B. A.** (2004). Gene regulation and DNA damage in the ageing human brain. *Nature* **429**, 883–91.
- Lugli, G., Torvik, V. I., Larson, J. and Smalheiser, N. R.** (2008). Expression of microRNAs and their precursors in synaptic fractions of adult mouse forebrain. *Journal of Neurochemistry* **106**, 650–61.
- MacLaren, E. J., Charlesworth, P., Coba, M. P. and Grant, S. G. N.** (2011). Knockdown of mental disorder susceptibility genes disrupts neuronal network physiology in vitro. *Molecular and cellular neurosciences* .

- Makeyev, E. V., Zhang, J., Carrasco, M. A. and Maniatis, T.** (2007). The MicroRNA miR-124 promotes neuronal differentiation by triggering brain-specific alternative pre-mRNA splicing. *Molecular Cell* **27**, 435–48.
- Malone, C. D. and Hannon, G. J.** (2009). Small RNAs as guardians of the genome. *Cell* **136**, 656–68.
- Manakov, S., Grant, S. and Enright, A.** (2009). Reciprocal regulation of microRNA and mRNA profiles in neuronal development and synapse formation. *BMC Genomics* **10**, 419.
- Maniataki, E. and Mourelatos, Z.** (2005). A human, ATP-independent, RISC assembly machine fueled by pre-miRNA. *Genes & Development* **19**, 2979–90.
- Maroney, P. A., Yu, Y., Fisher, J. and Nilsen, T. W.** (2006). Evidence that microRNAs are associated with translating messenger RNAs in human cells. *Nat Struct Mol Biol* **13**, 1102–7.
- Mathonnet, G., Fabian, M. R., Svitkin, Y. V., Parsyan, A., Huck, L., Murata, T., Biffo, S., Merrick, W. C., Darzynkiewicz, E., Pillai, R. S., Filipowicz, W., Duchaine, T. F. and Sonenberg, N.** (2007). MicroRNA inhibition of translation initiation in vitro by targeting the cap-binding complex eIF4F. *Science* **317**, 1764–7.
- Matranga, C., Tomari, Y., Shin, C., Bartel, D. P. and Zamore, P. D.** (2005). Passenger-strand cleavage facilitates assembly of siRNA into Ago2-containing RNAi enzyme complexes. *Cell* **123**, 607–20.
- Miska, E. A., Alvarez-Saavedra, E., Abbott, A. L., Lau, N. C., Hellman, A. B., McGonagle, S. M., Bartel, D. P., Ambros, V. R. and Horvitz, H. R.** (2007). Most *Caenorhabditis elegans* microRNAs are individually not essential for development or viability. *PLoS Genet* **3**, e215.
- Napoli, C., Lemieux, C. and Jorgensen, R.** (1990). Introduction of a Chimeric Chalcone Synthase Gene into Petunia Results in Reversible Co-Suppression of Homologous Genes in trans. *Plant Cell* **2**, 279–289.
- Nelson, P. G.** (1975). Nerve and muscle cells in culture. *Physiol Rev* **55**, 1–61.
- Ning, Z., Cox, A. J. and Mullikin, J. C.** (2001). SSAHA: a fast search method for large DNA databases. *Genome Res* **11**, 1725–9.

- Nottrott, S., Simard, M. J. and Richter, J. D.** (2006). Human let-7a miRNA blocks protein production on actively translating polyribosomes. *Nat Struct Mol Biol* **13**, 1108–14.
- Oelgeschläger, M., Larraín, J., Geissert, D. and Robertis, E. M. D.** (2000). The evolutionarily conserved BMP-binding protein Twisted gastrulation promotes BMP signalling. *Nature* **405**, 757–63.
- Olsen, P. H. and Ambros, V.** (1999). The lin-4 regulatory RNA controls developmental timing in *Caenorhabditis elegans* by blocking LIN-14 protein synthesis after the initiation of translation. *Dev Biol* **216**, 671–80.
- Ørom, U. A., Nielsen, F. C. and Lund, A. H.** (2008). MicroRNA-10a binds the 5'UTR of ribosomal protein mRNAs and enhances their translation. *Mol Cell* **30**, 460–71.
- Pasquinelli, A. E., Reinhart, B. J., Slack, F., Martindale, M. Q., Kuroda, M. I., Maller, B., Hayward, D. C., Ball, E. E., Degnan, B., Müller, P., Spring, J., Srinivasan, A., Fishman, M., Finnerty, J., Corbo, J., Levine, M., Leahy, P., Davidson, E. and Ruvkun, G.** (2000). Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA. *Nature* **408**, 86–9.
- Peters, L. and Meister, G.** (2007). Argonaute proteins: mediators of RNA silencing. *Molecular Cell* **26**, 611–23.
- Petersen, C. P., Bordeleau, M.-E., Pelletier, J. and Sharp, P. A.** (2006). Short RNAs repress translation after initiation in mammalian cells. *Molecular Cell* **21**, 533–42.
- Pillai, R. S., Artus, C. G. and Filipowicz, W.** (2004). Tethering of human Ago proteins to mRNA mimics the miRNA-mediated repression of protein synthesis. *RNA (New York, NY)* **10**, 1518–25.
- Pillai, R. S., Bhattacharyya, S. N., Artus, C. G., Zoller, T., Cougot, N., Basuyk, E., Bertrand, E. and Filipowicz, W.** (2005). Inhibition of translational initiation by Let-7 MicroRNA in human cells. *Science* **309**, 1573–6.
- Poliseno, L., Salmena, L., Riccardi, L., Fornari, A., Song, M. S., Hobbs, R. M., Sportoletti, P., Varmeh, S., Egia, A., Fedele, G., Rameh, L., Loda, M. and**

- Pandolfi, P. P.** (2010). Identification of the miR-106b 25 microRNA cluster as a proto-oncogenic PTEN-targeting intron that cooperates with its host gene MCM7 in transformation. *Sci Signal* **3**, ra29.
- Pruitt, K. D., Tatusova, T., Klimke, W. and Maglott, D. R.** (2009). NCBI Reference Sequences: current status, policy and new initiatives. *Nucleic acids research* **37**, D32–6.
- Qi, Y., Wang, J. K., McMillian, M. and Chikaraishi, D. M.** (1997). Characterization of a CNS cell line, CAD, in which morphological differentiation is initiated by serum deprivation. *J Neurosci* **17**, 1217–25.
- Quintavalle, M., Elia, L., Condorelli, G. and Courtneidge, S. A.** (2010). MicroRNA control of podosome formation in vascular smooth muscle cells in vivo and in vitro. *J Cell Biol* **189**, 13–22.
- Rajasethupathy, P., Fiumara, F., Sheridan, R., Betel, D., Puthanveettil, S. V., Russo, J. J., Sander, C., Tuschl, T. and Kandel, E.** (2009). Characterization of small RNAs in aplysia reveals a role for miR-124 in constraining synaptic plasticity through CREB. *Neuron* **63**, 803–17.
- Rao, Y., Lee, Y., Jarjoura, D., Ruppert, A. S., Liu, C.-G., Hsu, J. C. and Hagan, J. P.** (2008). A comparison of normalization techniques for microRNA microarray data. *Statistical Applications in Genetics and Molecular Biology* **7**, Article22.
- Reinhart, B. J., Slack, F. J., Basson, M., Pasquinelli, A. E., Bettinger, J. C., Rougvie, A. E., Horvitz, H. R. and Ruvkun, G.** (2000). The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* **403**, 901–6.
- Rodriguez, A., Vigorito, E., Clare, S., Warren, M. V., Couttet, P., Soond, D. R., Dongen, S. V., Grocock, R. J., Das, P. P., Miska, E. A., Vetrie, D., Okkenhaug, K., Enright, A. J., Dougan, G., Turner, M. and Bradley, A.** (2007). Requirement of bic/microRNA-155 for Normal Immune Function. *Science* **316**, 608–611.
- Romano, N. and Macino, G.** (1992). Quelling: transient inactivation of gene expression in *Neurospora crassa* by transformation with homologous sequences. *Mol Microbiol* **6**, 3343–53.

- RTeam** (2008). R: A language and environment for statistical computing. *R Foundation for Statistical Computing Vienna Austria.*
- Rybak, A., Fuchs, H., Smirnova, L., Brandt, C., Pohl, E. E., Nitsch, R. and Wulczyn, F. G.** (2008). A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* **10**, 987–93.
- Sayers, E. W., Barrett, T., Benson, D. A., Bolton, E., Bryant, S. H., Canese, K., Chetvernin, V., Church, D. M., Dicuccio, M., Federhen, S., Feolo, M., Fingerman, I. M., Geer, L. Y., Helberg, W., Kapustin, Y., Landsman, D., Lipman, D. J., Lu, Z., Madden, T. L., Madej, T., Maglott, D. R., Marchler-Bauer, A., Miller, V., Mizrahi, I., Ostell, J., Panchenko, A., Phan, L., Pruitt, K. D., Schuler, G. D., Sequeira, E., Sherry, S. T., Shumway, M., Sirotkin, K., Slotta, D., Souvorov, A., Starchenko, G., Tatusova, T. A., Wagner, L., Wang, Y., Wilbur, W. J., Yaschenko, E. and Ye, J.** (2010). Database resources of the National Center for Biotechnology Information. *Nucleic acids research* **39**, D38–51.
- Schaefer, A., O'Carroll, D., Tan, C. L., Hillman, D., Sugimori, M., Llinas, R. and Greengard, P.** (2007). Cerebellar neurodegeneration in the absence of microRNAs. *J Exp Med* **204**, 1553–8.
- Scherer, W., Syverton, J. and Gey, G.** (1953). Studies on the propagation in vitro of poliomyelitis viruses. IV. Viral multiplication in a stable strain of human malignant epithelial cells (strain HeLa) derived from an epidermoid carcinoma of the cervix. *J Exp Med* **97**, 695–710.
- Schratt, G.** (2009). microRNAs at the synapse. *Nat Rev Neurosci* **10**, 842–9.
- Schratt, G. M., Tuebing, F., Nigh, E. A., Kane, C. G., Sabatini, M. E., Kiebler, M. and Greenberg, M. E.** (2006). A brain-specific microRNA regulates dendritic spine development. *Nature* **439**, 283–9.
- Schroeder, A., Levins, C. G., Cortez, C., Langer, R. and Anderson, D. G.** (2010). Lipid-based nanotherapeutics for siRNA delivery. *J Intern Med* **267**, 9–21.
- Schwarz, D. S., Hutváigner, G., Du, T., Xu, Z., Aronin, N. and Zamore, P. D.** (2003). Asymmetry in the assembly of the RNAi enzyme complex. *Cell* **115**, 199–208.

- Seitz, H., Royo, H., Bortolin, M.-L., Lin, S.-P., Ferguson-Smith, A. C. and Cavaillé, J.** (2004). A large imprinted microRNA gene cluster at the mouse Dlk1-Gtl2 domain. *Genome Res* **14**, 1741–8.
- Selbach, M., Schwahnhäusser, B., Thierfelder, N., Fang, Z., Khanin, R. and Rajewsky, N.** (2008). Widespread changes in protein synthesis induced by microRNAs. *Nature* **455**, 58–63.
- Sempere, L. F., Cole, C. N., McPeek, M. A. and Peterson, K. J.** (2006). The phylogenetic distribution of metazoan microRNAs: insights into evolutionary complexity and constraint. *J Exp Zool B Mol Dev Evol* **306**, 575–88.
- Sharp, P. A.** (2009). The centrality of RNA. *Cell* **136**, 577–80.
- Shibata, M., Kurokawa, D., Nakao, H., Ohmura, T. and Aizawa, S.** (2008). MicroRNA-9 modulates Cajal-Retzius cell differentiation by suppressing Foxg1 expression in mouse medial pallium. *J Neurosci* **28**, 10415–21.
- Shkumatava, A., Stark, A., Sive, H. and Bartel, D. P.** (2009). Coherent but overlapping expression of microRNAs and their targets during vertebrate development. *Genes & Development* **23**, 466–81.
- Siegal, M. L. and Bergman, A.** (2002). Waddington's canalization revisited: developmental stability and evolution. *Proc Natl Acad Sci USA* **99**, 10528–32.
- Siegel, G., Obernosterer, G., Fiore, R., Oehmen, M., Bicker, S., Christensen, M., Khudayberdiev, S., Leuschner, P. F., Busch, C. J. L., Kane, C., Hübel, K., Dekker, F., Hedberg, C., Rengarajan, B., Drepper, C., Waldmann, H., Kauppinen, S., Greenberg, M. E., Draguhn, A., Rehmsmeier, M., Martinez, J. and Schratt, G. M.** (2009). A functional screen implicates microRNA-138-dependent regulation of the depalmitoylation enzyme APT1 in dendritic spine morphogenesis. *Nature* **11**, 705–16.
- Silber, J., Lim, D. A., Petritsch, C., Persson, A. I., Maunakea, A. K., Yu, M., Vandenberg, S. R., Ginzinger, D. G., James, C. D., Costello, J. F., Bergers, G., Weiss, W. A., Alvarez-Buylla, A. and Hodgson, J. G.** (2008). miR-124 and miR-137 inhibit proliferation of glioblastoma multiforme cells and induce differentiation of brain tumor stem cells. *BMC medicine* **6**, 14.

- Smith, C. J., Watson, C. F., Bird, C. R., Ray, J., Schuch, W. and Grierson, D.** (1990). Expression of a truncated tomato polygalacturonase gene inhibits expression of the endogenous gene in transgenic plants. *Mol Gen Genet* **224**, 477–81.
- Smyth, G.** (2004). Linear Models and Empirical Bayes Methods for Assessing Differential Expression in Microarray Experiments. *Statistical Applications in Genetics and Molecular Biology* **3**, Article 3.
- Sood, P., Krek, A., Zavolan, M., Macino, G. and Rajewsky, N.** (2006). Cell-type-specific signatures of microRNAs on target mRNA expression. *Proc Natl Acad Sci USA* **103**, 2746–51. Notes.
- Stark, A., Brennecke, J., Bushati, N., Russell, R. B. and Cohen, S. M.** (2005). Animal MicroRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell* **123**, 1133–46.
- Stark, A., Brennecke, J., Russell, R. B. and Cohen, S. M.** (2003). Identification of Drosophila MicroRNA targets. *PLoS Biol* **1**, E60.
- Svoboda, P., Stein, P., Hayashi, H. and Schultz, R. M.** (2000). Selective reduction of dormant maternal mRNAs in mouse oocytes by RNA interference. *Development* **127**, 4147–56.
- Taft, R. J., Glazov, E. A., Cloonan, N., Simons, C., Stephen, S., Faulkner, G. J., Lassmann, T., Forrest, A. R. R., Grimmond, S. M., Schroder, K., Irvine, K., Arakawa, T., Nakamura, M., Kubosaki, A., Hayashida, K., Kawazu, C., Murata, M., Nishiyori, H., Fukuda, S., Kawai, J., Daub, C. O., Hume, D. A., Suzuki, H., Orlando, V., Carninci, P., Hayashizaki, Y. and Mattick, J. S.** (2009). Tiny RNAs associated with transcription start sites in animals. *Nat Genet* **41**, 572–8.
- Taft, R. J., Simons, C., Nahkuri, S., Oey, H., Korbie, D. J., Mercer, T. R., Holst, J., Ritchie, W., Wong, J. J.-L., Rasko, J. E. J., Rokhsar, D. S., Degnan, B. M. and Mattick, J. S.** (2010). Nuclear-localized tiny RNAs are associated with transcription initiation and splice sites in metazoans. *Nat Struct Mol Biol* **17**, 1030–4.
- Thermann, R. and Hentze, M. W.** (2007). Drosophila miR2 induces pseudo-polysomes and inhibits translation initiation. *Nature* **447**, 875–8.

- Thomas-Chollier, M., Sand, O., Turatsinze, J.-V., Janky, R., Defrance, M., Vervisch, E., Brohée, S. and van Helden, J.** (2008). RSAT: regulatory sequence analysis tools. *Nucleic acids research* **36**, W119–27.
- Thomson, J. M., Newman, M., Parker, J. S., Morin-Kensicki, E. M., Wright, T. and Hammond, S. M.** (2006). Extensive post-transcriptional regulation of microRNAs and its implications for cancer. *Genes & Development* **20**, 2202–7.
- Valor, L. M., Charlesworth, P., Humphreys, L., Anderson, C. N. G. and Grant, S. G. N.** (2007). Network activity-independent coordinated gene expression program for synapse assembly. *Proc Natl Acad Sci USA* **104**, 4658–63.
- van der Krol, A. R., Mur, L. A., Beld, M., Mol, J. N. and Stuitje, A. R.** (1990). Flavonoid genes in petunia: addition of a limited number of gene copies may lead to a suppression of gene expression. *Plant Cell* **2**, 291–9.
- van Dongen, S.** (2000). Graph clustering by flow simulation. *PhD thesis University of Utrecht*.
- van Dongen, S., Abreu-Goodger, C. and Enright, A. J.** (2008). Detecting microRNA binding and siRNA off-target effects from expression data. *Nat Meth* **5**, 1023–5.
- van Rooij, E., Sutherland, L. B., Qi, X., Richardson, J. A., Hill, J. and Olson, E. N.** (2007). Control of stress-dependent cardiac growth and gene expression by a microRNA. *Science* **316**, 575–9.
- Vasudevan, S., Tong, Y. and Steitz, J. A.** (2007). Switching from repression to activation: microRNAs can up-regulate translation. *Science* **318**, 1931–4.
- Visvanathan, J., Lee, S., Lee, B., Lee, J. W. and Lee, S.-K.** (2007). The microRNA miR-124 antagonizes the anti-neural REST/SCP1 pathway during embryonic CNS development. *Genes & Development* **21**, 744–9.
- Voinnet, O.** (2009). Origin, biogenesis, and activity of plant microRNAs. *Cell* **136**, 669–87.
- Volinia, S., Galasso, M., Costinean, S., Tagliavini, L., Gamberoni, G., Drusco, A., Marchesini, J., Mascellani, N., Sana, M. E., Jarour, R. A., Desponts, C., Teitell, M., Baffa, R., Aqeilan, R., Iorio, M. V., Taccioli, C., Garzon,**

- R., Leva, G. D., Fabbri, M., Catozzi, M., Previati, M., Ambs, S., Palumbo, T., Garofalo, M., Veronese, A., Bottoni, A., Gasparini, P., Harris, C. C., Visone, R., Pekarsky, Y., de la Chapelle, A., Bloomston, M., Dillhoff, M., Rassenti, L. Z., Kipps, T. J., Huebner, K., Pichiorri, F., Lenze, D., Cairo, S., Buendia, M.-A., Pineau, P., Dejean, A., Zanesi, N., Rossi, S., Calin, G. A., Liu, C.-G., Palatini, J., Negrini, M., Vecchione, A., Rosenberg, A. and Croce, C. M. (2010). Reprogramming of miRNA networks in cancer and leukemia. *Genome Res* **20**, 589–99.
- Wagenaar, D. A., Pine, J. and Potter, S. M. (2006). Searching for plasticity in dissociated cortical cultures on multi-electrode arrays. *J Negat Results BioMed* **5**, 16.
- Wakiyama, M., Takimoto, K., Ohara, O. and Yokoyama, S. (2007). Let-7 microRNA-mediated mRNA deadenylation and translational repression in a mammalian cell-free system. *Genes & Development* **21**, 1857–62.
- Wang, B., Love, T. M., Call, M. E., Doench, J. G. and Novina, C. D. (2006). Recapitulation of short RNA-directed translational gene silencing in vitro. *Molecular Cell* **22**, 553–60.
- Wang, B., Yanez, A. and Novina, C. D. (2008a). MicroRNA-repressed mRNAs contain 40S but not 60S components. *Proc Natl Acad Sci USA* **105**, 5343–8.
- Wang, X. and Wang, X. (2006). Systematic identification of microRNA functions by combining target prediction and expression profiling. *Nucleic Acids Res* **34**, 1646–52.
- Wang, Y., Juranek, S., Li, H., Sheng, G., Tuschl, T. and Patel, D. J. (2008b). Structure of an argonaute silencing complex with a seed-containing guide DNA and target RNA duplex. *Nature* **456**, 921–6.
- Wang, Y., Medvid, R., Melton, C., Jaenisch, R. and Blelloch, R. (2007). DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nat Genet* **39**, 380–5.
- Wang, Y., Sheng, G., Juranek, S., Tuschl, T. and Patel, D. J. (2008c). Structure of the guide-strand-containing argonaute silencing complex. *Nature* **456**, 209–13.
- Wianny, F. and Zernicka-Goetz, M. (2000). Specific interference with gene function by double-stranded RNA in early mouse development. *Nat Cell Biol* **2**, 70–5.

- Wienholds, E., Kloosterman, W. P., Miska, E., Alvarez-Saavedra, E., Berezikov, E., de Bruijn, E., Horvitz, H. R., Kauppinen, S. and Plasterk, R. H. A.** (2005). MicroRNA expression in zebrafish embryonic development. *Science* **309**, 310–1.
- Wightman, B., Ha, I. and Ruvkun, G.** (1993). Posttranscriptional regulation of the heterochronic gene lin-14 by lin-4 mediates temporal pattern formation in *C. elegans*. *Cell* **75**, 855–62.
- Winter, J., Jung, S., Keller, S., Gregory, R. I. and Diederichs, S.** (2009). Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat Cell Biol* **11**, 228–34.
- Wu, C., Orozco, C., Boyer, J., Leglise, M., Goodale, J., Batalov, S., Hodge, C. L., Haase, J., Janes, J., Huss, J. W. and Su, A. I.** (2009a). BioGPS: an extensible and customizable portal for querying and organizing gene annotation resources. *Genome Biol* **10**, R130.
- Wu, C.-I., Shen, Y. and Tang, T.** (2009b). Evolution under canalization and the dual roles of microRNAs: a hypothesis. *Genome Res* **19**, 734–43.
- Xin, M., Small, E. M., Sutherland, L. B., Qi, X., McAnally, J., Plato, C. F., Richardson, J. A., Bassel-Duby, R. and Olson, E. N.** (2009). MicroRNAs miR-143 and miR-145 modulate cytoskeletal dynamics and responsiveness of smooth muscle cells to injury. *Genes & Development* **23**, 2166–78.
- Xu, P., Vernooy, S. Y., Guo, M. and Hay, B. A.** (2003). The Drosophila microRNA Mir-14 suppresses cell death and is required for normal fat metabolism. *Curr Biol* **13**, 790–5.
- Xu, Y. and Szoka, F. C.** (1996). Mechanism of DNA release from cationic liposome/DNA complexes used in cell transfection. *Biochemistry* **35**, 5616–23.
- Yang, J.-S., Maurin, T., Robine, N., Rasmussen, K. D., Jeffrey, K. L., Chandwani, R., Papapetrou, E. P., Sadelain, M., O'Carroll, D. and Lai, E. C.** (2010). Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc Natl Acad Sci USA* **107**, 15163–8.

- Yi, R., Qin, Y., Macara, I. G. and Cullen, B. R.** (2003). Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes & Development* **17**, 3011–6.
- Yoo, A. S., Staahl, B. T., Chen, L. and Crabtree, G. R.** (2009). MicroRNA-mediated switching of chromatin-remodelling complexes in neural development. *Nature* **460**, 642–6.
- Zabner, J., Fasbender, A. J., Moninger, T., Poellinger, K. A. and Welsh, M. J.** (1995). Cellular and molecular barriers to gene transfer by a cationic lipid. *J Biol Chem* **270**, 18997–9007.
- Zamore, P. D., Tuschl, T., Sharp, P. A. and Bartel, D. P.** (2000). RNAi: double-stranded RNA directs the ATP-dependent cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* **101**, 25–33.
- Zhang, G., Gurtu, V. and Kain, S. R.** (1996). An enhanced green fluorescent protein allows sensitive detection of gene transfer in mammalian cells. *Biochem Biophys Res Commun* **227**, 707–11.
- Zhao, C., Sun, G., Li, S. and Shi, Y.** (2009). A feedback regulatory loop involving microRNA-9 and nuclear receptor TLX in neural stem cell fate determination. *Nat Struct Mol Biol* **16**, 365–71.
- Zhao, X., He, X., Han, X., Yu, Y., Ye, F., Chen, Y., Hoang, T., Xu, X., Mi, Q.-S., Xin, M., Wang, F., Appel, B. and Lu, Q. R.** (2010). MicroRNA-mediated control of oligodendrocyte differentiation. *Neuron* **65**, 612–26.
- Zipprich, J. T., Bhattacharyya, S., Mathys, H. and Filipowicz, W.** (2009). Importance of the C-terminal domain of the human GW182 protein TNRC6C for translational repression. *RNA (New York, NY)* **15**, 781–93.