

References

- Amrani, N., Ganesan, R., Kervestin, S., Mangus, D.A., Ghosh, S. and Jacobson, A. (2004) A faux 3'-UTR promotes aberrant termination and triggers nonsense-mediated mRNA decay. *Nature*, **432**, 112-118.
- Anders, K.R., Grimson, A. and Anderson, P. (2003) SMG-5, required for C.elegans nonsense-mediated mRNA decay, associates with SMG-2 and protein phosphatase 2A. *Embo J*, **22**, 641-650.
- Austin, J. and Kimble, J. (1987) glp-1 is required in the germ line for regulation of the decision between mitosis and meiosis in *C. elegans*. *Cell*, **51**, 589-599.
- Austin, J. and Kimble, J. (1989) Transcript analysis of glp-1 and lin-12, homologous genes required for cell interactions during development of *C. elegans*. *Cell*, **58**, 565-571.
- Azzalin, C. M., Reichenbach, P., Khoriauli, L., Giulotto, E. and Lingner, J. (2007) Telomeric repeat containing RNA and RNA surveillance factors at mammalian chromosome ends. *Science* 318, 798-801.
- Banan, A., Fields, J.Z., Zhang, Y. and Keshavarzian, A. (2001) Key role of PKC and Ca²⁺ in EGF protection of microtubules and intestinal barrier against oxidants. *Am J Physiol Gastrointest Liver Physiol*, **280**, G828-843.
- Beer, M.A. and Tavazoie, S. (2004) Predicting gene expression from sequence. *Cell*, **117**, 185-198.
- Behm-Ansmant, I., Gatfield, D., Rehwinkel, J., Hilgers, V. and Izaurralde, E. (2007a) A conserved role for cytoplasmic poly(A)-binding protein 1 (PABPC1) in nonsense-mediated mRNA decay. *Embo J*, **26**, 1591-1601.
- Behm-Ansmant, I., Kashima, I., Rehwinkel, J., Sauliere, J., Wittkopp, N. and Izaurralde, E. (2007b) mRNA quality control: an ancient machinery recognizes and degrades mRNAs with nonsense codons. *FEBS Lett*, **581**, 2845-2853.
- Bender, A.M., Kirienko, N.V., Olson, S.K., Esko, J.D. and Fay, D.S. (2007) lin-35/Rb and the CoREST ortholog spr-1 coordinately regulate vulval morphogenesis and gonad development in *C. elegans*. *Dev Biol*, **302**, 448-462.
- Berry, L.W., Westlund, B. and Schedl, T. (1997) Germ-line tumor formation caused by activation of glp-1, a *Caenorhabditis elegans* member of the Notch family of receptors. *Development*, **124**, 925-936.
- Bertani, G. (1951). Studies on lysogenesis. I. The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol*, **62**, 293-300.
- Bertone, P., Stolc, V., Royce, T.E., Rozowsky, J.S., Urban, A.E., Zhu, X., Rinn, J.L., Tongprasit, W., Samanta, M., Weissman, S., Gerstein, M. and Snyder, M.

- (2004) Global identification of human transcribed sequences with genome tiling arrays. *Science*, **306**, 2242-2246.
- Blumenthal, T., Evans, D., Link, C.D., Guffanti, A., Lawson, D., Thierry-Mieg, J., Thierry-Mieg, D., Chiu, W.L., Duke, K., Kiraly, M. and Kim, S.K. (2002) A global analysis of *Caenorhabditis elegans* operons. *Nature*, **417**, 851-854.
- Boisvert, M.E. and Simard, M.J. (2008) RNAi pathway in *C. elegans*: the argonautes and collaborators. *Curr Top Microbiol Immunol*, **320**, 21-36.
- Bolstad, B.M., Irizarry, R.A., Astrand, M. and Speed, T.P. (2003) A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics*, **19**, 185-193.
- Booth, E.O., Van Driessche, N., Zhuchenko, O., Kuspa, A. and Shaulsky, G. (2005) Microarray phenotyping in *Dictyostelium* reveals a regulon of chemotaxis genes. *Bioinformatics*, **21**, 4371-4377.
- Brenner, S. (1974) The genetics of *Caenorhabditis elegans*. *Genetics*, **77**, 71-94.
- Buhler, M., Steiner, S., Mohn, F., Paillusson, A. and Muhlemann, O. (2006) EJC-independent degradation of nonsense immunoglobulin-mu mRNA depends on 3' UTR length. *Nat Struct Mol Biol*, **13**, 462-464.
- C. elegans* Sequencing Consortium. (1998) Genome sequence of the nematode *C. elegans*: a platform for investigating biology. *Science*, **282**, 2012-2018.
- Cali, B.M., Kuchma, S.L., Latham, J. and Anderson, P. (1999) smg-7 is required for mRNA surveillance in *Caenorhabditis elegans*. *Genetics*, **151**, 605-616.
- Calvi, L.M., Adams, G.B., Weibrech, K.W., Weber, J.M., Olson, D.P., Knight, M.C., Martin, R.P., Schipani, E., Divieti, P., Bringhurst, F.R., Milner, L.A., Kronenberg, H.M. and Scadden, D.T. (2003) Osteoblastic cells regulate the haematopoietic stem cell niche. *Nature*, **425**, 841-846.
- Chang, C., Hopper, N.A. and Sternberg, P.W. (2000) *Caenorhabditis elegans* SOS-1 is necessary for multiple RAS-mediated developmental signals. *Embo J*, **19**, 3283-3294.
- Chang, J.C. and Kan, Y.W. (1979) beta 0 thalassemia, a nonsense mutation in man. *Proc Natl Acad Sci U S A*, **76**, 2886-2889.
- Chang, Y.F., Imam, J.S. and Wilkinson, M.F. (2007) The nonsense-mediated decay RNA surveillance pathway. *Annu Rev Biochem*, **76**, 51-74.
- Chen, N. and Greenwald, I. (2004) The lateral signal for LIN-12/Notch in *C. elegans* vulval development comprises redundant secreted and transmembrane DSL proteins. *Dev Cell*, **6**, 183-192.
- Chiu, S.Y., Serin, G., Ohara, O. and Maquat, L.E. (2003) Characterization of human Smg5/7a: a protein with similarities to *Caenorhabditis elegans* SMG5 and SMG7 that functions in the dephosphorylation of Upf1. *Rna*, **9**, 77-87.

- Christensen, S., Kodoyianni, V., Bosenberg, M., Friedman, L. and Kimble, J. (1996) lag-1, a gene required for lin-12 and glp-1 signaling in *Caenorhabditis elegans*, is homologous to human CBF1 and *Drosophila* Su(H). *Development*, **122**, 1373-1383.
- Church, D.L., Guan, K.L. and Lambie, E.J. (1995) Three genes of the MAP kinase cascade, mek-2, mpk-1/sur-1 and let-60 ras, are required for meiotic cell cycle progression in *Caenorhabditis elegans*. *Development*, **121**, 2525-2535.
- Crittenden, S.L., Bernstein, D.S., Bachorik, J.L., Thompson, B.E., Gallegos, M., Petcherski, A.G., Moulder, G., Barstead, R., Wickens, M. and Kimble, J. (2002) A conserved RNA-binding protein controls germline stem cells in *Caenorhabditis elegans*. *Nature*, **417**, 660-663.
- Crittenden, S.L. and Kimble, J. (2008) Analysis of the *C. elegans* germline stem cell region. *Methods Mol Biol*, **450**, 27-44.
- Crittenden, S.L., Troemel, E.R., Evans, T.C. and Kimble, J. (1994) GLP-1 is localized to the mitotic region of the *C. elegans* germ line. *Development*, **120**, 2901-2911.
- Crooks, G.E., Hon, G., Chandonia, J.M. and Brenner, S.E. (2004) WebLogo: a sequence logo generator. *Genome Res*, **14**, 1188-1190.
- Crotty, T., Cai, J., Sakane, F., Taketomi, A., Prescott, S.M. and Topham, M.K. (2006) Diacylglycerol kinase delta regulates protein kinase C and epidermal growth factor receptor signaling. *Proc Natl Acad Sci U S A*, **103**, 15485-15490.
- Cui, Y., Hagan, K.W., Zhang, S. and Peltz, S.W. (1995) Identification and characterization of genes that are required for the accelerated degradation of mRNAs containing a premature translational termination codon. *Genes Dev*, **9**, 423-436.
- Cui, M., Chen, J., Myers, T.R., Hwang, B.J., Sternberg, P.W., Greenwald, I., Han, M. (2006) SynMuv genes redundantly inhibit *lin-3/EGF* expression to prevent inappropriate vulval induction in *C. elegans*. *Dev Cell*, **10**, 667-672.
- David, L., Huber, W., Granovskaia, M., Toedling, J., Palm, C.J., Bofkin, L., Jones, T., Davis, R.W. and Steinmetz, L.M. (2006) A high-resolution map of transcription in the yeast genome. *Proc Natl Acad Sci U S A*, **103**, 5320-5325.
- Doyle, T.G., Wen, C. and Greenwald, I. (2000) SEL-8, a nuclear protein required for LIN-12 and GLP-1 signaling in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*, **97**, 7877-7881.
- Duggan, D.J., Bittner, M., Chen, Y., Meltzer, P. and Trent, J.M. (1999) Expression profiling using cDNA microarrays. *Nat Genet*, **21**, 10-14.
- Eberle, A.B., Stalder, L., Mathys, H., Orozco, R.Z. and Muhlemann, O. (2008) Posttranscriptional gene regulation by spatial rearrangement of the 3' untranslated region. *PLoS Biol*, **6**, e92.

- Eckmann, C.R., Crittenden, S.L., Suh, N. and Kimble, J. (2004) GLD-3 and control of the mitosis/meiosis decision in the germline of *Caenorhabditis elegans*. *Genetics*, **168**, 147-160.
- Eckmann, C.R., Kraemer, B., Wickens, M. and Kimble, J. (2002) GLD-3, a bicaudal-C homolog that inhibits FBF to control germline sex determination in *C. elegans*. *Dev Cell*, **3**, 697-710.
- Eisenmann, D.M., Maloof, J.N., Simske, J.S., Kenyon, C. and Kim, S.K. (1998) The beta-catenin homolog BAR-1 and LET-60 Ras coordinately regulate the Hox gene lin-39 during *Caenorhabditis elegans* vulval development. *Development*, **125**, 3667-3680.
- Ercan, S., Giresi, P.G., Whittle, C.M., Zhang, X., Green, R.D. and Lieb, J.D. (2007) X chromosome repression by localization of the *C. elegans* dosage compensation machinery to sites of transcription initiation. *Nat Genet*, **39**, 403-408.
- Ferguson, E.L. and Horvitz, H.R. (1985) Identification and characterization of 22 genes that affect the vulval cell lineages of the nematode *Caenorhabditis elegans*. *Genetics*, **110**, 17-72.
- Ferguson, E.L. and Horvitz, H.R. (1989) The multivulva phenotype of certain *Caenorhabditis elegans* mutants results from defects in two functionally redundant pathways. *Genetics*, **123**, 109-121.
- Ferguson, E.L., Sternberg, P.W. and Horvitz, H.R. (1987) A genetic pathway for the specification of the vulval cell lineages of *Caenorhabditis elegans*. *Nature*, **326**, 259-267.
- Fiegler, H., Carr, P., Douglas, E.J., Burford, D.C., Hunt, S., Scott, C.E., Smith, J., Vetrie, D., Gorman, P., Tomlinson, I.P. and Carter, N.P. (2003) DNA microarrays for comparative genomic hybridization based on DOP-PCR amplification of BAC and PAC clones. *Genes Chromosomes Cancer*, **36**, 361-374.
- Filipowicz, W. (2005) RNAi: the nuts and bolts of the RISC machine. *Cell*, **122**, 17-20.
- Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E. and Mello, C.C. (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, **391**, 806-811.
- Fitzgerald, K. and Greenwald, I. (1995) Interchangeability of *Caenorhabditis elegans* DSL proteins and intrinsic signalling activity of their extracellular domains in vivo. *Development*, **121**, 4275-4282.
- Francis, R., Maine, E. and Schedl, T. (1995) Analysis of the multiple roles of gld-1 in germline development: interactions with the sex determination cascade and the glp-1 signaling pathway. *Genetics*, **139**, 607-630.

- Fraser, A.G., Kamath, R.S., Zipperlen, P., Martinez-Campos, M., Sohrmann, M. and Ahringer, J. (2000) Functional genomic analysis of *C. elegans* chromosome I by systematic RNA interference. *Nature*, **408**, 325-330.
- Freeman, T.C., Goldovsky, L., Brosch, M., van Dongen, S., Maziere, P., Grocock, R.J., Freilich, S., Thornton, J. and Enright, A.J. (2007) Construction, visualisation, and clustering of transcription networks from microarray expression data. *PLoS Comput Biol*, **3**, 2032-2042.
- Fribourg, S., Gatfield, D., Izaurrealde, E. and Conti, E. (2003) A novel mode of RBD-protein recognition in the Y14-Mago complex. *Nat Struct Biol*, **10**, 433-439.
- Gaiano, N. and Fishell, G. (2002) The role of notch in promoting glial and neural stem cell fates. *Annu Rev Neurosci*, **25**, 471-490.
- Gardner, L.B. (2008) Hypoxic inhibition of nonsense-mediated RNA decay regulates gene expression and the integrated stress response. *Mol Cell Biol*, **28**, 3729-3741.
- Gatfield, D. and Izaurrealde, E. (2004) Nonsense-mediated messenger RNA decay is initiated by endonucleolytic cleavage in *Drosophila*. *Nature*, **429**, 575-578.
- Gatfield, D., Unterholzner, L., Ciccarelli, F.D., Bork, P. and Izaurrealde, E. (2003) Nonsense-mediated mRNA decay in *Drosophila*: at the intersection of the yeast and mammalian pathways. *Embo J*, **22**, 3960-3970.
- Gehen, S. C., Staversky, R. J., Bambara, R. A., Keng, P. C. & O'Reilly, M. A. (2008) hSMG-1 and ATM sequentially and independently regulate the G(1) checkpoint during oxidative stress. *Oncogene*, **27**, 4065-4074.
- Gehring, N.H., Neu-Yilik, G., Schell, T., Hentze, M.W. and Kulozik, A.E. (2003) Y14 and hUpf3b form an NMD-activating complex. *Mol Cell*, **11**, 939-949.
- Greenwald, I.S., Sternberg, P.W. and Horvitz, H.R. (1983) The lin-12 locus specifies cell fates in *Caenorhabditis elegans*. *Cell*, **34**, 435-444.
- Grimson, A., O'Connor, S., Newman, C.L. and Anderson, P. (2004) SMG-1 is a phosphatidylinositol kinase-related protein kinase required for nonsense-mediated mRNA Decay in *Caenorhabditis elegans*. *Mol Cell Biol*, **24**, 7483-7490.
- Guan, Q., Zheng, W., Tang, S., Liu, X., Zinkel, R.A., Tsui, K.W., Yandell, B.S. and Culbertson, M.R. (2006) Impact of nonsense-mediated mRNA decay on the global expression profile of budding yeast. *PLoS Genet*, **2**, e203.
- Han, M., Aroian, R.V. and Sternberg, P.W. (1990) The let-60 locus controls the switch between vulval and nonvulval cell fates in *Caenorhabditis elegans*. *Genetics*, **126**, 899-913.
- Hanada, K., Zhang, X., Borevitz, J.O., Li, W.H. and Shiu, S.H. (2007) A large number of novel coding small open reading frames in the intergenic regions of

- the *Arabidopsis thaliana* genome are transcribed and/or under purifying selection. *Genome Res*, **17**, 632-640.
- Hannon, G.J. (2002) RNA interference. *Nature*, **418**, 244-251.
- Hansen, D., Hubbard, E.J. and Schedl, T. (2004a) Multi-pathway control of the proliferation versus meiotic development decision in the *Caenorhabditis elegans* germline. *Dev Biol*, **268**, 342-357.
- Hansen, D., Wilson-Berry, L., Dang, T. and Schedl, T. (2004b) Control of the proliferation versus meiotic development decision in the *C. elegans* germline through regulation of GLD-1 protein accumulation. *Development*, **131**, 93-104.
- Harrow, J., Denoeud, F., Frankish, A., Reymond, A., Chen, C.K., Chrast, J., Lagarde, J., Gilbert, J.G., Storey, R., Swarbreck, D., Rossier, C., UCLA, C., Hubbard, T., Antonarakis, S.E. and Guigo, R. (2006) GENCODE: producing a reference annotation for ENCODE. *Genome Biol*, **7 Suppl 1**, S4 1-9.
- He, F., Li, X., Spatrik, P., Casillo, R., Dong, S. and Jacobson, A. (2003) Genome-wide analysis of mRNAs regulated by the nonsense-mediated and 5' to 3' mRNA decay pathways in yeast. *Mol Cell*, **12**, 1439-1452.
- Hedgecock, E.M. and Herman, R.K. (1995) The ncl-1 gene and genetic mosaics of *Caenorhabditis elegans*. *Genetics*, **141**, 989-1006.
- Henderson, S.T., Gao, D., Christensen, S. and Kimble, J. (1997) Functional domains of LAG-2, a putative signaling ligand for LIN-12 and GLP-1 receptors in *Caenorhabditis elegans*. *Mol Biol Cell*, **8**, 1751-1762.
- Heo, J.S. and Han, H.J. (2006) PKC and MAPKs pathways mediate EGF-induced stimulation of 2-deoxyglucose uptake in mouse embryonic stem cells. *Cell Physiol Biochem*, **17**, 145-158.
- Herman, R.K. and Hedgecock, E.M. (1990) Limitation of the size of the vulval primordium of *Caenorhabditis elegans* by lin-15 expression in surrounding hypodermis. *Nature*, **348**, 169-171.
- Hodgkin, J., Papp, A., Pulak, R., Ambros, V. and Anderson, P. (1989) A new kind of informational suppression in the nematode *Caenorhabditis elegans*. *Genetics*, **123**, 301-313.
- Horak, C.E. and Snyder, M. (2002) ChIP-chip: a genomic approach for identifying transcription factor binding sites. *Methods Enzymol*, **350**, 469-483.
- Horvitz, H.R. and Sulston, J.E. (1980) Isolation and genetic characterization of cell-lineage mutants of the nematode *Caenorhabditis elegans*. *Genetics*, **96**, 435-454.
- Hsu, V., Zobel, C.L., Lambie, E.J., Schedl, T. and Kornfeld, K. (2002) *Caenorhabditis elegans* lin-45 raf is essential for larval viability, fertility and the induction of vulval cell fates. *Genetics*, **160**, 481-492.

- Huang, L.S., Tzou, P. and Sternberg, P.W. (1994) The lin-15 locus encodes two negative regulators of *Caenorhabditis elegans* vulval development. *Mol Biol Cell*, **5**, 395-411.
- Hubbard, E.J., Dong, Q. and Greenwald, I. (1996) Evidence for physical and functional association between EMB-5 and LIN-12 in *Caenorhabditis elegans*. *Science*, **273**, 112-115.
- Hubbard, E.J. and Greenstein, D. (2005) Introduction to the germ line. *WormBook*, 1-4.
- Hughes, T.R., Mao, M., Jones, A.R., Burchard, J., Marton, M.J., Shannon, K.W., Lefkowitz, S.M., Ziman, M., Schelter, J.M., Meyer, M.R., Kobayashi, S., Davis, C., Dai, H., He, Y.D., Stepaniants, S.B., Cavet, G., Walker, W.L., West, A., Coffey, E., Shoemaker, D.D., Stoughton, R., Blanchard, A.P., Friend, S.H. and Linsley, P.S. (2001) Expression profiling using microarrays fabricated by an ink-jet oligonucleotide synthesizer. *Nat Biotechnol*, **19**, 342-347.
- Hughes, T.R., Marton, M.J., Jones, A.R., Roberts, C.J., Stoughton, R., Armour, C.D., Bennett, H.A., Coffey, E., Dai, H., He, Y.D., Kidd, M.J., King, A.M., Meyer, M.R., Slade, D., Lum, P.Y., Stepaniants, S.B., Shoemaker, D.D., Gachotte, D., Chakraburty, K., Simon, J., Bard, M. and Friend, S.H. (2000) Functional discovery via a compendium of expression profiles. *Cell*, **102**, 109-126.
- Ishida, S., Shigemoto-Mogami, Y., Kagechika, H., Shudo, K., Ozawa, S., Sawada, J., Ohno, Y. and Inoue, K. (2003) Clinically potential subclasses of retinoid synergists revealed by gene expression profiling. *Mol Cancer Ther*, **2**, 49-58.
- Jiang, M., Ryu, J., Kiraly, M., Duke, K., Reinke, V. and Kim, S.K. (2001) Genome-wide analysis of developmental and sex-regulated gene expression profiles in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*, **98**, 218-223.
- Joshua-Tor, L. (2006) The Argonautes. *Cold Spring Harb Symp Quant Biol*, **71**, 67-72.
- Kadyk, L.C. and Kimble, J. (1998) Genetic regulation of entry into meiosis in *Caenorhabditis elegans*. *Development*, **125**, 1803-1813.
- Kamath, R.S., Fraser, A.G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin, A., Le Bot, N., Moreno, S., Sohrmann, M., Welchman, D.P., Zipperlen, P. and Ahringer, J. (2003) Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature*, **421**, 231-237.
- Kamath, R.S., Martinez-Campos, M., Zipperlen, P., Fraser, A.G. and Ahringer, J. (2001) Effectiveness of specific RNA-mediated interference through ingested double-stranded RNA in *Caenorhabditis elegans*. *Genome Biol*, **2**, RESEARCH0002.
- Katz, W.S., Hill, R.J., Clandinin, T.R. and Sternberg, P.W. (1995) Different levels of the *C. elegans* growth factor LIN-3 promote distinct vulval precursor fates. *Cell*, **82**, 297-307.

- Kaygun, H. and Marzluff, W. F. (2005) Regulated degradation of replication-dependent histone mRNAs requires both ATR and Upf1. *Nature Struct. Mol. Biol.*, **12**, 794–800.
- Keene, J.D., Komisarow, J.M. and Friedersdorf, M.B. (2006) RIP-Chip: the isolation and identification of mRNAs, microRNAs and protein components of ribonucleoprotein complexes from cell extracts. *Nat Protoc.*, **1**, 302-307.
- Khajavi, M., Inoue, K. and Lupski, J.R. (2006) Nonsense-mediated mRNA decay modulates clinical outcome of genetic disease. *Eur J Hum Genet*, **14**, 1074-1081.
- Kim, Y. K., Furic, L., Desgroseillers, L. and Maquat, L. E. (2005) Mammalian Staufen1 recruits Upf1 to specific mRNA 3UTRs so as to elicit mRNA decay. *Cell*, **120**, 195–208.
- Kimble, J. (1981) Alterations in cell lineage following laser ablation of cells in the somatic gonad of *Caenorhabditis elegans*. *Dev Biol*, **87**, 286-300.
- Kimble, J. and Hirsh, D. (1979) The postembryonic cell lineages of the hermaphrodite and male gonads in *Caenorhabditis elegans*. *Dev Biol*, **70**, 396-417.
- Kimble, J.E. and White, J.G. (1981) On the control of germ cell development in *Caenorhabditis elegans*. *Dev Biol*, **81**, 208-219.
- Koch, C.M., Andrews, R.M., Flieck, P., Dillon, S.C., Karaoz, U., Clelland, G.K., Wilcox, S., Beare, D.M., Fowler, J.C., Couttet, P., James, K.D., Lefebvre, G.C., Bruce, A.W., Dovey, O.M., Ellis, P.D., Dhami, P., Langford, C.F., Weng, Z., Birney, E., Carter, N.P., Vtrie, D. and Dunham, I. (2007) The landscape of histone modifications across 1% of the human genome in five human cell lines. *Genome Res*, **17**, 691-707.
- Kozak, M. (1984) Point mutations close to the AUG initiator codon affect the efficiency of translation of rat preproinsulin in vivo. *Nature*, **308**, 241-246.
- Kozak, M. (1986) Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell*, **44**, 283-292.
- Kozak, M. (1987) At least six nucleotides preceding the AUG initiator codon enhance translation in mammalian cells. *J Mol Biol*, **196**, 947-950.
- Kraemer, B., Crittenden, S., Gallegos, M., Moulder, G., Barstead, R., Kimble, J. and Wickens, M. (1999) NANOS-3 and FBF proteins physically interact to control the sperm-oocyte switch in *Caenorhabditis elegans*. *Curr Biol*, **9**, 1009-1018.
- Lambie, E.J. and Kimble, J. (1991) Two homologous regulatory genes, lin-12 and glp-1, have overlapping functions. *Development*, **112**, 231-240.
- Lamont, L.B., Crittenden, S.L., Bernstein, D., Wickens, M. and Kimble, J. (2004) FBF-1 and FBF-2 regulate the size of the mitotic region in the *C. elegans* germline. *Dev Cell*, **7**, 697-707.

- Lareau, L.F., Inada, M., Green, R.E., Wengrod, J.C. and Brenner, S.E. (2007) Unproductive splicing of SR genes associated with highly conserved and ultraconserved DNA elements. *Nature*, **446**, 926-929.
- Lee, B.S. and Culbertson, M.R. (1995) Identification of an additional gene required for eukaryotic nonsense mRNA turnover. *Proc Natl Acad Sci U S A*, **92**, 10354-10358.
- Lee, M.H. and Schedl, T. (2004) Translation repression by GLD-1 protects its mRNA targets from nonsense-mediated mRNA decay in *C. elegans*. *Genes Dev*, **18**, 1047-1059.
- Lee, M.Y., Lee, S.H., Kim, Y.H., Heo, J.S., Park, S.H., Lee, J.H. and Han, H.J. (2006a) Effect of EGF on [³H]-thymidine incorporation and cell cycle regulatory proteins in primary cultured chicken hepatocytes: Involvement of Ca²⁺/PKC and MAPKs. *J Cell Biochem*, **99**, 1677-1687.
- Lee, M.Y., Park, S.H., Lee, Y.J., Heo, J.S., Lee, J.H. and Han, H.J. (2006b) EGF-induced inhibition of glucose transport is mediated by PKC and MAPK signal pathways in primary cultured chicken hepatocytes. *Am J Physiol Gastrointest Liver Physiol*, **291**, G744-750.
- Leeds, P., Peltz, S.W., Jacobson, A. and Culbertson, M.R. (1991) The product of the yeast UPF1 gene is required for rapid turnover of mRNAs containing a premature translational termination codon. *Genes Dev*, **5**, 2303-2314.
- Lehner, B., Crombie, C., Tischler, J., Fortunato, A. and Fraser, A.G. (2006) Systematic mapping of genetic interactions in *Caenorhabditis elegans* identifies common modifiers of diverse signaling pathways. *Nat Genet*, **38**, 896-903.
- Lejeune, F., Li, X. and Maquat, L.E. (2003) Nonsense-mediated mRNA decay in mammalian cells involves decapping, deadenylation, and exonucleolytic activities. *Mol Cell*, **12**, 675-687.
- Lelivelt, M.J. and Culbertson, M.R. (1999) Yeast Upf proteins required for RNA surveillance affect global expression of the yeast transcriptome. *Mol Cell Biol*, **19**, 6710-6719.
- Lewis, B.P., Green, R.E. and Brenner, S.E. (2003) Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans. *Proc Natl Acad Sci U S A*, **100**, 189-192.
- Lipshutz, R.J., Fodor, S.P., Gingras, T.R. and Lockhart, D.J. (1999) High density synthetic oligonucleotide arrays. *Nat Genet*, **21**, 20-24.
- Longman, D., Johnstone, I.L. and Caceres, J.F. (2000) Functional characterization of SR and SR-related genes in *Caenorhabditis elegans*. *Embo J*, **19**, 1625-1637.
- Longman, D., Plasterk, R.H., Johnstone, I.L. and Caceres, J.F. (2007) Mechanistic insights and identification of two novel factors in the *C. elegans* NMD pathway. *Genes Dev*, **21**, 1075-1085.

- Losson, R. and Lacroute, F. (1979) Interference of nonsense mutations with eukaryotic messenger RNA stability. *Proc Natl Acad Sci U S A*, **76**, 5134-5137.
- Lykke-Andersen, J., Shu, M.D. and Steitz, J.A. (2001) Communication of the position of exon-exon junctions to the mRNA surveillance machinery by the protein RNPS1. *Science*, **293**, 1836-1839.
- Lyne, R., Burns, G., Mata, J., Penkett, C.J., Rustici, G., Chen, D., Langford, C., Vetrie, D. and Bahler, J. (2003) Whole-genome microarrays of fission yeast: characteristics, accuracy, reproducibility, and processing of array data. *BMC Genomics*, **4**, 27.
- Manak, J.R., Dike, S., Sementchenko, V., Kapranov, P., Biemar, F., Long, J., Cheng, J., Bell, I., Ghosh, S., Piccolboni, A. and Gingeras, T.R. (2006) Biological function of unannotated transcription during the early development of *Drosophila melanogaster*. *Nat Genet*, **38**, 1151-1158.
- Mango, S.E. (2001) Stop making nonSense: the *C. elegans* smg genes. *Trends Genet*, **17**, 646-653.
- Marin, V.A. and Evans, T.C. (2003) Translational repression of a *C. elegans* Notch mRNA by the STAR/KH domain protein GLD-1. *Development*, **130**, 2623-2632.
- Matzke, M.A. and Birchler, J.A. (2005) RNAi-mediated pathways in the nucleus. *Nat Rev Genet*, **6**, 24-35.
- Medghalchi, S.M., Frischmeyer, P.A., Mendell, J.T., Kelly, A.G., Lawler, A.M. and Dietz, H.C. (2001) Rent1, a trans-effector of nonsense-mediated mRNA decay, is essential for mammalian embryonic viability. *Hum Mol Genet*, **10**, 99-105.
- Mellor, H. and Parker, P.J. (1998) The extended protein kinase C superfamily. *Biochem J*, **332 (Pt 2)**, 281-292.
- Mendell, J.T., Sharifi, N.A., Meyers, J.L., Martinez-Murillo, F. and Dietz, H.C. (2004) Nonsense surveillance regulates expression of diverse classes of mammalian transcripts and mutes genomic noise. *Nat Genet*, **36**, 1073-1078.
- Merida, I., Avila-Flores, A. and Merino, E. (2008) Diacylglycerol kinases: at the hub of cell signalling. *Biochem J*, **409**, 1-18.
- Mitchell, P. and Tollervey, D. (2003) An NMD pathway in yeast involving accelerated deadenylation and exosome-mediated 3'-->5' degradation. *Mol Cell*, **11**, 1405-1413.
- Morrison, M., Harris, K.S. and Roth, M.B. (1997) smg mutants affect the expression of alternatively spliced SR protein mRNAs in *Caenorhabditis elegans*. *Proc Natl Acad Sci U S A*, **94**, 9782-9785.

- Nagalakshmi, U., Wang, Z., Waern, K., Shou, C., Raha, D., Gerstein, M. and Snyder, M. (2008) The transcriptional landscape of the yeast genome defined by RNA sequencing. *Science*, **320**, 1344-1349.
- Ni JZ, Grate L, Donohue JP, Preston C, Nobida N, O'Brien G, Shiue L, Clark TA, Blume JE, Ares M, Jr. 2007. Ultraconserved elements are associated with homeostatic control of splicing regulators by alternative splicing and nonsense-mediated decay. *Genes Dev*, **21**, 708-718.
- Nishizuka, Y. (1984) The role of protein kinase C in cell surface signal transduction and tumour promotion. *Nature*, **308**, 693-698.
- Ohmachi, M., Rocheleau, C.E., Church, D., Lambie, E., Schedl, T. and Sundaram, M.V. (2002) *C. elegans* ksr-1 and ksr-2 have both unique and redundant functions and are required for MPK-1 ERK phosphorylation. *Curr Biol*, **12**, 427-433.
- Ohnishi, T., Yamashita, A., Kashima, I., Schell, T., Anders, K.R., Grimson, A., Hachiya, T., Hentze, M.W., Anderson, P. and Ohno, S. (2003) Phosphorylation of hUPF1 induces formation of mRNA surveillance complexes containing hSMG-5 and hSMG-7. *Mol Cell*, **12**, 1187-1200.
- Olivas, W. and Parker, R. (2000) The Puf3 protein is a transcript-specific regulator of mRNA degradation in yeast. *Embo J*, **19**, 6602-6611.
- Oliveira, V., Romanow, W.J., Geisen, C., Otterness, D.M., Mercurio, F., Wang, H.G., Dalton, W.S. and Abraham, R.T. (2008) A protective role for the human SMG-1 kinase against tumor necrosis factor-alpha-induced apoptosis. *J Biol Chem*, **283**, 13174-84.
- Page, M.F., Carr, B., Anders, K.R., Grimson, A. and Anderson, P. (1999) SMG-2 is a phosphorylated protein required for mRNA surveillance in *Caenorhabditis elegans* and related to Upf1p of yeast. *Mol Cell Biol*, **19**, 5943-5951.
- Pepper, A.S., Lo, T.W., Killian, D.J., Hall, D.H. and Hubbard, E.J. (2003) The establishment of *Caenorhabditis elegans* germline pattern is controlled by overlapping proximal and distal somatic gonad signals. *Dev Biol*, **259**, 336-350.
- Petcherski, A.G. and Kimble, J. (2000) LAG-3 is a putative transcriptional activator in the *C. elegans* Notch pathway. *Nature*, **405**, 364-368.
- Petricoin, E.F., 3rd, Hackett, J.L., Lesko, L.J., Puri, R.K., Gutman, S.I., Chumakov, K., Woodcock, J., Feigal, D.W., Jr., Zoon, K.C. and Sistare, F.D. (2002) Medical applications of microarray technologies: a regulatory science perspective. *Nat Genet*, **32 Suppl**, 474-479.
- Poulin, G., Dong, Y., Fraser, A.G., Hopper, N.A. and Ahringer, J. (2005) Chromatin regulation and sumoylation in the inhibition of Ras-induced vulval development in *Caenorhabditis elegans*. *Embo J*, **24**, 2613-2623.

- Qiao, L., Lissemore, J.L., Shu, P., Smardon, A., Gelber, M.B. and Maine, E.M. (1995) Enhancers of *glp-1*, a gene required for cell-signaling in *Caenorhabditis elegans*, define a set of genes required for germline development. *Genetics*, **141**, 551-569.
- Redon, R., Ishikawa, S., Fitch, K.R., Feuk, L., Perry, G.H., Andrews, T.D., Fiegler, H., Shapero, M.H., Carson, A.R., Chen, W., Cho, E.K., Dallaire, S., Freeman, J.L., Gonzalez, J.R., Gratacos, M., Huang, J., Kalaitzopoulos, D., Komura, D., MacDonald, J.R., Marshall, C.R., Mei, R., Montgomery, L., Nishimura, K., Okamura, K., Shen, F., Somerville, M.J., Tchinda, J., Valsesia, A., Woodwark, C., Yang, F., Zhang, J., Zerjal, T., Zhang, J., Armengol, L., Conrad, D.F., Estivill, X., Tyler-Smith, C., Carter, N.P., Aburatani, H., Lee, C., Jones, K.W., Scherer, S.W. and Hurles, M.E. (2006) Global variation in copy number in the human genome. *Nature*, **444**, 444-454.
- Rehwinkel, J., Letunic, I., Raes, J., Bork, P. and Izaurralde, E. (2005) Nonsense-mediated mRNA decay factors act in concert to regulate common mRNA targets. *Rna*, **11**, 1530-1544.
- Reinke, V. (2002) Functional exploration of the *C. elegans* genome using DNA microarrays. *Nat Genet*, **32 Suppl**, 541-546.
- Reinke, V., Gil, I.S., Ward, S. and Kazmer, K. (2004) Genome-wide germline-enriched and sex-biased expression profiles in *Caenorhabditis elegans*. *Development*, **131**, 311-323.
- Reinke, V., Smith, H.E., Nance, J., Wang, J., Van Doren, C., Begley, R., Jones, S.J., Davis, E.B., Scherer, S., Ward, S. and Kim, S.K. (2000) A global profile of germline gene expression in *C. elegans*. *Mol Cell*, **6**, 605-616.
- Reinke, V. and White, K.P. (2002) Developmental genomic approaches in model organisms. *Annu Rev Genomics Hum Genet*, **3**, 153-178.
- Rodriguez-Gabriel, M.A., Watt, S., Bahler, J. and Russell, P. (2006) Upf1, an RNA helicase required for nonsense-mediated mRNA decay, modulates the transcriptional response to oxidative stress in fission yeast. *Mol Cell Biol*, **26**, 6347-6356.
- Rual, J.F., Ceron, J., Koreth, J., Hao, T., Nicot, A.S., Hirozane-Kishikawa, T., Vandenhoute, J., Orkin, S.H., Hill, D.E., van den Heuvel, S. and Vidal, M. (2004) Toward improving *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library. *Genome Res*, **14**, 2162-2168.
- Ruiz-Echevarria, M.J. and Peltz, S.W. (2000) The RNA binding protein Pub1 modulates the stability of transcripts containing upstream open reading frames. *Cell*, **101**, 741-751.
- Ryder, S.P., Frater, L.A., Abramovitz, D.L., Goodwin, E.B. and Williamson, J.R. (2004) RNA target specificity of the STAR/GSG domain post-transcriptional regulatory protein GLD-1. *Nat Struct Mol Biol*, **11**, 20-28.

- Saltzman AL, Kim YK, Pan Q, Fagnani MM, Maquat LE, Blencowe BJ. 2008. Regulation of multiple core spliceosomal proteins by alternative splicing-coupled nonsense-mediated mRNA decay. *Mol Cell Biol*, **28**, 4320-4330.
- Schena, M., Shalon, D., Davis, R.W. and Brown, P.O. (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science*, **270**, 467-470.
- Schneider, T.D. and Stephens, R.M. (1990) Sequence logos: a new way to display consensus sequences. *Nucleic Acids Res*, **18**, 6097-6100.
- Schroeter, E.H., Kisslinger, J.A. and Kopan, R. (1998) Notch-1 signalling requires ligand-induced proteolytic release of intracellular domain. *Nature*, **393**, 382-386.
- Shaye, D.D. and Greenwald, I. (2002) Endocytosis-mediated downregulation of LIN-12/Notch upon Ras activation in *Caenorhabditis elegans*. *Nature*, **420**, 686-690.
- Simske, J.S. and Kim, S.K. (1995) Sequential signalling during *Caenorhabditis elegans* vulval induction. *Nature*, **375**, 142-146.
- Singh, G., Rebbapragada, I. and Lykke-Andersen, J. (2008) A competition between stimulators and antagonists of Upf complex recruitment governs human nonsense-mediated mRNA decay. *PLoS Biol*, **6**, e111.
- Sriraman, V., Modi, S.R., Bodenburg, Y., Denner, L.A. and Urban, R.J. (2008) Identification of ERK and JNK as signaling mediators on protein kinase C activation in cultured granulosa cells. *Mol Cell Endocrinol*.
- Sternberg, P.W. (1988) Lateral inhibition during vulval induction in *Caenorhabditis elegans*. *Nature*, **335**, 551-554.
- Sternberg, P.W. and Horvitz, H.R. (1986) Pattern formation during vulval development in *C. elegans*. *Cell*, **44**, 761-772.
- Stiernagle, T. (2006) Maintenance of *C. elegans*. *Wormbook*, **11**, 1-11.
- Sulston, J.E. and Horvitz, H.R. (1977) Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Dev Biol*, **56**, 110-156.
- Sulston, J.E., Schierenberg, E., White, J.G. and Thomson, J.N. (1983) The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev Biol*, **100**, 64-119.
- Sulston, J.E. and White, J.G. (1980) Regulation and cell autonomy during postembryonic development of *Caenorhabditis elegans*. *Dev Biol*, **78**, 577-597.
- Sundaram, M. and Greenwald, I. (1993) Suppressors of a lin-12 hypomorph define genes that interact with both lin-12 and glp-1 in *Caenorhabditis elegans*. *Genetics*, **135**, 765-783.

- Tabara, H., Grishok, A. and Mello, C.C. (1998) RNAi in *C. elegans*: soaking in the genome sequence. *Science*, **282**, 430-431.
- Tan, P.B., Lackner, M.R. and Kim, S.K. (1998) MAP kinase signaling specificity mediated by the LIN-1 Ets/LIN-31 WH transcription factor complex during *C. elegans* vulval induction. *Cell*, **93**, 569-580.
- Tarraga, J., Medina, I., Carbonell, J., Huerta-Cepas, J., Minguez, P., Alloza, E., Al-Shahrour, F., Vegas-Azcarate, S., Goetz, S., Escobar, P., Garcia-Garcia, F., Conesa, A., Montaner, D. and Dopazo, J. (2008) GEPAS, a web-based tool for microarray data analysis and interpretation. *Nucleic Acids Res*, **36**, W308-314.
- The FANTOM Consortium and RIKEN Genome Exploration Research Group and Genome Science Group (Genome Network Project Core Group). (2006) The Transcriptional Landscape of the Mammalian Genome. *Science*, **309**, 1559-1563.
- Timmons, L., Court, D.L. and Fire, A. (2001) Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene*, **263**, 103-112.
- Timmons, L. and Fire, A. (1998) Specific interference by ingested dsRNA. *Nature*, **395**, 854.
- Wang, L., Eckmann, C.R., Kadyk, L.C., Wickens, M. and Kimble, J. (2002) A regulatory cytoplasmic poly(A) polymerase in *Caenorhabditis elegans*. *Nature*, **419**, 312-316.
- Watt, F.M. and Hogan, B.L. (2000) Out of Eden: stem cells and their niches. *Science*, **287**, 1427-1430.
- Wilhelm, B.T., Marguerat, S., Watt, S., Schubert, F., Wood, V., Goodhead, I., Penkett, C.J., Rogers, J. and Bahler, J. (2008) Dynamic repertoire of a eukaryotic transcriptome surveyed at single-nucleotide resolution. *Nature*, **453**, 1239-1243.
- Winston, W.M., Molodowitch, C. and Hunter, C.P. (2002) Systemic RNAi in *C. elegans* requires the putative transmembrane protein SID-1. *Science*, **295**, 2456-2459.
- Winston, W.M., Sutherlin, M., Wright, A.J., Feinberg, E.H. and Hunter, C.P. (2007) *Caenorhabditis elegans* SID-2 is required for environmental RNA interference. *Proc Natl Acad Sci U S A*, **104**, 10565-10570.
- Wreden, C., Verrotti, A.C., Schisa, J.A., Lieberfarb, M.E. and Strickland, S. (1997) Nanos and pumilio establish embryonic polarity in *Drosophila* by promoting posterior deadenylation of hunchback mRNA. *Development*, **124**, 3015-3023.
- Wu, Y. and Han, M. (1994) Suppression of activated Let-60 ras protein defines a role of *Caenorhabditis elegans* Sur-1 MAP kinase in vulval differentiation. *Genes Dev*, **8**, 147-159.

- Wultsch, T., Chourbaji, S., Fritzen, S., Kittelt, S., Grunblatt, E., Gerlach, M., Gutknecht, L., Chizat, F., Golfier, G., Schmitt, A., Gass, P., Lesch, K.P. and Reif, A. (2007) Behavioural and expressional phenotyping of nitric oxide synthase-I knockdown animals. *J Neural Transm Suppl*, 69-85.
- Yamashita, A., Kashima, I. and Ohno, S. (2005) The role of SMG-1 in nonsense-mediated mRNA decay. *Biochim Biophys Acta*, **1754**, 305-315.
- Yang, Y.H., Dudoit, S., Luu, P., Lin, D.M., Peng, V., Ngai, J. and Speed, T.P. (2002) Normalization for cDNA microarray data: a robust composite method addressing single and multiple slide systematic variation. *Nucleic Acids Res*, **30**, e15.
- Yochem, J. and Greenwald, I. (1989) glp-1 and lin-12, genes implicated in distinct cell-cell interactions in *C. elegans*, encode similar transmembrane proteins. *Cell*, **58**, 553-563.
- Yochem, J., Weston, K. and Greenwald, I. (1988) The *Caenorhabditis elegans* lin-12 gene encodes a transmembrane protein with overall similarity to *Drosophila* Notch. *Nature*, **335**, 547-550.
- Yoo, A.S., Bais, C. and Greenwald, I. (2004) Crosstalk between the EGFR and LIN-12/Notch pathways in *C. elegans* vulval development. *Science*, **303**, 663-666.
- Zamore, P.D. and Haley, B. (2005) Ribo-gnome: the big world of small RNAs. *Science*, **309**, 1519-1524.
- Zhang, B., Gallegos, M., Puoti, A., Durkin, E., Fields, S., Kimble, J. and Wickens, M.P. (1997) A conserved RNA-binding protein that regulates sexual fates in the *C. elegans* hermaphrodite germ line. *Nature*, **390**, 477-484.
- Zien, A., Gebhard, P.M., Fundel, K. and Aigner, T. (2007) Phenotyping of chondrocytes in vivo and in vitro using cDNA array technology. *Clin Orthop Relat Res*, **460**, 226-233.