

## REFERENCES

- Abe, K., Hashiyama, M., Macgregor, G., and Yamamura, K. (1996). Purification of primordial germ cells from TNAPbeta-geo mouse embryos using FACS-gal. *Dev Biol* *180*, 468-472.
- Adolfsson, J., Mansson, R., Buza-Vidas, N., Hultquist, A., Liuba, K., Jensen, C.T., Bryder, D., Yang, L., Borge, O.J., Thoren, L.A., *et al.* (2005). Identification of Flt3+ lympho-myeloid stem cells lacking erythro-megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell* *121*, 295-306.
- Akashi, K., Kondo, M., Cheshier, S., Shizuru, J., Gandy, K., Domen, J., Mebius, R., Traver, D., and Weissman, I.L. (1999). Lymphoid development from stem cells and the common lymphocyte progenitors. *Cold Spring Harb Symp Quant Biol* *64*, 1-12.
- Akashi, K., Kondo, M., and Weissman, I.L. (1998). Role of interleukin-7 in T-cell development from hematopoietic stem cells. *Immunol Rev* *165*, 13-28.
- Albu, D.I., Feng, D., Bhattacharya, D., Jenkins, N.A., Copeland, N.G., Liu, P., and Avram, D. (2007). BCL11B is required for positive selection and survival of double-positive thymocytes. *J Exp Med* *204*, 3003-3015.
- Alici, E., Sutlu, T., Bjorkstrand, B., Gilljam, M., Stellan, B., Nahi, H., Quezada, H.C., Gahrton, G., Ljunggren, H.G., and Dilber, M.S. (2008). Autologous antitumor activity by NK cells expanded from myeloma patients using GMP-compliant components. *Blood* *111*, 3155-3162.
- Allman, D., Sambandam, A., Kim, S., Miller, J.P., Pagan, A., Well, D., Meraz, A., and Bhandoola, A. (2003). Thymopoiesis independent of common lymphoid progenitors. *Nat Immunol* *4*, 168-174.

Amsen, D., Antov, A., and Flavell, R.A. (2009). The different faces of Notch in T-helper-cell differentiation. *Nat Rev Immunol* 9, 116-124.

Anderson, G., Jenkinson, W.E., Jones, T., Parnell, S.M., Kinsella, F.A., White, A.J., Pongracz, J.E., Rossi, S.W., and Jenkinson, E.J. (2006). Establishment and functioning of intrathymic microenvironments. *Immunol Rev* 209, 10-27.

Anderson, G., Moore, N.C., Owen, J.J., and Jenkinson, E.J. (1996). Cellular interactions in thymocyte development. *Annu Rev Immunol* 14, 73-99.

Antoch, M.P., Song, E.J., Chang, A.M., Vitaterna, M.H., Zhao, Y., Wilsbacher, L.D., Sangoram, A.M., King, D.P., Pinto, L.H., and Takahashi, J.S. (1997). Functional identification of the mouse circadian Clock gene by transgenic BAC rescue. *Cell* 89, 655-667.

Arlotta, P., Molyneaux, B.J., Chen, J., Inoue, J., Kominami, R., and Macklis, J.D. (2005). Neuronal subtype-specific genes that control corticospinal motor neuron development in vivo. *Neuron* 45, 207-221.

Arlotta, P., Molyneaux, B.J., Jabaudon, D., Yoshida, Y., and Macklis, J.D. (2008). Ctip2 controls the differentiation of medium spiny neurons and the establishment of the cellular architecture of the striatum. *J Neurosci* 28, 622-632.

Assa-Kunik, E., Torres, I.L., Schejter, E.D., Johnston, D.S., and Shilo, B.Z. (2007). *Drosophila* follicle cells are patterned by multiple levels of Notch signaling and antagonism between the Notch and JAK/STAT pathways. *Development* 134, 1161-1169.

Assarsson, E., Kambayashi, T., Sandberg, J.K., Hong, S., Taniguchi, M., Van Kaer, L., Ljunggren, H.G., and Chambers, B.J. (2000). CD8<sup>+</sup> T cells rapidly acquire NK1.1 and NK cell-associated molecules upon stimulation in vitro and in vivo. *J Immunol* 165, 3673-3679.

Avram, D., Fields, A., Pretty On Top, K., Nevriy, D.J., Ishmael, J.E., and Leid, M. (2000). Isolation of a novel family of C(2)H(2) zinc finger proteins implicated in transcriptional repression mediated by chicken ovalbumin upstream promoter transcription factor (COUP-TF) orphan nuclear receptors. *J Biol Chem* 275, 10315-10322.

Bain, G., Maandag, E.C., Izon, D.J., Amsen, D., Kruisbeek, A.M., Weintraub, B.C., Krop, I., Schlissel, M.S., Feeney, A.J., van Roon, M., *et al.* (1994). E2A proteins are required for proper B cell development and initiation of immunoglobulin gene rearrangements. *Cell* 79, 885-892.

Bajoghli, B., Aghaallaei, N., Hess, I., Rode, I., Netuschil, N., Tay, B.H., Venkatesh, B., Yu, J.K., Kaltenbach, S.L., Holland, N.D., *et al.* (2009). Evolution of genetic networks underlying the emergence of thymopoiesis in vertebrates. *Cell* 138, 186-197.

Barrett, A.J., and Le Blanc, K. (2010). Immunotherapy prospects for acute myeloid leukaemia. *Clin Exp Immunol* 161, 223-232.

Barton, K., Muthusamy, N., Fischer, C., Ting, C.N., Walunas, T.L., Lanier, L.L., and Leiden, J.M. (1998). The Ets-1 transcription factor is required for the development of natural killer cells in mice. *Immunity* 9, 555-563.

Bell, J.J., and Bhandoola, A. (2008). The earliest thymic progenitors for T cells possess myeloid lineage potential. *Nature* 452, 764-767.

Benczik, M., and Gaffen, S.L. (2004). The interleukin (IL)-2 family cytokines: survival and proliferation signaling pathways in T lymphocytes. *Immunol Invest* 33, 109-142.

Bendelac, A. (1995). Positive selection of mouse NK1+ T cells by CD1-expressing cortical thymocytes. *J Exp Med* 182, 2091-2096.

Benlagha, K., Kyin, T., Beavis, A., Teyton, L., and Bendelac, A. (2002). A thymic precursor to the NK T cell lineage. *Science* 296, 553-555.

Benlagha, K., Wei, D.G., Veiga, J., Teyton, L., and Bendelac, A. (2005). Characterization of the early stages of thymic NKT cell development. *J Exp Med* 202, 485-492.

Bernard, O.A., Busson-LeConiat, M., Ballerini, P., Mauchauffe, M., Della Valle, V., Monni, R., Nguyen Khac, F., Mercher, T., Penard-Lacronique, V., Pasturaud, P., *et al.* (2001). A new recurrent and specific cryptic translocation, t(5;14)(q35;q32), is associated with expression of the Hox11L2 gene in T acute lymphoblastic leukemia. *Leukemia* 15, 1495-1504.

Betz, U.A., Vosshenrich, C.A., Rajewsky, K., and Muller, W. (1996). Bypass of lethality with mosaic mice generated by Cre-loxP-mediated recombination. *Curr Biol* 6, 1307-1316.

Blott, E.J., and Griffiths, G.M. (2002). Secretory lysosomes. *Nat Rev Mol Cell Biol* 3, 122-131.

Boos, M.D., Yokota, Y., Eberl, G., and Kee, B.L. (2007). Mature natural killer cell and lymphoid tissue-inducing cell development requires Id2-mediated suppression of E protein activity. *J Exp Med* 204, 1119-1130.

Borrego, F., Masilamani, M., Marusina, A.I., Tang, X., and Coligan, J.E. (2006). The CD94/NKG2 family of receptors: from molecules and cells to clinical relevance. *Immunol Res* 35, 263-278.

Bradley, A., Evans, M., Kaufman, M.H., and Robertson, E. (1984). Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature* 309, 255-256.

Brinster, R.L., Chen, H.Y., Trumbauer, M., Senechal, A.W., Warren, R., and Palmiter, R.D. (1981). Somatic expression of herpes thymidine kinase in mice following injection of a fusion gene into eggs. *Cell* 27, 223-231.

Brocard, J., Warot, X., Wendling, O., Messaddeq, N., Vonesch, J.L., Chambon, P., and Metzger, D. (1997). Spatio-temporally controlled site-specific somatic mutagenesis in the mouse. *Proc Natl Acad Sci U S A* 94, 14559-14563.

Bunz, F., Dutriaux, A., Lengauer, C., Waldman, T., Zhou, S., Brown, J.P., Sedivy, J.M., Kinzler, K.W., and Vogelstein, B. (1998). Requirement for p53 and p21 to sustain G2 arrest after DNA damage. *Science* 282, 1497-1501.

Burbach, B.J., Medeiros, R.B., Mueller, K.L., and Shimizu, Y. (2007). T-cell receptor signaling to integrins. *Immunol Rev* 218, 65-81.

Busslinger, M. (2004). Transcriptional control of early B cell development. *Annu Rev Immunol* 22, 55-79.

Capecchi, M.R. (1989). Altering the genome by homologous recombination. *Science* 244, 1288-1292.

Caraux, A., Kim, N., Bell, S.E., Zompi, S., Ranson, T., Lesjean-Pottier, S., Garcia-Ojeda, M.E., Turner, M., and Colucci, F. (2006). Phospholipase C-gamma2 is essential for NK cell cytotoxicity and innate immunity to malignant and virally infected cells. *Blood* 107, 994-1002.

Carlens, S., Gilljam, M., Chambers, B.J., Aschan, J., Guven, H., Ljunggren, H.G., Christensson, B., and Dilber, M.S. (2001). A new method for in vitro expansion of cytotoxic human CD3-CD56+ natural killer cells. *Hum Immunol* 62, 1092-1098.

Carotta, S., Brady, J., Wu, L., and Nutt, S.L. (2006). Transient Notch signaling induces NK cell potential in Pax5-deficient pro-B cells. *Eur J Immunol* 36, 3294-3304.

Carpenter, A.C., and Bosselut, R. (2010). Decision checkpoints in the thymus. *Nat Immunol* 11, 666-673.

Carrio, R., Rolle, C.E., and Malek, T.R. (2007). Non-redundant role for IL-7R signaling for the survival of CD8<sup>+</sup> memory T cells. *Eur J Immunol* 37, 3078-3088.

Casey, T.P. (1966). Autoimmunity in strains of New Zealand mice. *NZ Med J* 65, 105-110.

Chambers, S.M., Boles, N.C., Lin, K.Y., Tierney, M.P., Bowman, T.V., Bradfute, S.B., Chen, A.J., Merchant, A.A., Sirin, O., Weksberg, D.C., *et al.* (2007). Hematopoietic fingerprints: an expression database of stem cells and their progeny. *Cell Stem Cell* 1, 578-591.

Chan, W., Costantino, N., Li, R., Lee, S.C., Su, Q., Melvin, D., Court, D.L., and Liu, P. (2007). A recombineering based approach for high-throughput conditional knockout targeting vector construction. *Nucleic Acids Res* 35, e64.

Chang, C., Rodriguez, A., Carretero, M., Lopez-Botet, M., Phillips, J.H., and Lanier, L.L. (1995). Molecular characterization of human CD94: a type II membrane glycoprotein related to the C-type lectin superfamily. *Eur J Immunol* 25, 2433-2437.

Chang, C.C., and Ferrone, S. (2007). Immune selective pressure and HLA class I antigen defects in malignant lesions. *Cancer Immunol Immunother* 56, 227-236.

Cherrier, T., Suzanne, S., Redel, L., Calao, M., Marban, C., Samah, B., Mukerjee, R., Schwartz, C., Gras, G., Sawaya, B.E., *et al.* (2009). p21(WAF1) gene promoter is epigenetically silenced by CTIP2 and SUV39H1. *Oncogene* 28, 3380-3389.

Ciofani, M., and Zuniga-Pflucker, J.C. (2007). The thymus as an inductive site for T lymphopoiesis. *Annu Rev Cell Dev Biol* 23, 463-493.

Cismasiu, V.B., Adamo, K., Gecewicz, J., Duque, J., Lin, Q., and Avram, D. (2005). BCL11B functionally associates with the NuRD complex in T lymphocytes to repress targeted promoter. *Oncogene* 24, 6753-6764.

Cismasiu, V.B., Duque, J., Paskaleva, E., Califano, D., Ghanta, S., Young, H.A., and Avram, D. (2009). BCL11B enhances TCR/CD28-triggered NF-kappaB activation through up-regulation of Cot kinase gene expression in T-lymphocytes. *Biochem J* 417, 457-466.

Cismasiu, V.B., Ghanta, S., Duque, J., Albu, D.I., Chen, H.M., Kasturi, R., and Avram, D. (2006). BCL11B participates in the activation of IL2 gene expression in CD4+ T lymphocytes. *Blood* 108, 2695-2702.

Cobaleda, C., Jochum, W., and Busslinger, M. (2007). Conversion of mature B cells into T cells by dedifferentiation to uncommitted progenitors. *Nature* 449, 473-477.

Colucci, F., Caligiuri, M.A., and Di Santo, J.P. (2003). What does it take to make a natural killer? *Nat Rev Immunol* 3, 413-425.

Colucci, F., Samson, S.I., DeKoter, R.P., Lantz, O., Singh, H., and Di Santo, J.P. (2001). Differential requirement for the transcription factor PU.1 in the generation of natural killer cells versus B and T cells. *Blood* 97, 2625-2632.

Colucci, F., Soudais, C., Rosmaraki, E., Vanes, L., Tybulewicz, V.L., and Di Santo, J.P. (1999). Dissecting NK cell development using a novel alymphoid mouse model: investigating the role of the c-abl proto-oncogene in murine NK cell differentiation. *J Immunol* 162, 2761-2765.

Cook, H.L., Lytle, J.R., Mischo, H.E., Li, M.J., Rossi, J.J., Silva, D.P., Desrosiers, R.C., and Steitz, J.A. (2005). Small nuclear RNAs encoded by Herpesvirus saimiri upregulate the expression of genes linked to T cell activation in virally transformed T cells. *Curr Biol* 15, 974-979.

Cronin, S.J., and Penninger, J.M. (2007). From T-cell activation signals to signaling control of anti-cancer immunity. *Immunol Rev* 220, 151-168.

Cunningham, J.J., and Roussel, M.F. (2001). Cyclin-dependent kinase inhibitors in the development of the central nervous system. *Cell Growth Differ* 12, 387-396.

Dave, V.P., Allman, D., Keefe, R., Hardy, R.R., and Kappes, D.J. (1998). HD mice: a novel mouse mutant with a specific defect in the generation of CD4(+) T cells. *Proc Natl Acad Sci U S A* 95, 8187-8192.

David-Fung, E.S., Butler, R., Buzi, G., Yui, M.A., Diamond, R.A., Anderson, M.K., Rowen, L., and Rothenberg, E.V. (2009). Transcription factor expression dynamics of early T-lymphocyte specification and commitment. *Dev Biol* 325, 444-467.

de Alboran, I.M., O'Hagan, R.C., Gartner, F., Malynn, B., Davidson, L., Rickert, R., Rajewsky, K., DePinho, R.A., and Alt, F.W. (2001). Analysis of C-MYC function in normal cells via conditional gene-targeted mutation. *Immunity* 14, 45-55.

Degli-Esposti, M.A., and Smyth, M.J. (2005). Close encounters of different kinds: dendritic cells and NK cells take centre stage. *Nat Rev Immunol* 5, 112-124.

Desplats, P.A., Lambert, J.R., and Thomas, E.A. (2008). Functional roles for the striatal-enriched transcription factor, Bcl11b, in the control of striatal gene expression and transcriptional dysregulation in Huntington's disease. *Neurobiol Dis* 31, 298-308.

Di Santo, J.P. (2006). Natural killer cell developmental pathways: a question of balance. *Annu Rev Immunol* 24, 257-286.

Di Santo, J.P. (2008). Natural killer cells: diversity in search of a niche. *Nat Immunol* 9, 473-475.

Di Santo, J.P., and Vosshenrich, C.A. (2006). Bone marrow versus thymic pathways of natural killer cell development. *Immunol Rev* 214, 35-46.



Ding, S., Wu, X., Li, G., Han, M., Zhuang, Y., and Xu, T. (2005). Efficient transposition of the piggyBac (PB) transposon in mammalian cells and mice. *Cell* *122*, 473-483.

Dudley, E.C., Petrie, H.T., Shah, L.M., Owen, M.J., and Hayday, A.C. (1994). T cell receptor beta chain gene rearrangement and selection during thymocyte development in adult mice. *Immunity* *1*, 83-93.

Dunn, G.P., Old, L.J., and Schreiber, R.D. (2004). The three Es of cancer immunoediting. *Annu Rev Immunol* *22*, 329-360.

Evans, M.J., and Kaufman, M.H. (1981). Establishment in culture of pluripotential cells from mouse embryos. *Nature* *292*, 154-156.

Feltri, M.L., D'Antonio, M., Quattrini, A., Numerato, R., Arona, M., Previtali, S., Chiu, S.Y., Messing, A., and Wrabetz, L. (1999). A novel P0 glycoprotein transgene activates expression of lacZ in myelin-forming Schwann cells. *Eur J Neurosci* *11*, 1577-1586.

Ferlazzo, G., and Munz, C. (2009). Dendritic cell interactions with NK cells from different tissues. *J Clin Immunol* *29*, 265-273.

Feyerabend, T.B., Terszowski, G., Tietz, A., Blum, C., Luche, H., Gossler, A., Gale, N.W., Radtke, F., Fehling, H.J., and Rodewald, H.R. (2009). Deletion of Notch1 converts pro-T cells to dendritic cells and promotes thymic B cells by cell-extrinsic and cell-intrinsic mechanisms. *Immunity* *30*, 67-79.

Forsberg, E.C., Downs, K.M., and Bresnick, E.H. (2000). Direct interaction of NF-E2 with hypersensitive site 2 of the beta-globin locus control region in living cells. *Blood* *96*, 334-339.

Fowlkes, B.J., and Robey, E.A. (2002). A reassessment of the effect of activated Notch1 on CD4 and CD8 T cell development. *J Immunol* *169*, 1817-1821.

Fujimoto, S., Ikawa, T., Kina, T., and Yokota, Y. (2007). Forced expression of Id2 in fetal thymic T cell progenitors allows some of their progeny to adopt NK cell fate. *Int Immunol* *19*, 1175-1182.

Fung-Leung, W.P., Schilham, M.W., Rahemtulla, A., Kundig, T.M., Vollenweider, M., Potter, J., van Ewijk, W., and Mak, T.W. (1991). CD8 is needed for development of cytotoxic T cells but not helper T cells. *Cell* *65*, 443-449.

Gadue, P., and Stein, P.L. (2002). NK T cell precursors exhibit differential cytokine regulation and require Itk for efficient maturation. *J Immunol* *169*, 2397-2406.

Ganguli-Indra, G., Liang, X., Hyter, S., Leid, M., Hanifin, J., and Indra, A.K. (2009). Expression of COUP-TF-interacting protein 2 (CTIP2) in human atopic dermatitis and allergic contact dermatitis skin. *Exp Dermatol* *18*, 994-996.

Gascoyne, D.M., Long, E., Veiga-Fernandes, H., de Boer, J., Williams, O., Seddon, B., Coles, M., Kioussis, D., and Brady, H.J. (2009). The basic leucine zipper transcription factor E4BP4 is essential for natural killer cell development. *Nat Immunol* *10*, 1118-1124.

Glaser, S., Anastassiadis, K., and Stewart, A.F. (2005). Current issues in mouse genome engineering. *Nat Genet* *37*, 1187-1193.

Glick, B. (1979). The Avian Immune System. *Avian Diseases* *23*, 282-289.

Go, R., Hirose, S., Morita, S., Yamamoto, T., Katsuragi, Y., Mishima, Y., and Kominami, R. (2010). Bcl11b heterozygosity promotes clonal expansion and differentiation arrest of thymocytes in gamma-irradiated mice. *Cancer Sci* *101*, 1347-1353.

Godfrey, D.I., Kennedy, J., Suda, T., and Zlotnik, A. (1993). A developmental pathway involving four phenotypically and functionally distinct subsets of CD3-CD4-CD8- triple-negative adult mouse thymocytes defined by CD44 and CD25 expression. *J Immunol* *150*, 4244-4252.

Godfrey, D.I., MacDonald, H.R., Kronenberg, M., Smyth, M.J., and Van Kaer, L. (2004). NKT cells: what's in a name? *Nat Rev Immunol* 4, 231-237.

Godin, I., and Cumano, A. (2002). The hare and the tortoise: an embryonic haematopoietic race. *Nat Rev Immunol* 2, 593-604.

Godin, I., Dieterlen-Lievre, F., and Cumano, A. (1995). Emergence of multipotent hemopoietic cells in the yolk sac and paraaortic splanchnopleura in mouse embryos, beginning at 8.5 days postcoitus. *Proc Natl Acad Sci U S A* 92, 773-777.

Golonzhka, O., Leid, M., Indra, G., and Indra, A.K. (2007). Expression of COUP-TF-interacting protein 2 (CTIP2) in mouse skin during development and in adulthood. *Gene Expr Patterns* 7, 754-760.

Golonzhka, O., Liang, X., Messaddeq, N., Bornert, J.M., Campbell, A.L., Metzger, D., Chambon, P., Ganguli-Indra, G., Leid, M., and Indra, A.K. (2009a). Dual role of COUP-TF-interacting protein 2 in epidermal homeostasis and permeability barrier formation. *J Invest Dermatol* 129, 1459-1470.

Golonzhka, O., Metzger, D., Bornert, J.M., Bay, B.K., Gross, M.K., Kioussi, C., and Leid, M. (2009b). *Ctip2/Bcl11b* controls ameloblast formation during mammalian odontogenesis. *Proc Natl Acad Sci U S A* 106, 4278-4283.

Gorelik, E., Wiltrot, R.H., Okumura, K., Habu, S., and Herberman, R.B. (1982). Role of NK cells in the control of metastatic spread and growth of tumor cells in mice. *Int J Cancer* 30, 107-112.

Gorer, P.A., Lyman, S., and Snell, G.D. (1948). Studies on the genetic and antigenic basis of tumour transplantation: linkage between a histocompatibility gene and 'Fused' in mice. *Proc R Soc Lond B* 135, 499-505.

Gossen, M., and Bujard, H. (1992). Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proc Natl Acad Sci U S A* 89, 5547-5551.

Grabarczyk, P., Przybylski, G.K., Depke, M., Volker, U., Bahr, J., Assmus, K., Broker, B.M., Walther, R., and Schmidt, C.A. (2007). Inhibition of BCL11B expression leads to apoptosis of malignant but not normal mature T cells. *Oncogene* 26, 3797-3810.

Graf, T., and Enver, T. (2009). Forcing cells to change lineages. *Nature* 462, 587-594.  
Griffiths, G.M., Tsun, A., and Stinchcombe, J.C. (2010). The immunological synapse: a focal point for endocytosis and exocytosis. *J Cell Biol* 189, 399-406.

Guan, C., Ye, C., Yang, X., and Gao, J. (2010). A review of current large-scale mouse knockout efforts. *Genesis* 48, 73-85.

Guerra, N., Tan, Y.X., Joncker, N.T., Choy, A., Gallardo, F., Xiong, N., Knoblaugh, S., Cado, D., Greenberg, N.M., and Raulet, D.H. (2008). NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. *Immunity* 28, 571-580.

Guo, P., Hirano, M., Herrin, B.R., Li, J., Yu, C., Sadlonova, A., and Cooper, M.D. (2009). Dual nature of the adaptive immune system in lampreys. *Nature* 459, 796-801.

Guo, Y., Maillard, I., Chakraborti, S., Rothenberg, E.V., and Speck, N.A. (2008). Core binding factors are necessary for natural killer cell development and cooperate with Notch signaling during T-cell specification. *Blood* 112, 480-492.

Gutierrez-Avino, F.J., Ferres-Marco, D., and Dominguez, M. (2009). The position and function of the Notch-mediated eye growth organizer: the roles of JAK/STAT and four-jointed. *EMBO Rep* 10, 1051-1058.

Hadjantonakis, A.K., Gertsenstein, M., Ikawa, M., Okabe, M., and Nagy, A. (1998). Generating green fluorescent mice by germline transmission of green fluorescent ES cells. *Mech Dev* 76, 79-90.

Hameyer, D., Loonstra, A., Eshkind, L., Schmitt, S., Antunes, C., Groen, A., Bindels, E., Jonkers, J., Krimpenfort, P., Meuwissen, R., *et al.* (2007). Toxicity of ligand-dependent Cre recombinases and generation of a conditional Cre deleter mouse allowing mosaic recombination in peripheral tissues. *Physiol Genomics* 31, 32-41.

Han, H., Tanigaki, K., Yamamoto, N., Kuroda, K., Yoshimoto, M., Nakahata, T., Ikuta, K., and Honjo, T. (2002). Inducible gene knockout of transcription factor recombination signal binding protein-J reveals its essential role in T versus B lineage decision. *Int Immunol* 14, 637-645.

Hardy, R.R., and Hayakawa, K. (2001). B cell development pathways. *Annu Rev Immunol* 19, 595-621.

Harman, B.C., Jenkinson, E.J., and Anderson, G. (2003). Microenvironmental regulation of Notch signalling in T cell development. *Semin Immunol* 15, 91-97.

Harrison, P.R. (1976). Analysis of erythropoiesis at the molecular level. *Nature* 262, 353-356.

He, X., Park, K., and Kappes, D.J. (2010). The role of ThPOK in control of CD4/CD8 lineage commitment. *Annu Rev Immunol* 28, 295-320.

He, X., Park, K., Wang, H., Zhang, Y., Hua, X., Li, Y., and Kappes, D.J. (2008). CD4-CD8 lineage commitment is regulated by a silencer element at the ThPOK transcription-factor locus. *Immunity* 28, 346-358.

Herberman, R.B., Nunn, M.E., Holden, H.T., and Lavrin, D.H. (1975). Natural cytotoxic reactivity of mouse lymphoid cells against syngeneic and allogeneic tumors. II. Characterization of effector cells. *Int J Cancer* 16, 230-239.

Hickman-Davis, J.M., and Davis, I.C. (2006). Transgenic mice. *Paediatr Respir Rev* 7, 49-53.

Ho, I.C., Tai, T.S., and Pai, S.Y. (2009). GATA3 and the T-cell lineage: essential functions before and after T-helper-2-cell differentiation. *Nat Rev Immunol* 9, 125-135.

Hogan, S.P., and Rothenberg, M.E. (2008). Dietary allergenic proteins and intestinal immunity: a shift from oral tolerance to sensitization. *Clin Exp Allergy* 38, 229-232.

Huang, J., Garrett, K.P., Pelayo, R., Zuniga-Pflucker, J.C., Petrie, H.T., and Kincaide, P.W. (2005). Propensity of adult lymphoid progenitors to progress to DN2/3 stage thymocytes with Notch receptor ligation. *J Immunol* 175, 4858-4865.

Huang, W.Y., Aramburu, J., Douglas, P.S., and Izumo, S. (2000). Transgenic expression of green fluorescence protein can cause dilated cardiomyopathy. *Nat Med* 6, 482-483.

Huntington, N.D., Vosshenrich, C.A., and Di Santo, J.P. (2007). Developmental pathways that generate natural-killer-cell diversity in mice and humans. *Nat Rev Immunol* 7, 703-714.

Iguchi-Manaka, A., Kai, H., Yamashita, Y., Shibata, K., Tahara-Hanaoka, S., Honda, S., Yasui, T., Kikutani, H., Shibuya, K., and Shibuya, A. (2008). Accelerated tumor growth in mice deficient in DNAM-1 receptor. *J Exp Med* 205, 2959-2964.

Ikawa, T., Fujimoto, S., Kawamoto, H., Katsura, Y., and Yokota, Y. (2001). Commitment to natural killer cells requires the helix-loop-helix inhibitor Id2. *Proc Natl Acad Sci U S A* 98, 5164-5169.

Ikawa, T., Hirose, S., Masuda, K., Kakugawa, K., Satoh, R., Shibano-Satoh, A., Kominami, R., Katsura, Y., and Kawamoto, H. (2010). An essential developmental checkpoint for production of the T cell lineage. *Science* 329, 93-96.

Ikawa, T., Kawamoto, H., Fujimoto, S., and Katsura, Y. (1999). Commitment of common T/Natural killer (NK) progenitors to unipotent T and NK progenitors in the

murine fetal thymus revealed by a single progenitor assay. *J Exp Med* *190*, 1617-1626.

Intlekofer, A.M., Takemoto, N., Wherry, E.J., Longworth, S.A., Northrup, J.T., Palanivel, V.R., Mullen, A.C., Gasink, C.R., Kaech, S.M., Miller, J.D., *et al.* (2005). Effector and memory CD8<sup>+</sup> T cell fate coupled by T-bet and eomesodermin. *Nat Immunol* *6*, 1236-1244.

Iwasaki, H., and Akashi, K. (2007). Myeloid lineage commitment from the hematopoietic stem cell. *Immunity* *26*, 726-740.

Izon, D.J., Aster, J.C., He, Y., Weng, A., Karnell, F.G., Patriub, V., Xu, L., Bakkour, S., Rodriguez, C., Allman, D., *et al.* (2002). Deltex1 redirects lymphoid progenitors to the B cell lineage by antagonizing Notch1. *Immunity* *16*, 231-243.

Jahn, T., Sindhu, S., Gooch, S., Seipel, P., Lavori, P., Leifheit, E., and Weinberg, K. (2007). Direct interaction between Kit and the interleukin-7 receptor. *Blood* *110*, 1840-1847.

Kaech, S.M., Tan, J.T., Wherry, E.J., Konieczny, B.T., Surh, C.D., and Ahmed, R. (2003). Selective expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived memory cells. *Nat Immunol* *4*, 1191-1198.

Kamimura, K., Mishima, Y., Obata, M., Endo, T., Aoyagi, Y., and Kominami, R. (2007a). Lack of Bcl11b tumor suppressor results in vulnerability to DNA replication stress and damages. *Oncogene* *26*, 5840-5850.

Kamimura, K., Ohi, H., Kubota, T., Okazuka, K., Yoshikai, Y., Wakabayashi, Y., Aoyagi, Y., Mishima, Y., and Kominami, R. (2007b). Haploinsufficiency of Bcl11b for suppression of lymphomagenesis and thymocyte development. *Biochem Biophys Res Commun* *355*, 538-542.

Kamizono, S., Duncan, G.S., Seidel, M.G., Morimoto, A., Hamada, K., Grosveld, G., Akashi, K., Lind, E.F., Haight, J.P., Ohashi, P.S., *et al.* (2009). Nfil3/E4bp4 is

required for the development and maturation of NK cells in vivo. *J Exp Med* 206, 2977-2986.

Karanam, N.K., Grabarczyk, P., Hammer, E., Scharf, C., Venz, S., Gesell-Salazar, M., Barthlen, W., Przybylski, G.K., Schmidt, C.A., and Volker, U. (2010). Proteome Analysis Reveals New Mechanisms of Bcl11b-loss Driven Apoptosis. *J Proteome Res* [Epub ahead of print].

Karasuyama, H., Kudo, A., and Melchers, F. (1990). The proteins encoded by the VpreB and lambda 5 pre-B cell-specific genes can associate with each other and with mu heavy chain. *J Exp Med* 172, 969-972.

Karasuyama, H., Rolink, A., and Melchers, F. (1993). A complex of glycoproteins is associated with VpreB/lambda 5 surrogate light chain on the surface of mu heavy chain-negative early precursor B cell lines. *J Exp Med* 178, 469-478.

Karre, K. (2008). Natural killer cell recognition of missing self. *Nat Immunol* 9, 477-480.

Karre, K., Ljunggren, H.G., Piontek, G., and Kiessling, R. (1986). Selective rejection of H-2-deficient lymphoma variants suggests alternative immune defence strategy. *Nature* 319, 675-678.

Kastner, P., Chan, S., Vogel, W.K., Zhang, L.J., Topark-Ngarm, A., Golonzhka, O., Jost, B., Le Gras, S., Gross, M.K., and Leid, M. (2010). Bcl11b represses a mature T-cell gene expression program in immature CD4(+)CD8(+) thymocytes. *Eur J Immunol*.

Kiessling, R., Haller, O., Fenyo, E.M., Steinitz, M., and Klein, G. (1978). Mouse natural killer (NK) cell activity against human cell lines is not influenced by superinfection of the target cell with xenotropic murine C-type virus. *Int J Cancer* 21, 460-465.



Kiessling, R., Klein, E., Pross, H., and Wigzell, H. (1975). "Natural" killer cells in the mouse. II. Cytotoxic cells with specificity for mouse Moloney leukemia cells. Characteristics of the killer cell. *Eur J Immunol* 5, 117-121.

Kiessling, R., Petrányi, G., Karre, K., Jondal, M., Tracey, D., and Wigzell, H. (1976). Killer cells: a functional comparison between natural, immune T-cell and antibody-dependent in vitro systems. *J Exp Med* 143, 772-780.

Kim, J.B., Greber, B., Arauzo-Bravo, M.J., Meyer, J., Park, K.I., Zaehres, H., and Scholer, H.R. (2009). Direct reprogramming of human neural stem cells by OCT4. *Nature* 461, 649-643.

Kim, S., Iizuka, K., Kang, H.S., Dokun, A., French, A.R., Greco, S., and Yokoyama, W.M. (2002). In vivo developmental stages in murine natural killer cell maturation. *Nat Immunol* 3, 523-528.

Kitamura, D., Roes, J., Kuhn, R., and Rajewsky, K. (1991). A B cell-deficient mouse by targeted disruption of the membrane exon of the immunoglobulin mu chain gene. *Nature* 350, 423-426.

Kleinjan, D.A., and van Heyningen, V. (2005). Long-range control of gene expression: emerging mechanisms and disruption in disease. *Am J Hum Genet* 76, 8-32.

Klug, A. (2010). The discovery of zinc fingers and their applications in gene regulation and genome manipulation. *Annu Rev Biochem* 79, 213-231.

Koller, B.H., Marrack, P., Kappler, J.W., and Smithies, O. (1990). Normal development of mice deficient in beta 2M, MHC class I proteins, and CD8<sup>+</sup> T cells. *Science* 248, 1227-1230.

Koller, B.H., and Smithies, O. (1989). Inactivating the beta 2-microglobulin locus in mouse embryonic stem cells by homologous recombination. *Proc Natl Acad Sci U S A* 86, 8932-8935.

Kondo, M., Weissman, I.L., and Akashi, K. (1997). Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell* 91, 661-672.

Kopan, R., and Ilagan, M.X. (2009). The canonical Notch signaling pathway: unfolding the activation mechanism. *Cell* 137, 216-233.

Krejci, A., Bernard, F., Housden, B.E., Collins, S., and Bray, S.J. (2009). Direct response to Notch activation: signaling crosstalk and incoherent logic. *Sci Signal* 2, ra1.

Kuprash, D.V., Alimzhanov, M.B., Tumanov, A.V., Grivennikov, S.I., Shakhov, A.N., Drutskaya, L.N., Marino, M.W., Turetskaya, R.L., Anderson, A.O., Rajewsky, K., *et al.* (2002). Redundancy in tumor necrosis factor (TNF) and lymphotoxin (LT) signaling in vivo: mice with inactivation of the entire TNF/LT locus versus single-knockout mice. *Mol Cell Biol* 22, 8626-8634.

Lakshmikanth, T., Burke, S., Ali, T.H., Kimpfler, S., Ursini, F., Ruggeri, L., Capanni, M., Umansky, V., Paschen, A., Sucker, A., *et al.* (2009). NCRs and DNAM-1 mediate NK cell recognition and lysis of human and mouse melanoma cell lines in vitro and in vivo. *J Clin Invest* 119, 1251-1263.

Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., *et al.* (2001). Initial sequencing and analysis of the human genome. *Nature* 409, 860-921.

Laouar, Y., Sutterwala, F.S., Gorelik, L., and Flavell, R.A. (2005). Transforming growth factor-beta controls T helper type 1 cell development through regulation of natural killer cell interferon-gamma. *Nat Immunol* 6, 600-607.

Lee, D.H., Iwanski, G.B., and Thoennissen, N.H. (2010). Cucurbitacin: ancient compound shedding new light on cancer treatment. *ScientificWorldJournal* 10, 413-418.

Lee, P.P., Fitzpatrick, D.R., Beard, C., Jessup, H.K., Lehar, S., Makar, K.W., Perez-Melgosa, M., Sweetser, M.T., Schlissel, M.S., Nguyen, S., *et al.* (2001). A critical role for Dnmt1 and DNA methylation in T cell development, function, and survival. *Immunity* 15, 763-774.

Lewis, R.S. (2001). Calcium signaling mechanisms in T lymphocytes. *Annu Rev Immunol* 19, 497-521.

Li, L., Leid, M., and Rothenberg, E.V. (2010). An early T cell lineage commitment checkpoint dependent on the transcription factor Bcl11b. *Science* 329, 89-93.

Lin, H., and Grosschedl, R. (1995). Failure of B-cell differentiation in mice lacking the transcription factor EBF. *Nature* 376, 263-267.

Liu, P., Keller, J.R., Ortiz, M., Tessarollo, L., Rachel, R.A., Nakamura, T., Jenkins, N.A., and Copeland, N.G. (2003). Bcl11a is essential for normal lymphoid development. *Nat Immunol* 4, 525-532.

Ljunggren, H.G., and Karre, K. (1985). Host resistance directed selectively against H-2-deficient lymphoma variants. Analysis of the mechanism. *J Exp Med* 162, 1745-1759.

Lodolce, J.P., Boone, D.L., Chai, S., Swain, R.E., Dassopoulos, T., Trettin, S., and Ma, A. (1998). IL-15 receptor maintains lymphoid homeostasis by supporting lymphocyte homing and proliferation. *Immunity* 9, 669-676.

M. Bielschowsky, B.J.H.a.J.H. (1959). Spontaneous haemolytic anaemia in mice of the NZB/B1 strain. *Proc Univ of Otago Med School* 37, 9-11.

MacLeod, R.A., Nagel, S., Kaufmann, M., Janssen, J.W., and Drexler, H.G. (2003). Activation of HOX11L2 by juxtaposition with 3'-BCL11B in an acute lymphoblastic leukemia cell line (HPB-ALL) with t(5;14)(q35;q32.2). *Genes Chromosomes Cancer* 37, 84-91.

Maillard, I., Fang, T., and Pear, W.S. (2005). Regulation of lymphoid development, differentiation, and function by the Notch pathway. *Annu Rev Immunol* 23, 945-974.

Mak, T.W., Hakem, A., McPherson, J.P., Shehabeldin, A., Zabolcki, E., Migon, E., Duncan, G.S., Bouchard, D., Wakeham, A., Cheung, A., *et al.* (2000). Brcal required for T cell lineage development but not TCR loci rearrangement. *Nat Immunol* 1, 77-82.

Mak, T.W., Penninger, J.M., and Ohashi, P.S. (2001). Knockout mice: a paradigm shift in modern immunology. *Nat Rev Immunol* 1, 11-19.

Manz, M.G., Traver, D., Miyamoto, T., Weissman, I.L., and Akashi, K. (2001). Dendritic cell potentials of early lymphoid and myeloid progenitors. *Blood* 97, 3333-3341.

Marban, C., Suzanne, S., Dequiedt, F., de Walque, S., Redel, L., Van Lint, C., Aunis, D., and Rohr, O. (2007). Recruitment of chromatin-modifying enzymes by CTIP2 promotes HIV-1 transcriptional silencing. *EMBO J* 26, 412-423.

Mardis, E.R. (2008). The impact of next-generation sequencing technology on genetics. *Trends Genet* 24, 133-141.

Marsland, B.J., and Kopf, M. (2008). T-cell fate and function: PKC-theta and beyond. *Trends Immunol* 29, 179-185.

Martin, G.R. (1981). Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc Natl Acad Sci U S A* 78, 7634-7638.

Masuda, K., Kakugawa, K., Nakayama, T., Minato, N., Katsura, Y., and Kawamoto, H. (2007). T cell lineage determination precedes the initiation of TCR beta gene rearrangement. *J Immunol* 179, 3699-3706.

Matsumoto, Y., Kosugi, S., Shinbo, T., Chou, D., Ohashi, M., Wakabayashi, Y., Sakai, K., Okumoto, M., Mori, N., Aizawa, S., *et al.* (1998). Allelic loss analysis of gamma-ray-induced mouse thymic lymphomas: two candidate tumor suppressor gene loci on chromosomes 12 and 16. *Oncogene* *16*, 2747-2754.

Menzel, S., Garner, C., Gut, I., Matsuda, F., Yamaguchi, M., Heath, S., Foglio, M., Zelenika, D., Boland, A., Rooks, H., *et al.* (2007). A QTL influencing F cell production maps to a gene encoding a zinc-finger protein on chromosome 2p15. *Nat Genet* *39*, 1197-1199.

Metzger, D., and Chambon, P. (2001). Site- and time-specific gene targeting in the mouse. *Methods* *24*, 71-80.

Miaw, S.C., Choi, A., Yu, E., Kishikawa, H., and Ho, I.C. (2000). ROG, repressor of GATA, regulates the expression of cytokine genes. *Immunity* *12*, 323-333.

Michie, A.M., Carlyle, J.R., Schmitt, T.M., Ljutic, B., Cho, S.K., Fong, Q., and Zuniga-Pflucker, J.C. (2000). Clonal characterization of a bipotent T cell and NK cell progenitor in the mouse fetal thymus. *J Immunol* *164*, 1730-1733.

Mikkola, I., Heavey, B., Horcher, M., and Busslinger, M. (2002). Reversion of B cell commitment upon loss of Pax5 expression. *Science* *297*, 110-113.

Miller, J.S., Soignier, Y., Panoskaltsis-Mortari, A., McNearney, S.A., Yun, G.H., Fautsch, S.K., McKenna, D., Le, C., Defor, T.E., Burns, L.J., *et al.* (2005). Successful adoptive transfer and in vivo expansion of human haploidentical NK cells in patients with cancer. *Blood* *105*, 3051-3057.

Mlodzik, M., Hiromi, Y., Weber, U., Goodman, C.S., and Rubin, G.M. (1990). The *Drosophila* seven-up gene, a member of the steroid receptor gene superfamily, controls photoreceptor cell fates. *Cell* *60*, 211-224.

Mohrs, M., Shinkai, K., Mohrs, K., and Locksley, R.M. (2001). Analysis of type 2 immunity in vivo with a bicistronic IL-4 reporter. *Immunity* *15*, 303-311.

Morris, L., Klanke, C., Lang, S., Lim, F.Y., and Crombleholme, T. (2010). TdTomato and EGFP identification in histological sections: insight and alternatives. *Biotech Histochem*.

Morrison, S.J., and Weissman, I.L. (1994). The long-term repopulating subset of hematopoietic stem cells is deterministic and isolatable by phenotype. *Immunity* *1*, 661-673.

Muul, L.M., Spiess, P.J., Director, E.P., and Rosenberg, S.A. (1987). Identification of specific cytolytic immune responses against autologous tumor in humans bearing malignant melanoma. *J Immunol* *138*, 989-995.

Nagel, S., Kaufmann, M., Drexler, H.G., and MacLeod, R.A. (2003). The cardiac homeobox gene NKX2-5 is deregulated by juxtaposition with BCL11B in pediatric T-ALL cell lines via a novel t(5;14)(q35.1;q32.2). *Cancer Res* *63*, 5329-5334.

Nagy, A. (2000). Cre recombinase: the universal reagent for genome tailoring. *Genesis* *26*, 99-109.

Nagy, A., Rossant, J., Nagy, R., Abramow-Newerly, W., and Roder, J.C. (1993). Derivation of completely cell culture-derived mice from early-passage embryonic stem cells. *Proc Natl Acad Sci U S A* *90*, 8424-8428.

Nakamura, T., Largaespada, D.A., Shaughnessy, J.D., Jr., Jenkins, N.A., and Copeland, N.G. (1996). Cooperative activation of Hoxa and Pbx1-related genes in murine myeloid leukaemias. *Nat Genet* *12*, 149-153.

Nakamura, T., Yamazaki, Y., Saiki, Y., Moriyama, M., Largaespada, D.A., Jenkins, N.A., and Copeland, N.G. (2000). Evi9 encodes a novel zinc finger protein that physically interacts with BCL6, a known human B-cell proto-oncogene product. *Mol Cell Biol* *20*, 3178-3186.

Nakano, T., Kodama, H., and Honjo, T. (1994). Generation of lymphohematopoietic cells from embryonic stem cells in culture. *Science* 265, 1098-1101.

Nelms, K.A., and Goodnow, C.C. (2001). Genome-wide ENU mutagenesis to reveal immune regulators. *Immunity* 15, 409-418.

Nikolic-Zugic, J., and Moore, M.W. (1989). T cell receptor expression on immature thymocytes with in vivo and in vitro precursor potential. *Eur J Immunol* 19, 1957-1960.

Nolan, G.P., Fiering, S., Nicolas, J.F., and Herzenberg, L.A. (1988). Fluorescence-activated cell analysis and sorting of viable mammalian cells based on beta-D-galactosidase activity after transduction of *Escherichia coli lacZ*. *Proc Natl Acad Sci U S A* 85, 2603-2607.

Northrup, D.L., and Allman, D. (2008). Transcriptional regulation of early B cell development. *Immunol Res* 42, 106-117.

Okazuka, K., Wakabayashi, Y., Kashihara, M., Inoue, J., Sato, T., Yokoyama, M., Aizawa, S., Aizawa, Y., Mishima, Y., and Kominami, R. (2005). p53 prevents maturation of T cell development to the immature CD4-CD8<sup>+</sup> stage in *Bcl11b*<sup>-/-</sup> mice. *Biochem Biophys Res Commun* 328, 545-549.

Oldham, R.K. (1983). Natural killer cells: artifact to reality: an odyssey in biology. *Cancer Metastasis Rev* 2, 323-336.

Omori, M., Yamashita, M., Inami, M., Ukai-Tadenuma, M., Kimura, M., Nigo, Y., Hosokawa, H., Hasegawa, A., Taniguchi, M., and Nakayama, T. (2003). CD8 T cell-specific downregulation of histone hyperacetylation and gene activation of the IL-4 gene locus by ROG, repressor of GATA. *Immunity* 19, 281-294.

Pai, S.Y., Truitt, M.L., Ting, C.N., Leiden, J.M., Glimcher, L.H., and Ho, I.C. (2003). Critical roles for transcription factor GATA-3 in thymocyte development. *Immunity* 19, 863-875.

Park, K., He, X., Lee, H.O., Hua, X., Li, Y., Wiest, D., and Kappes, D.J. (2010). TCR-mediated ThPOK induction promotes development of mature (CD24-) gammadelta thymocytes. *EMBO J* 29, 2329-2341.

Pearce, E.L., Mullen, A.C., Martins, G.A., Krawczyk, C.M., Hutchins, A.S., Zediak, V.P., Banica, M., DiCioccio, C.B., Gross, D.A., Mao, C.A., *et al.* (2003). Control of effector CD8<sup>+</sup> T cell function by the transcription factor Eomesodermin. *Science* 302, 1041-1043.

Pellicci, D.G., Hammond, K.J., Uldrich, A.P., Baxter, A.G., Smyth, M.J., and Godfrey, D.I. (2002). A natural killer T (NKT) cell developmental pathway involving a thymus-dependent NK1.1(-)CD4(+) CD1d-dependent precursor stage. *J Exp Med* 195, 835-844.

Pentao Liu, P.L., Shannon Burke (2010). Critical Roles of Bcl11b in T Cell Development and Maintenance of T Cell Identity. *Immunological Reviews* *In press*.

Petrie, H.T., Hugo, P., Scollay, R., and Shortman, K. (1990). Lineage relationships and developmental kinetics of immature thymocytes: CD3, CD4, and CD8 acquisition in vivo and in vitro. *J Exp Med* 172, 1583-1588.

Przybylski, G.K., Dik, W.A., Wanzeck, J., Grabarczyk, P., Majunke, S., Martin-Subero, J.I., Siebert, R., Dolken, G., Ludwig, W.D., Verhaaf, B., *et al.* (2005). Disruption of the BCL11B gene through inv(14)(q11.2q32.31) results in the expression of BCL11B-TRDC fusion transcripts and is associated with the absence of wild-type BCL11B transcripts in T-ALL. *Leukemia* 19, 201-208.

Pui, J.C., Allman, D., Xu, L., DeRocco, S., Karnell, F.G., Bakkour, S., Lee, J.Y., Kadesch, T., Hardy, R.R., Aster, J.C., *et al.* (1999). Notch1 expression in early lymphopoiesis influences B versus T lineage determination. *Immunity* 11, 299-308.



Puzanov, I.J., Bennett, M., and Kumar, V. (1996). IL-15 can substitute for the marrow microenvironment in the differentiation of natural killer cells. *J Immunol* *157*, 4282-4285.

Qi, Q., and August, A. (2007). Keeping the (kinase) party going: SLP-76 and ITK dance to the beat. *Sci STKE* *2007*, pe39.

Radtke, F., Fasnacht, N., and Macdonald, H.R. (2010). Notch signaling in the immune system. *Immunity* *32*, 14-27.

Radtke, F., Wilson, A., Stark, G., Bauer, M., van Meerwijk, J., MacDonald, H.R., and Aguet, M. (1999). Deficient T cell fate specification in mice with an induced inactivation of Notch1. *Immunity* *10*, 547-558.

Rajewsky, K., Gu, H., Kuhn, R., Betz, U.A., Muller, W., Roes, J., and Schwenk, F. (1996). Conditional gene targeting. *J Clin Invest* *98*, 600-603.

Ramalho-Santos, M., Yoon, S., Matsuzaki, Y., Mulligan, R.C., and Melton, D.A. (2002). "Stemness": transcriptional profiling of embryonic and adult stem cells. *Science* *298*, 597-600.

Robbins, P.F., Dudley, M.E., Wunderlich, J., El-Gamil, M., Li, Y.F., Zhou, J., Huang, J., Powell, D.J., Jr., and Rosenberg, S.A. (2004). Cutting edge: persistence of transferred lymphocyte clonotypes correlates with cancer regression in patients receiving cell transfer therapy. *J Immunol* *173*, 7125-7130.

Robertson, E., Bradley, A., Kuehn, M., and Evans, M. (1986). Germ-line transmission of genes introduced into cultured pluripotential cells by retroviral vector. *Nature* *323*, 445-448.

Robertson, M.J. (2002). Role of chemokines in the biology of natural killer cells. *J Leukoc Biol* *71*, 173-183.

Rolink, A.G., Balciunaite, G., Demoliere, C., and Ceredig, R. (2006). The potential involvement of Notch signaling in NK cell development. *Immunol Lett* 107, 50-57.

Rolink, A.G., Nutt, S.L., Melchers, F., and Busslinger, M. (1999). Long-term in vivo reconstitution of T-cell development by Pax5-deficient B-cell progenitors. *Nature* 401, 603-606.

Rosenberg, S.A., and Dudley, M.E. (2009). Adoptive cell therapy for the treatment of patients with metastatic melanoma. *Curr Opin Immunol* 21, 233-240.

Rosenberg, S.A., Lotze, M.T., Muul, L.M., Leitman, S., Chang, A.E., Ettinghausen, S.E., Matory, Y.L., Skibber, J.M., Shiloni, E., Vetto, J.T., *et al.* (1985). Observations on the systemic administration of autologous lymphokine-activated killer cells and recombinant interleukin-2 to patients with metastatic cancer. *N Engl J Med* 313, 1485-1492.

Rosenberg, S.A., Restifo, N.P., Yang, J.C., Morgan, R.A., and Dudley, M.E. (2008). Adoptive cell transfer: a clinical path to effective cancer immunotherapy. *Nat Rev Cancer* 8, 299-308.

Rosenberg, S.A., Spiess, P., and Lafreniere, R. (1986). A new approach to the adoptive immunotherapy of cancer with tumor-infiltrating lymphocytes. *Science* 233, 1318-1321.

Rosenberg, S.A., Yang, J.C., and Restifo, N.P. (2004). Cancer immunotherapy: moving beyond current vaccines. *Nat Med* 10, 909-915.

Rosenberg, S.A., Yang, J.C., White, D.E., and Steinberg, S.M. (1998). Durability of complete responses in patients with metastatic cancer treated with high-dose interleukin-2: identification of the antigens mediating response. *Ann Surg* 228, 307-319.

Rosenwasser, L.J., and Rothenberg, M.E. (2010). IL-5 pathway inhibition in the treatment of asthma and Churg-Strauss syndrome. *J Allergy Clin Immunol* 125, 1245-1246.

Rothenberg, E.V. (2007). Negotiation of the T lineage fate decision by transcription-factor interplay and microenvironmental signals. *Immunity* 26, 690-702.

Rothenberg, E.V., Moore, J.E., and Yui, M.A. (2008). Launching the T-cell-lineage developmental programme. *Nat Rev Immunol* 8, 9-21.

Rothenberg, E.V., and Taghon, T. (2005). Molecular genetics of T cell development. *Annu Rev Immunol* 23, 601-649.

Ruggeri, L., Capanni, M., Urbani, E., Perruccio, K., Shlomchik, W.D., Tosti, A., Posati, S., Rogaia, D., Frassoni, F., Aversa, F., *et al.* (2002). Effectiveness of donor natural killer cell alloreactivity in mismatched hematopoietic transplants. *Science* 295, 2097-2100.

Russell, W.L., Kelly, E.M., Hunsicker, P.R., Bangham, J.W., Maddux, S.C., and Phipps, E.L. (1979). Specific-locus test shows ethylnitrosourea to be the most potent mutagen in the mouse. *Proc Natl Acad Sci U S A* 76, 5818-5819.

Salem, M.L., El-Demellawy, M., and El-Azm, A.R. (2010). The potential use of Toll-like receptor agonists to restore the dysfunctional immunity induced by hepatitis C virus. *Cell Immunol* 262, 96-104.

Sankaran, V.G., Menne, T.F., Xu, J., Akie, T.E., Lettre, G., Van Handel, B., Mikkola, H.K., Hirschhorn, J.N., Cantor, A.B., and Orkin, S.H. (2008). Human fetal hemoglobin expression is regulated by the developmental stage-specific repressor BCL11A. *Science* 322, 1839-1842.

Sankaran, V.G., Xu, J., Ragoczy, T., Ippolito, G.C., Walkley, C.R., Maika, S.D., Fujiwara, Y., Ito, M., Groudine, M., Bender, M.A., *et al.* (2009). Developmental and species-divergent globin switching are driven by BCL11A. *Nature* 460, 1093-1097.

Satterwhite, E., Sonoki, T., Willis, T.G., Harder, L., Nowak, R., Arriola, E.L., Liu, H., Price, H.P., Gesk, S., Steinemann, D., *et al.* (2001). The BCL11 gene family: involvement of BCL11A in lymphoid malignancies. *Blood* 98, 3413-3420.

Schmitt, T.M., and Zuniga-Pflucker, J.C. (2002). Induction of T cell development from hematopoietic progenitor cells by delta-like-1 in vitro. *Immunity* 17, 749-756.

Senawong, T., Peterson, V.J., Avram, D., Shepherd, D.M., Frye, R.A., Minucci, S., and Leid, M. (2003). Involvement of the histone deacetylase SIRT1 in chicken ovalbumin upstream promoter transcription factor (COUP-TF)-interacting protein 2-mediated transcriptional repression. *J Biol Chem* 278, 43041-43050.

Senawong, T., Peterson, V.J., and Leid, M. (2005). BCL11A-dependent recruitment of SIRT1 to a promoter template in mammalian cells results in histone deacetylation and transcriptional repression. *Arch Biochem Biophys* 434, 316-325.

Setoguchi, R., Tachibana, M., Naoe, Y., Muroi, S., Akiyama, K., Tezuka, C., Okuda, T., and Taniuchi, I. (2008). Repression of the transcription factor Th-POK by Runx complexes in cytotoxic T cell development. *Science* 319, 822-825.

Shaner, N.C., Campbell, R.E., Steinbach, P.A., Giepmans, B.N., Palmer, A.E., and Tsien, R.Y. (2004). Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein. *Nat Biotechnol* 22, 1567-1572.

Shinbo, T., Matsuki, A., Matsumoto, Y., Kosugi, S., Takahashi, Y., Niwa, O., and Kominami, R. (1999). Allelic loss mapping and physical delineation of a region harboring a putative thymic lymphoma suppressor gene on mouse chromosome 12. *Oncogene* 18, 4131-4136.

Silver, L.M. (1985). *Mouse Genetics, Concepts and Applications*. Oxford University Press.

Smithies, O., Gregg, R.G., Boggs, S.S., Koralewski, M.A., and Kucherlapati, R.S. (1985). Insertion of DNA sequences into the human chromosomal beta-globin locus by homologous recombination. *Nature* 317, 230-234.

Smyth, M.J., Dunn, G.P., and Schreiber, R.D. (2006). Cancer immunosurveillance and immunoediting: the roles of immunity in suppressing tumor development and shaping tumor immunogenicity. *Adv Immunol* 90, 1-50.

Snell, G.D., and Reed, S. (1993). William Ernest Castle, pioneer mammalian geneticist. *Genetics* 133, 751-753.

Snell, G.D.a.H., G.F. (1951). Alleles at the histocompatibility-2 locus in the mouse as determined by tumor transplantation. *Genetics* 36, 306-310.

Sotillos, S., Diaz-Meco, M.T., Moscat, J., and Castelli-Gair Hombria, J. (2008). Polarized subcellular localization of Jak/STAT components is required for efficient signaling. *Curr Biol* 18, 624-629.

Spits, H., Couwenberg, F., Bakker, A.Q., Weijer, K., and Uittenbogaart, C.H. (2000). Id2 and Id3 inhibit development of CD34(+) stem cells into predendritic cell (pre-DC)2 but not into pre-DC1. Evidence for a lymphoid origin of pre-DC2. *J Exp Med* 192, 1775-1784.

Stinchcombe, J.C., Majorovits, E., Bossi, G., Fuller, S., and Griffiths, G.M. (2006). Centrosome polarization delivers secretory granules to the immunological synapse. *Nature* 443, 462-465.

Stremmel, C., Exley, M., Balk, S., Hohenberger, W., and Kuchroo, V.K. (2001). Characterization of the phenotype and function of CD8(+), alpha / beta(+) NKT cells from tumor-bearing mice that show a natural killer cell activity and lyse multiple tumor targets. *Eur J Immunol* 31, 2818-2828.

Suzuki, A., Yamaguchi, M.T., Ohteki, T., Sasaki, T., Kaisho, T., Kimura, Y., Yoshida, R., Wakeham, A., Higuchi, T., Fukumoto, M., *et al.* (2001). T cell-specific

loss of Pten leads to defects in central and peripheral tolerance. *Immunity* *14*, 523-534.

Suzuki, H., Duncan, G.S., Takimoto, H., and Mak, T.W. (1997). Abnormal development of intestinal intraepithelial lymphocytes and peripheral natural killer cells in mice lacking the IL-2 receptor beta chain. *J Exp Med* *185*, 499-505.

Szabo, S.J., Kim, S.T., Costa, G.L., Zhang, X., Fathman, C.G., and Glimcher, L.H. (2000). A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* *100*, 655-669.

Szabo, S.J., Sullivan, B.M., Stemmann, C., Satoskar, A.R., Sleckman, B.P., and Glimcher, L.H. (2002). Distinct effects of T-bet in TH1 lineage commitment and IFN-gamma production in CD4 and CD8 T cells. *Science* *295*, 338-342.

Taghon, T., Yui, M.A., Pant, R., Diamond, R.A., and Rothenberg, E.V. (2006). Developmental and molecular characterization of emerging beta- and gammadelta-selected pre-T cells in the adult mouse thymus. *Immunity* *24*, 53-64.

Taghon, T.N., David, E.S., Zuniga-Pflucker, J.C., and Rothenberg, E.V. (2005). Delayed, asynchronous, and reversible T-lineage specification induced by Notch/Delta signaling. *Genes Dev* *19*, 965-978.

Takada, K., and Jameson, S.C. (2009). Naive T cell homeostasis: from awareness of space to a sense of place. *Nat Rev Immunol* *9*, 823-832.

Takahashi, K., Tanabe, K., Ohnuki, M., Narita, M., Ichisaka, T., Tomoda, K., and Yamanaka, S. (2007). Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* *131*, 861-872.

Takahashi, M., Altschmied, L., and Hillen, W. (1986). Kinetic and equilibrium characterization of the Tet repressor-tetracycline complex by fluorescence measurements. Evidence for divalent metal ion requirement and energy transfer. *J Mol Biol* *187*, 341-348.

Takeda, K., Cretney, E., Hayakawa, Y., Ota, T., Akiba, H., Ogasawara, K., Yagita, H., Kinoshita, K., Okumura, K., and Smyth, M.J. (2005). TRAIL identifies immature natural killer cells in newborn mice and adult mouse liver. *Blood* *105*, 2082-2089.

Taniguchi, S., Kawano, T., Kakunaga, T., and Baba, T. (1986). Differences in expression of a variant actin between low and high metastatic B16 melanoma. *J Biol Chem* *261*, 6100-6106.

Taniuchi, I., Osato, M., Egawa, T., Sunshine, M.J., Bae, S.C., Komori, T., Ito, Y., and Littman, D.R. (2002). Differential requirements for Runx proteins in CD4 repression and epigenetic silencing during T lymphocyte development. *Cell* *111*, 621-633.

Telling, G.C., Tremblay, P., Torchia, M., Dearmond, S.J., Cohen, F.E., and Prusiner, S.B. (1997). N-terminally tagged prion protein supports prion propagation in transgenic mice. *Protein Sci* *6*, 825-833.

Thibault, S.T., Singer, M.A., Miyazaki, W.Y., Milash, B., Dompe, N.A., Singh, C.M., Buchholz, R., Demsky, M., Fawcett, R., Francis-Lang, H.L., *et al.* (2004). A complementary transposon tool kit for *Drosophila melanogaster* using P and piggyBac. *Nat Genet* *36*, 283-287.

Thomas, K.R., and Capecchi, M.R. (1986). Introduction of homologous DNA sequences into mammalian cells induces mutations in the cognate gene. *Nature* *324*, 34-38.

Topark-Ngarm, A., Golonzhka, O., Peterson, V.J., Barrett, B., Jr., Martinez, B., Crofoot, K., Filtz, T.M., and Leid, M. (2006). CTIP2 associates with the NuRD complex on the promoter of p57KIP2, a newly identified CTIP2 target gene. *J Biol Chem* *281*, 32272-32283.

Triezenberg, S.J., Kingsbury, R.C., and McKnight, S.L. (1988). Functional dissection of VP16, the trans-activator of herpes simplex virus immediate early gene expression. *Genes Dev* *2*, 718-729.

Trono, D., Van Lint, C., Rouzioux, C., Verdin, E., Barre-Sinoussi, F., Chun, T.W., and Chomont, N. (2010). HIV persistence and the prospect of long-term drug-free remissions for HIV-infected individuals. *Science* 329, 174-180.

Tun, T., Hamaguchi, Y., Matsunami, N., Furukawa, T., Honjo, T., and Kawaichi, M. (1994). Recognition sequence of a highly conserved DNA binding protein RBP-J kappa. *Nucleic Acids Res* 22, 965-971.

Turman, M.A., Yabe, T., McSherry, C., Bach, F.H., and Houchins, J.P. (1993). Characterization of a novel gene (NKG7) on human chromosome 19 that is expressed in natural killer cells and T cells. *Hum Immunol* 36, 34-40.

Tydell, C.C., David-Fung, E.S., Moore, J.E., Rowen, L., Taghon, T., and Rothenberg, E.V. (2007). Molecular dissection of prethymic progenitor entry into the T lymphocyte developmental pathway. *J Immunol* 179, 421-438.

Uda, M., Galanello, R., Sanna, S., Lettre, G., Sankaran, V.G., Chen, W., Usala, G., Busonero, F., Maschio, A., Albai, G., *et al.* (2008). Genome-wide association study shows BCL11A associated with persistent fetal hemoglobin and amelioration of the phenotype of beta-thalassemia. *Proc Natl Acad Sci U S A* 105, 1620-1625.

Urbanek, P., Wang, Z.Q., Fetka, I., Wagner, E.F., and Busslinger, M. (1994). Complete block of early B cell differentiation and altered patterning of the posterior midbrain in mice lacking Pax5/BSAP. *Cell* 79, 901-912.

Urnov, F.D., Miller, J.C., Lee, Y.L., Beausejour, C.M., Rock, J.M., Augustus, S., Jamieson, A.C., Porteus, M.H., Gregory, P.D., and Holmes, M.C. (2005). Highly efficient endogenous human gene correction using designed zinc-finger nucleases. *Nature* 435, 646-651.

van der Bruggen, P., Traversari, C., Chomez, P., Lurquin, C., De Plaen, E., Van den Eynde, B., Knuth, A., and Boon, T. (1991). A gene encoding an antigen recognized by cytolytic T lymphocytes on a human melanoma. *Science* 254, 1643-1647.



van der Wees, J., Matharu, P.J., de Roos, K., Destree, O.H., Godsave, S.F., Durston, A.J., and Sweeney, G.E. (1996). Developmental expression and differential regulation by retinoic acid of *Xenopus* COUP-TF-A and COUP-TF-B. *Mech Dev* 54, 173-184.

Vegh, P., Winckler, J., and Melchers, F. (2010). Long-term "in vitro" proliferating mouse hematopoietic progenitor cell lines. *Immunol Lett* 130, 32-35.

Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., Smith, H.O., Yandell, M., Evans, C.A., Holt, R.A., *et al.* (2001). The sequence of the human genome. *Science* 291, 1304-1351.

Vosshenrich, C.A., Garcia-Ojeda, M.E., Samson-Villeger, S.I., Pasqualetto, V., Enault, L., Richard-Le Goff, O., Corcuff, E., Guy-Grand, D., Rocha, B., Cumano, A., *et al.* (2006). A thymic pathway of mouse natural killer cell development characterized by expression of GATA-3 and CD127. *Nat Immunol* 7, 1217-1224.

Wada, H., Masuda, K., Satoh, R., Kakugawa, K., Ikawa, T., Katsura, Y., and Kawamoto, H. (2008). Adult T-cell progenitors retain myeloid potential. *Nature* 452, 768-772.

Wagner, E.F., Stewart, T.A., and Mintz, B. (1981a). The human beta-globin gene and a functional viral thymidine kinase gene in developing mice. *Proc Natl Acad Sci U S A* 78, 5016-5020.

Wagner, T.E., Hoppe, P.C., Jollick, J.D., Scholl, D.R., Hodinka, R.L., and Gault, J.B. (1981b). Microinjection of a rabbit beta-globin gene into zygotes and its subsequent expression in adult mice and their offspring. *Proc Natl Acad Sci U S A* 78, 6376-6380.

Wakabayashi, Y., Inoue, J., Takahashi, Y., Matsuki, A., Kosugi-Okano, H., Shinbo, T., Mishima, Y., Niwa, O., and Kominami, R. (2003a). Homozygous deletions and point mutations of the *Rit1/Bcl11b* gene in gamma-ray induced mouse thymic lymphomas. *Biochem Biophys Res Commun* 301, 598-603.

Wakabayashi, Y., Watanabe, H., Inoue, J., Takeda, N., Sakata, J., Mishima, Y., Hitomi, J., Yamamoto, T., Utsuyama, M., Niwa, O., *et al.* (2003b). Bcl11b is required for differentiation and survival of alphabeta T lymphocytes. *Nat Immunol* 4, 533-539.

Waldmann, T.A. (2006). The biology of interleukin-2 and interleukin-15: implications for cancer therapy and vaccine design. *Nat Rev Immunol* 6, 595-601.

Waldmann, T.A., and Tagaya, Y. (1999). The multifaceted regulation of interleukin-15 expression and the role of this cytokine in NK cell differentiation and host response to intracellular pathogens. *Annu Rev Immunol* 17, 19-49.

Wallace, H.A., Marques-Kranc, F., Richardson, M., Luna-Crespo, F., Sharpe, J.A., Hughes, J., Wood, W.G., Higgs, D.R., and Smith, A.J. (2007). Manipulating the mouse genome to engineer precise functional syntenic replacements with human sequence. *Cell* 128, 197-209.

Walzer, T., Blery, M., Chaix, J., Fuseri, N., Chasson, L., Robbins, S.H., Jaeger, S., Andre, P., Gauthier, L., Daniel, L., *et al.* (2007). Identification, activation, and selective in vivo ablation of mouse NK cells via NKp46. *Proc Natl Acad Sci U S A* 104, 3384-3389.

Wang, D., Claus, C.L., Vaccarelli, G., Braunstein, M., Schmitt, T.M., Zuniga-Pflucker, J.C., Rothenberg, E.V., and Anderson, M.K. (2006). The basic helix-loop-helix transcription factor HEBAlt is expressed in pro-T cells and enhances the generation of T cell precursors. *J Immunol* 177, 109-119.

Wang, H., Feng, J., Qi, C., and Morse, H.C., 3rd (2008a). An ENU-induced mutation in the lymphotoxin alpha gene impairs organogenesis of lymphoid tissues in C57BL/6 mice. *Biochem Biophys Res Commun* 370, 461-467.

Wang, W., Lin, C., Lu, D., Ning, Z., Cox, T., Melvin, D., Wang, X., Bradley, A., and Liu, P. (2008b). Chromosomal transposition of PiggyBac in mouse embryonic stem cells. *Proc Natl Acad Sci U S A* 105, 9290-9295.

Ware, C.F., Crowe, P.D., Grayson, M.H., Androlewicz, M.J., and Browning, J.L. (1992). Expression of surface lymphotoxin and tumor necrosis factor on activated T, B, and natural killer cells. *J Immunol* *149*, 3881-3888.

Washburn, T., Schweighoffer, E., Gridley, T., Chang, D., Fowlkes, B.J., Cado, D., and Robey, E. (1997). Notch activity influences the alphabeta versus gammadelta T cell lineage decision. *Cell* *88*, 833-843.

Waterston, R.H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J.F., Agarwal, P., Agarwala, R., Ainscough, R., Alexandersson, M., An, P., *et al.* (2002). Initial sequencing and comparative analysis of the mouse genome. *Nature* *420*, 520-562.

Wilde, J.I., and Watson, S.P. (2001). Regulation of phospholipase C gamma isoforms in haematopoietic cells: why one, not the other? *Cell Signal* *13*, 691-701.

Wildin, R.S., Wang, H.U., Forbush, K.A., and Perlmutter, R.M. (1995). Functional dissection of the murine lck distal promoter. *J Immunol* *155*, 1286-1295.

Winnard, P.T., Jr., Kluth, J.B., and Raman, V. (2006). Noninvasive optical tracking of red fluorescent protein-expressing cancer cells in a model of metastatic breast cancer. *Neoplasia* *8*, 796-806.

Wolfer, A., Wilson, A., Nemir, M., MacDonald, H.R., and Radtke, F. (2002). Inactivation of Notch1 impairs VDJbeta rearrangement and allows pre-TCR-independent survival of early alpha beta Lineage Thymocytes. *Immunity* *16*, 869-879.

Yee, C., Thompson, J.A., Byrd, D., Riddell, S.R., Roche, P., Celis, E., and Greenberg, P.D. (2002). Adoptive T cell therapy using antigen-specific CD8+ T cell clones for the treatment of patients with metastatic melanoma: in vivo persistence, migration, and antitumor effect of transferred T cells. *Proc Natl Acad Sci U S A* *99*, 16168-16173.

Yokota, Y., Mansouri, A., Mori, S., Sugawara, S., Adachi, S., Nishikawa, S., and Gruss, P. (1999). Development of peripheral lymphoid organs and natural killer cells depends on the helix-loop-helix inhibitor Id2. *Nature* 397, 702-706.

Yoshida, T., Ng, S.Y., Zuniga-Pflucker, J.C., and Georgopoulos, K. (2006). Early hematopoietic lineage restrictions directed by Ikaros. *Nat Immunol* 7, 382-391.

Young, D.C., Kingsley, S.D., Ryan, K.A., and Dutko, F.J. (1993). Selective inactivation of eukaryotic beta-galactosidase in assays for inhibitors of HIV-1 TAT using bacterial beta-galactosidase as a reporter enzyme. *Anal Biochem* 215, 24-30.

Yu, J., Vodyanik, M.A., Smuga-Otto, K., Antosiewicz-Bourget, J., Frane, J.L., Tian, S., Nie, J., Jonsdottir, G.A., Ruotti, V., Stewart, R., *et al.* (2007). Induced pluripotent stem cell lines derived from human somatic cells. *Science* 318, 1917-1920.

Yucel, R., Karsunky, H., Klein-Hitpass, L., and Moroy, T. (2003). The transcriptional repressor Gfi1 affects development of early, uncommitted c-Kit<sup>+</sup> T cell progenitors and CD4/CD8 lineage decision in the thymus. *J Exp Med* 197, 831-844.

Yui, M.A., and Rothenberg, E.V. (2004). Deranged early T cell development in immunodeficient strains of nonobese diabetic mice. *J Immunol* 173, 5381-5391.

Zhang, S., Rozell, M., Verma, R.K., Albu, D.I., Califano, D., Vanvalkenburgh, J., Merchant, A., Rangel-Moreno, J., Randall, T.D., Jenkins, N.A., *et al.* (2010). Antigen-specific clonal expansion and cytolytic effector function of CD8<sup>+</sup> T lymphocytes depend on the transcription factor Bcl11b. *J Exp Med* 207, 1687-1699.

Zheng, W., and Flavell, R.A. (1997). The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* 89, 587-596.

Zhuang, Y., Soriano, P., and Weintraub, H. (1994). The helix-loop-helix gene E2A is required for B cell formation. *Cell* 79, 875-884.

Zijlstra, M., Bix, M., Simister, N.E., Loring, J.M., Raulet, D.H., and Jaenisch, R. (1990). Beta 2-microglobulin deficient mice lack CD4-8+ cytolytic T cells. *Nature* 344, 742-746.

Zitvogel, L., Tesniere, A., and Kroemer, G. (2006). Cancer despite immunosurveillance: immunoselection and immunosubversion. *Nat Rev Immunol* 6, 715-727.

Zwirner, N.W., and Domaica, C.I. (2010). Cytokine regulation of natural killer cell effector functions. *Biofactors*.