

BIBLIOGRAPHY

- Abbott, R. J. M., Spendlove, I., Roversi, P., Fitzgibbon, H., Knott, V., Teriete, P., McDonnell, J. M., Handford, P. A., and Lea, S. M. (2007). Structural and functional characterization of a novel T cell receptor co-regulatory protein complex, CD97-CD55. *J. Biol. Chem.*, 282(30):22023–22032.
- Alkhatib, G. (2009). The biology of CCR5 and CXCR4. *Curr. Opin. HIV AIDS*, 4(2):96–103.
- Altman, J. D., Moss, P. A., Goulder, P. J., Barouch, D. H., McHeyzer-Williams, M. G., Bell, J. I., McMichael, A. J., and Davis, M. M. (1996). Phenotypic analysis of antigen-specific T lymphocytes. *Science*, 274(5284):94–96.
- Araç, D., Boucard, A. A., Bolliger, M. F., Nguyen, J., Soltis, S. M., Südhof, T. C., and Brunger, A. T. (2012). A novel evolutionarily conserved domain of cell-adhesion GPCRs mediates autoproteolysis. *EMBO J.*, 31(6):1364–1378.
- Bausch-Fluck, D., Hofmann, A., Bock, T., Frei, A. P., Cerciello, F., Jacobs, A., Moest, H., Omasits, U., Gundry, R. L., Yoon, C., Schiess, R., Schmidt, A., Mirkowska, P., Härtlová, A., Van Eyk, J. E., Bourquin, J.-P., Aebersold, R., Boheler, K. R., Zandstra, P., and Wollscheid, B. (2015). A mass Spectrometric-Derived cell surface protein atlas. *PLoS One*, 10(4):e0121314.
- Bjarnadóttir, T. K., Fredriksson, R., Höglund, P. J., Gloriam, D. E., Lagerström, M. C., and Schiöth, H. B. (2004). The human and mouse repertoire of the adhesion family of g-protein-coupled receptors. *Genomics*, 84(1):23–33.
- Brandsma, I. and Gent, D. C. (2012). Pathway choice in DNA double strand break repair: observations of a balancing act. *Genome Integr.*, 3(1):9.
- Brodsky, R. A., Mukhina, G. L., Li, S., Nelson, K. L., Chiurazzi, P. L., Buckley, J. T., and Borowitz, M. J. (2000). Improved detection and characterization of paroxysmal nocturnal hemoglobinuria using fluorescent aerolysin. *Am. J. Clin. Pathol.*, 114(3):459–466.
- Brown, M. H. and Barclay, A. N. (1994). Expression of immunoglobulin and scavenger receptor superfamily domains as chimeric proteins with domains 3 and 4 of CD4 for ligand analysis. *Protein Eng.*, 7(4):515–521.

- Bushell, K. M., Söllner, C., Schuster-Boeckler, B., Bateman, A., and Wright, G. J. (2008). Large-scale screening for novel low-affinity extracellular protein interactions. *Genome Res.*, 18(4):622–630.
- Chakraborty, S., Ji, H., Kabadi, A. M., Gersbach, C. A., Christoforou, N., and Leong, K. W. (2014). A CRISPR/Cas9-based system for reprogramming cell lineage specification. *Stem Cell Reports*, 3(6):940–947.
- Chan, H. M. and La Thangue, N. B. (2001). p300/CBP proteins: HATs for transcriptional bridges and scaffolds. *J. Cell Sci.*, 114(Pt 13):2363–2373.
- Chavez, A., Scheiman, J., Vora, S., Pruitt, B. W., Tuttle, M., P R Iyer, E., Lin, S., Kiani, S., Guzman, C. D., Wiegand, D. J., Ter-Ovanesyan, D., Braff, J. L., Davidsohn, N., Housden, B. E., Perrimon, N., Weiss, R., Aach, J., Collins, J. J., and Church, G. M. (2015). Highly efficient cas9-mediated transcriptional programming. *Nat. Methods*, 12(4):326–328.
- Chavez, A., Tuttle, M., Pruitt, B. W., Ewen-Campen, B., Chari, R., Ter-Ovanesyan, D., Haque, S. J., Cecchi, R. J., Kowal, E. J. K., Buchthal, J., Housden, B. E., Perrimon, N., Collins, J. J., and Church, G. (2016). Comparison of cas9 activators in multiple species. *Nat. Methods*, 13(7):563–567.
- Chen, B., Gilbert, L. A., Cimini, B. A., Schnitzbauer, J., Zhang, W., Li, G.-W., Park, J., Blackburn, E. H., Weissman, J. S., Qi, L. S., and Huang, B. (2013). Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system. *Cell*, 155(7):1479–1491.
- Cheng, A. W., Wang, H., Yang, H., Shi, L., Katz, Y., Theunissen, T. W., Rangarajan, S., Shivalila, C. S., Dadon, D. B., and Jaenisch, R. (2013). Multiplexed activation of endogenous genes by CRISPR-on, an RNA-guided transcriptional activator system. *Cell Res.*, 23:1163.
- Collins, A. V., Brodie, D. W., Gilbert, R. J. C., Iaboni, A., Manso-Sancho, R., Walse, B., Stuart, D. I., van der Merwe, P. A., and Davis, S. J. (2002). The interaction properties of costimulatory molecules revisited. *Immunity*, 17(2):201–210.
- Contardi, E., Palmisano, G. L., Tazzari, P. L., Martelli, A. M., Falà, F., Fabbì, M., Kato, T., Lucarelli, E., Donati, D., Polito, L., Bolognesi, A., Ricci, F., Salvi, S., Gargaglione, V., Mantero, S., Alberghini, M., Ferrara, G. B., and Pistillo, M. P. (2005). CTLA-4 is constitutively expressed on tumor cells and can trigger apoptosis upon ligand interaction. *Int. J. Cancer*, 117(4):538–550.
- Crosnier, C., Bustamante, L. Y., Bartholdson, S. J., Bei, A. K., Theron, M., Uchikawa, M., Mboup, S., Ndir, O., Kwiatkowski, D. P., Duraisingh, M. T., Rayner, J. C., and Wright, G. J. (2011). Basigin is a receptor essential for erythrocyte invasion by plasmodium falciparum. *Nature*, 480(7378):534–537.
- Czajkowsky, D. M., Hu, J., Shao, Z., and Pleass, R. J. (2012). Fc-fusion proteins: new developments and future perspectives. *EMBO Mol. Med.*, 4(10):1015–1028.

- da Cunha, J. P. C., Galante, P. A. F., de Souza, J. E., de Souza, R. F., Carvalho, P. M., Ohara, D. T., Moura, R. P., Oba-Shinja, S. M., Marie, S. K. N., Silva, Jr, W. A., Perez, R. O., Stransky, B., Pieprzyk, M., Moore, J., Caballero, O., Gama-Rodrigues, J., Habr-Gama, A., Kuo, W. P., Simpson, A. J., Camargo, A. A., Old, L. J., and de Souza, S. J. (2009). Bioinformatics construction of the human cell surfaceome. *Proc. Natl. Acad. Sci. U. S. A.*, 106(39):16752–16757.
- Dancy, B. M. and Cole, P. A. (2015). Protein lysine acetylation by p300/CBP. *Chem. Rev.*, 115(6):2419–2452.
- Dayalan Naidu, S. and Dinkova-Kostova, A. T. (2017). Regulation of the mammalian heat shock factor 1. *FEBS J.*, 284(11):1606–1627.
- Duman, J. G., Tzeng, C. P., Tu, Y.-K., Munjal, T., Schwechter, B., Ho, T. S.-Y., and Tolia, K. F. (2013). The adhesion-GPCR BAI1 regulates synaptogenesis by controlling the recruitment of the Par3/Tiam1 polarity complex to synaptic sites. *J. Neurosci.*, 33(16):6964–6978.
- Durinck, S., Spellman, P. T., Birney, E., and Huber, W. (2009). Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomart. *Nat. Protoc.*, 4(8):1184–1191.
- Einhauer, A. and Jungbauer, A. (2001). The FLAG™ peptide, a versatile fusion tag for the purification of recombinant proteins. *J. Biochem. Biophys. Methods*, 49(1):455–465.
- Ellis, E. L. and Delbrück, M. (1939). THE GROWTH OF BACTERIOPHAGE. *J. Gen. Physiol.*, 22(3):365–384.
- Feige, M. J. and Hendershot, L. M. (2011). Disulfide bonds in ER protein folding and homeostasis. *Curr. Opin. Cell Biol.*, 23(2):167–175.
- Ferguson, M. A. J., Hart, G. W., and Kinoshita, T. (2017). Glycosylphosphatidylinositol anchors. In Varki, A., Cummings, R. D., Esko, J. D., Stanley, P., Hart, G. W., Aebi, M., Darvill, A. G., Kinoshita, T., Packer, N. H., Prestegard, J. H., Schnaar, R. L., and Seeberger, P. H., editors, *Essentials of Glycobiology*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor (NY).
- Fields, S. and Song, O. (1989). A novel genetic system to detect protein-protein interactions. *Nature*, 340(6230):245–246.
- Fournier, A. E., GrandPre, T., and Strittmatter, S. M. (2001). Identification of a receptor mediating nogo-66 inhibition of axonal regeneration. *Nature*, 409(6818):341–346.
- Frei, A. P., Jeon, O.-Y., Kilcher, S., Moest, H., Henning, L. M., Jost, C., Plückthun, A., Mercer, J., Aebersold, R., Carreira, E. M., and Wollscheid, B. (2012). Direct identification of ligand-receptor interactions on living cells and tissues. *Nat. Biotechnol.*, 30(10):997–1001.
- Futschik, M. E., Chaurasia, G., and Herzel, H. (2007). Comparison of human protein-protein interaction maps. *Bioinformatics*, 23(5):605–611.

- Gilbert, L. A., Horlbeck, M. A., Adamson, B., Villalta, J. E., Chen, Y., Whitehead, E. H., Guimaraes, C., Panning, B., Ploegh, H. L., Bassik, M. C., Qi, L. S., Kampmann, M., and Weissman, J. S. (2014). Genome-Scale CRISPR-Mediated control of gene repression and activation. *Cell*, 159(3):647–661.
- Gilbert, L. A., Larson, M. H., Morsut, L., Liu, Z., Brar, G. A., Torres, S. E., Stern-Ginossar, N., Brandman, O., Whitehead, E. H., Doudna, J. A., Lim, W. A., Weissman, J. S., and Qi, L. S. (2013). CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes. *Cell*, 154(2):442–451.
- Haeussler, M., Schönig, K., Eckert, H., Eschstruth, A., Mianné, J., Renaud, J.-B., Schneider-Maunoury, S., Shkumatava, A., Teboul, L., Kent, J., Joly, J.-S., and Concordet, J.-P. (2016). Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol.*, 17(1):148.
- Hardwick, J. M., Tse, L., Applegren, N., Nicholas, J., and Veluona, M. A. (1992). The Epstein-Barr virus R transactivator (rta) contains a complex, potent activation domain with properties different from those of VP16. *J. Virol.*, 66(9):5500–5508.
- Harrow, J., Frankish, A., Gonzalez, J. M., Tapanari, E., Diekhans, M., Kokocinski, F., Aken, B. L., Barrell, D., Zadissa, A., Searle, S., Barnes, I., Bignell, A., Boychenko, V., Hunt, T., Kay, M., Mukherjee, G., Rajan, J., Despacio-Reyes, G., Saunders, G., Steward, C., Harte, R., Lin, M., Howald, C., Tanzer, A., Derrien, T., Chrast, J., Walters, N., Balasubramanian, S., Pei, B., Tress, M., Rodriguez, J. M., Ezkurdia, I., van Baren, J., Brent, M., Haussler, D., Kellis, M., Valencia, A., Reymond, A., Gerstein, M., Guigó, R., and Hubbard, T. J. (2012). GENCODE: the reference human genome annotation for the ENCODE project. *Genome Res.*, 22(9):1760–1774.
- Hauser, A. S., Chavali, S., Masuho, I., Jahn, L. J., Martemyanov, K. A., Gloriam, D. E., and Babu, M. M. (2018). Pharmacogenomics of GPCR drug targets. *Cell*, 172(1-2):41–54.e19.
- Heaton, B. E., Kennedy, E. M., Dumm, R. E., Harding, A. T., Sacco, M. T., Sachs, D., and Heaton, N. S. (2017). A CRISPR activation screen identifies a pan-avian influenza virus inhibitory host factor. *Cell Rep.*, 20(7):1503–1512.
- Hellebrekers, D. M. E. I., Castermans, K., Viré, E., Dings, R. P. M., Hoebbers, N. T. H., Mayo, K. H., Oude Egbrink, M. G. A., Molema, G., Fuks, F., van Engeland, M., and Griffioen, A. W. (2006). Epigenetic regulation of tumor endothelial cell anergy: silencing of intercellular adhesion molecule-1 by histone modifications. *Cancer Res.*, 66(22):10770–10777.
- Hilton, I. B., D’Ippolito, A. M., Vockley, C. M., Thakore, P. I., Crawford, G. E., Reddy, T. E., and Gersbach, C. A. (2015). Epigenome editing by a CRISPR-Cas9-based acetyltransferase activates genes from promoters and enhancers. *Nat. Biotechnol.*, 33(5):510–517.
- Hochuli, E., Bannwarth, W., Döbeli, H., Gentz, R., and Stüber, D. (1988). Genetic approach to facilitate purification of recombinant proteins with a novel metal chelate adsorbent. *Biotechnology*, 6:1321.

- Horlbeck, M. A., Gilbert, L. A., Villalta, J. E., Adamson, B., Pak, R. A., Chen, Y., Fields, A. P., Park, C. Y., Corn, J. E., Kampmann, M., and Weissman, J. S. (2016). Compact and highly active next-generation libraries for CRISPR-mediated gene repression and activation. *Elife*, 5.
- Huttlin, E. L., Ting, L., Bruckner, R. J., Gebreab, F., Gygi, M. P., Szpyt, J., Tam, S., Zarraga, G., Colby, G., Baltier, K., Dong, R., Guarani, V., Vaites, L. P., Ordureau, A., Rad, R., Erickson, B. K., Wühr, M., Chick, J., Zhai, B., Kolippakkam, D., Mintseris, J., Obar, R. A., Harris, T., Artavanis-Tsakonas, S., Sowa, M. E., De Camilli, P., Paulo, J. A., Harper, J. W., and Gygi, S. P. (2015). The BioPlex network: A systematic exploration of the human interactome. *Cell*, 162(2):425–440.
- Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., and Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*, 337(6096):816–821.
- Jinno, A. and Park, P. W. (2015). Role of glycosaminoglycans in infectious disease. *Methods Mol. Biol.*, 1229:567–585.
- Josefin Bartholdson, S., Bustamante, L. Y., Crosnier, C., Johnson, S., Lea, S., Rayner, J. C., and Wright, G. J. (2012). Semaphorin-7A is an erythrocyte receptor for p. falciparum Merozoite-Specific TRAP homolog, MTRAP. *PLoS Pathog.*, 8(11):e1003031.
- Kent, W. J. (2002). BLAT—The BLAST-Like alignment tool. *Genome Res.*, 12(4):656–664.
- Konermann, S., Brigham, M. D., Trevino, A. E., Joung, J., Abudayyeh, O. O., Barcena, C., Hsu, P. D., Habib, N., Gootenberg, J. S., Nishimasu, H., Nureki, O., and Zhang, F. (2015). Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature*, 517(7536):583–588.
- Leeuwenberg, J. F., Smeets, E. F., Neefjes, J. J., Shaffer, M. A., Cinek, T., Jeunhomme, T. M., Ahern, T. J., and Buurman, W. A. (1992). E-selectin and intercellular adhesion molecule-1 are released by activated human endothelial cells in vitro. *Immunology*, 77(4):543–549.
- Lema Tomé, C. M., Palma, E., Ferluga, S., Lowther, W. T., Hantgan, R., Wykosky, J., and Debinski, W. (2012). Structural and functional characterization of monomeric EphrinA1 binding site to EphA2 receptor. *J. Biol. Chem.*, 287(17):14012–14022.
- Li, W., Xu, H., Xiao, T., Cong, L., Love, M. I., Zhang, F., Irizarry, R. A., Liu, J. S., Brown, M., and Liu, X. S. (2014). MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9 knockout screens. *Genome Biol.*, 15(12):554.
- Lin, H. H., Stacey, M., Saxby, C., Knott, V., Chaudhry, Y., Evans, D., Gordon, S., McKnight, A. J., Handford, P., and Lea, S. (2001). Molecular analysis of the epidermal growth factor-like short consensus repeat domain-mediated protein-protein interactions: dissection of the CD97-CD55 complex. *J. Biol. Chem.*, 276(26):24160–24169.
- Liu, B. P., Fournier, A., GrandPré, T., and Strittmatter, S. M. (2002). Myelin-associated glycoprotein as a functional ligand for the nogo-66 receptor. *Science*, 297(5584):1190–1193.

- Liu, H., Wei, Z., Dominguez, A., Li, Y., Wang, X., and Qi, L. S. (2015). CRISPR-ERA: a comprehensive design tool for CRISPR-mediated gene editing, repression and activation. *Bioinformatics*, 31(22):3676–3678.
- Livak, K. J. and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2(-delta delta C(T)) method. *Methods*, 25(4):402–408.
- Loignon, M., Perret, S., Kelly, J., Boulais, D., Cass, B., Bisson, L., Afkhamizarreh, F., and Durocher, Y. (2008). Stable high volumetric production of glycosylated human recombinant IFNalpha2b in HEK293 cells. *BMC Biotechnol.*, 8:65.
- Lopalco, L. (2010). CCR5: From natural resistance to a new Anti-HIV strategy. *Viruses*, 2(2):574–600.
- Luo, R., Jeong, S.-J., Jin, Z., Strokes, N., Li, S., and Piao, X. (2011). G protein-coupled receptor 56 and collagen III, a receptor-ligand pair, regulates cortical development and lamination. *Proc. Natl. Acad. Sci. U. S. A.*, 108(31):12925–12930.
- Ma, H., Wu, Y., Dang, Y., Choi, J.-G., Zhang, J., and Wu, H. (2014). Pol III promoters to express small RNAs: Delineation of transcription initiation. *Mol. Ther. Nucleic Acids*, 3:e161.
- Maeder, M. L., Linder, S. J., Cascio, V. M., Fu, Y., Ho, Q. H., and Joung, J. K. (2013). CRISPR RNA-guided activation of endogenous human genes. *Nat. Methods*, 10(10):977–979.
- Makarova, K. S., Wolf, Y. I., Alkhnbashi, O. S., Costa, F., Shah, S. A., Saunders, S. J., Barrangou, R., Brouns, S. J. J., Charpentier, E., Haft, D. H., Horvath, P., Moineau, S., Mojica, F. J. M., Terns, R. M., Terns, M. P., White, M. F., Yakunin, A. F., Garrett, R. A., van der Oost, J., Backofen, R., and Koonin, E. V. (2015). An updated evolutionary classification of CRISPR-Cas systems. *Nat. Rev. Microbiol.*, 13(11):722–736.
- Martin, S., Söllner, C., Charoensawan, V., Adryan, B., Thisse, B., Thisse, C., Teichmann, S., and Wright, G. J. (2010). Construction of a large extracellular protein interaction network and its resolution by spatiotemporal expression profiling. *Mol. Cell. Proteomics*, 9(12):2654–2665.
- Michel, M. C., Wieland, T., and Tsujimoto, G. (2009). How reliable are g-protein-coupled receptor antibodies? *Naunyn. Schmiedeberg's Arch. Pharmacol.*, 379(4):385–388.
- Mojica, F. J. M., Díez-Villaseñor, C., García-Martínez, J., and Soria, E. (2005). Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic elements. *J. Mol. Evol.*, 60(2):174–182.
- Mullican, S. E., Lin-Schmidt, X., Chin, C.-N., Chavez, J. A., Furman, J. L., Armstrong, A. A., Beck, S. C., South, V. J., Dinh, T. Q., Cash-Mason, T. D., Cavanaugh, C. R., Nelson, S., Huang, C., Hunter, M. J., and Rangwala, S. M. (2017). GFRAL is the receptor for GDF15 and the ligand promotes weight loss in mice and nonhuman primates. *Nat. Med.*, 23(10):1150–1157.

- Nam, J.-W., Rissland, O. S., Koppstein, D., Abreu-Goodger, C., Jan, C. H., Agarwal, V., Yildirim, M. A., Rodriguez, A., and Bartel, D. P. (2014). Global analyses of the effect of different cellular contexts on microRNA targeting. *Mol. Cell*, 53(6):1031–1043.
- Nielsen, S., Yuzenkova, Y., and Zenkin, N. (2013). Mechanism of eukaryotic RNA polymerase III transcription termination. *Science*, 340(6140):1577–1580.
- Ong, S. H., Li, Y., Koike-Yusa, H., and Yusa, K. (2017). Optimised metrics for CRISPR-KO screens with second-generation gRNA libraries. *Sci. Rep.*, 7(1):7384.
- Ord, R. L., Rodriguez, M., and Lobo, C. A. (2015). Malaria invasion ligand RH5 and its prime candidacy in blood-stage malaria vaccine design. *Hum. Vaccin. Immunother.*, 11(6):1465–1473.
- O’Sullivan, M. L., de Wit, J., Savas, J. N., Comoletti, D., Otto-Hitt, S., Yates, 3rd, J. R., and Ghosh, A. (2012). FLRT proteins are endogenous latrophilin ligands and regulate excitatory synapse development. *Neuron*, 73(5):903–910.
- Özkan, E., Carrillo, R. A., Eastman, C. L., Weiszmann, R., Waghray, D., Johnson, K. G., Zinn, K., Celniker, S. E., and Garcia, K. C. (2013). An extracellular interactome of immunoglobulin and LRR proteins reveals receptor-ligand networks. *Cell*, 154(1):228–239.
- Paavola, K. J., Sidik, H., Zuchero, J. B., Eckart, M., and Talbot, W. S. (2014). Type IV collagen is an activating ligand for the adhesion G protein-coupled receptor GPR126. *Sci. Signal.*, 7(338):ra76.
- Park, D., Tosello-Tramont, A.-C., Elliott, M. R., Lu, M., Haney, L. B., Ma, Z., Klibanov, A. L., Mandell, J. W., and Ravichandran, K. S. (2007). BAI1 is an engulfment receptor for apoptotic cells upstream of the ELMO/Dock180/Rac module. *Nature*, 450(7168):430–434.
- Paulick, M. G. and Bertozzi, C. R. (2008). The glycosylphosphatidylinositol anchor: a complex membrane-anchoring structure for proteins. *Biochemistry*, 47(27):6991–7000.
- Petersen, S. C., Luo, R., Liebscher, I., Giera, S., Jeong, S.-J., Mogha, A., Ghidinelli, M., Feltri, M. L., Schöneberg, T., Piao, X., and Monk, K. R. (2015). The adhesion GPCR GPR126 has distinct, domain-dependent functions in schwann cell development mediated by interaction with laminin-211. *Neuron*, 85(4):755–769.
- Petersen, T. N., Brunak, S., von Heijne, G., and Nielsen, H. (2011). SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat. Methods*, 8(10):785–786.
- Pierleoni, A., Martelli, P. L., and Casadio, R. (2008). PredGPI: a GPI-anchor predictor. *BMC Bioinformatics*, 9:392.
- Pizarro-Cerdá, J. and Cossart, P. (2006). Bacterial adhesion and entry into host cells. *Cell*, 124(4):715–727.
- Quail, M. A., Otto, T. D., Gu, Y., Harris, S. R., Skelly, T. F., McQuillan, J. A., Swerdlow, H. P., and Oyola, S. O. (2011). Optimal enzymes for amplifying sequencing libraries. *Nat. Methods*, 9(1):10–11.

- Robak, L. A., Venkatesh, K., Lee, H., Raiker, S. J., Duan, Y., Lee-Osbourne, J., Hofer, T., Mage, R. G., Rader, C., and Giger, R. J. (2009). Molecular basis of the interactions of the nogo-66 receptor and its homolog NgR2 with myelin-associated glycoprotein: development of NgROMNI-Fc, a novel antagonist of CNS myelin inhibition. *J. Neurosci.*, 29(18):5768–5783.
- Rodriguez, J. M., Maietta, P., Ezkurdia, I., Pietrelli, A., Wesselink, J.-J., Lopez, G., Valencia, A., and Tress, M. L. (2013). APPRIS: annotation of principal and alternative splice isoforms. *Nucleic Acids Res.*, 41(Database issue):D110–7.
- Rolland, T., Taşan, M., Charlotteaux, B., Pevzner, S. J., Zhong, Q., Sahni, N., Yi, S., Lemmens, I., Fontanillo, C., Mosca, R., Kamburov, A., Ghiassian, S. D., Yang, X., Ghamsari, L., Balcha, D., Begg, B. E., Braun, P., Brehme, M., Broly, M. P., Carvunis, A.-R., Convery-Zupan, D., Corominas, R., Coulombe-Huntington, J., Dann, E., Dreze, M., Dricot, A., Fan, C., Franzosa, E., Gebreab, F., Gutierrez, B. J., Hardy, M. F., Jin, M., Kang, S., Kiros, R., Lin, G. N., Luck, K., MacWilliams, A., Menche, J., Murray, R. R., Palagi, A., Poulin, M. M., Rambout, X., Rasla, J., Reichert, P., Romero, V., Ruysinck, E., Sahalie, J. M., Scholz, A., Shah, A. A., Sharma, A., Shen, Y., Spirohn, K., Tam, S., Tejada, A. O., Trigg, S. A., Twizere, J.-C., Vega, K., Walsh, J., Cusick, M. E., Xia, Y., Barabási, A.-L., Iakoucheva, L. M., Aloy, P., De Las Rivas, J., Tavernier, J., Calderwood, M. A., Hill, D. E., Hao, T., Roth, F. P., and Vidal, M. (2014). A proteome-scale map of the human interactome network. *Cell*, 159(5):1212–1226.
- Safaei, M., Clark, A. J., Ivan, M. E., Oh, M. C., Bloch, O., Sun, M. Z., Oh, T., and Parsa, A. T. (2013). CD97 is a multifunctional leukocyte receptor with distinct roles in human cancers (review). *Int. J. Oncol.*, 43(5):1343–1350.
- Sambrook, J. and Russell, D. W. (2006). Isolation of DNA fragments from polyacrylamide gels by the crush and soak method. *CSH Protoc.*, 2006(1).
- Sansom, D. M. (2000). CD28, CTLA-4 and their ligands: who does what and to whom? *Immunology*, 101(2):169–177.
- Sanson, K. R., Hanna, R. E., Hegde, M., Donovan, K. F., Strand, C., Sullender, M. E., Vaimberg, E. W., Goodale, A., Root, D. E., Piccioni, F., and Doench, J. G. (2018). Up, down, and out: optimized libraries for CRISPRa, CRISPRi, and CRISPR-knockout genetic screens.
- Sapranaukas, R., Gasiunas, G., Fremaux, C., Barrangou, R., Horvath, P., and Siksnys, V. (2011). The streptococcus thermophilus CRISPR/Cas system provides immunity in escherichia coli. *Nucleic Acids Res.*, 39(21):9275–9282.
- Selvaraj, P., Plunkett, M. L., Dustin, M., Sanders, M. E., Shaw, S., and Springer, T. A. (1987). The T lymphocyte glycoprotein CD2 binds the cell surface ligand LFA-3. *Nature*, 326(6111):400–403.
- Seyfried, T. N. and Huysentruyt, L. C. (2013). On the origin of cancer metastasis. *Crit. Rev. Oncog.*, 18(1-2):43–73.
- Sharma, S., Bartholdson, S. J., Couch, A. C., Yusa, K., and Wright, G. J. (2018). Genome-scale identification of cellular pathways required for cell surface recognition. *Genome Res.*

- Silva, J.-P., Lelianova, V. G., Ermolyuk, Y. S., Vysokov, N., Hitchen, P. G., Berninghausen, O., Rahman, M. A., Zangrandi, A., Fidalgo, S., Tonevitsky, A. G., Dell, A., Volynski, K. E., and Ushkaryov, Y. A. (2011). Latrophilin 1 and its endogenous ligand lasso/teneurin-2 form a high-affinity transsynaptic receptor pair with signaling capabilities. *Proc. Natl. Acad. Sci. U. S. A.*, 108(29):12113–12118.
- Simonsen, H. and Lodish, H. F. (1994). Cloning by function: expression cloning in mammalian cells. *Trends Pharmacol. Sci.*, 15(12):437–441.
- Snider, J., Kittanakom, S., Curak, J., and Stagljar, I. (2010). Split-ubiquitin based membrane yeast two-hybrid (MYTH) system: a powerful tool for identifying protein-protein interactions. *J. Vis. Exp.*, (36).
- Snider, J. and Stagljar, I. (2016). Membrane yeast Two-Hybrid (MYTH) mapping of Full-Length membrane protein interactions. *Cold Spring Harb. Protoc.*, 2016(1):db.top077560.
- Sorek, R., Kunin, V., and Hugenholtz, P. (2008). CRISPR—a widespread system that provides acquired resistance against phages in bacteria and archaea. *Nat. Rev. Microbiol.*, 6(3):181–186.
- Stephenson, J. R., Paavola, K. J., Schaefer, S. A., Kaur, B., Van Meir, E. G., and Hall, R. A. (2013). Brain-specific angiogenesis inhibitor-1 signaling, regulation, and enrichment in the postsynaptic density. *J. Biol. Chem.*, 288(31):22248–22256.
- Sternberg, S. H., Redding, S., Jinek, M., Greene, E. C., and Doudna, J. A. (2014). DNA interrogation by the CRISPR RNA-guided endonuclease cas9. *Nature*, 507(7490):62–67.
- Stoveken, H. M., Hajduczuk, A. G., Xu, L., and Tall, G. G. (2015). Adhesion G protein-coupled receptors are activated by exposure of a cryptic tethered agonist. *Proc. Natl. Acad. Sci. U. S. A.*, 112(19):6194–6199.
- Stupack, D. G. and Cheresch, D. A. (2002). Get a ligand, get a life: integrins, signaling and cell survival. *J. Cell Sci.*, 115(Pt 19):3729–3738.
- Sun, Y., Gallagher-Jones, M., Barker, C., and Wright, G. J. (2012). A benchmarked protein microarray-based platform for the identification of novel low-affinity extracellular protein interactions. *Anal. Biochem.*, 424(1):45–53.
- Tanenbaum, M. E., Gilbert, L. A., Qi, L. S., Weissman, J. S., and Vale, R. D. (2014). A protein-tagging system for signal amplification in gene expression and fluorescence imaging. *Cell*, 159(3):635–646.
- Tanner, W. and Lehle, L. (1987). Protein glycosylation in yeast. *Biochim. Biophys. Acta*, 906(1):81–99.
- Thul, P. J. and Lindskog, C. (2018). The human protein atlas: A spatial map of the human proteome. *Protein Sci.*, 27(1):233–244.
- Tumbar, T., Sudlow, G., and Belmont, A. S. (1999). Large-scale chromatin unfolding and remodeling induced by VP16 acidic activation domain. *J. Cell Biol.*, 145(7):1341–1354.

- Turner, L., Lavstsen, T., Berger, S. S., Wang, C. W., Petersen, J. E. V., Avril, M., Brazier, A. J., Freeth, J., Jespersen, J. S., Nielsen, M. A., Magistrado, P., Lusingu, J., Smith, J. D., Higgins, M. K., and Theander, T. G. (2013). Severe malaria is associated with parasite binding to endothelial protein C receptor. *Nature*, 498(7455):502–505.
- Uhlén, M., Fagerberg, L., Hallström, B. M., Lindskog, C., Oksvold, P., Mardinoglu, A., Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., Olsson, I., Edlund, K., Lundberg, E., Navani, S., Szigartyo, C. A.-K., Odeberg, J., Djureinovic, D., Takanen, J. O., Hober, S., Alm, T., Edqvist, P.-H., Berling, H., Tegel, H., Mulder, J., Rockberg, J., Nilsson, P., Schwenk, J. M., Hamsten, M., von Feilitzen, K., Forsberg, M., Persson, L., Johansson, F., Zwahlen, M., von Heijne, G., Nielsen, J., and Pontén, F. (2015). Proteomics. tissue-based map of the human proteome. *Science*, 347(6220):1260419.
- Ulloa-Aguirre, A., Timossi, C., Damián-Matsumura, P., and Dias, J. A. (1999). Role of glycosylation in function of follicle-stimulating hormone. *Endocrine*, 11(3):205–215.
- Vallon, M. and Essler, M. (2006). Proteolytically processed soluble tumor endothelial marker (TEM) 5 mediates endothelial cell survival during angiogenesis by linking integrin alpha(v)beta3 to glycosaminoglycans. *J. Biol. Chem.*, 281(45):34179–34188.
- van der Merwe, P. A. and Barclay, A. N. (1994). Transient intercellular adhesion: the importance of weak protein-protein interactions. *Trends Biochem. Sci.*, 19(9):354–358.
- van der Merwe, P. A., Bodian, D. L., Daenke, S., Linsley, P., and Davis, S. J. (1997). CD80 (b7-1) binds both CD28 and CTLA-4 with a low affinity and very fast kinetics. *J. Exp. Med.*, 185(3):393–403.
- van Essen, D., Engist, B., Natoli, G., and Sacconi, S. (2009). Two modes of transcriptional activation at native promoters by NF-kappaB p65. *PLoS Biol.*, 7(3):e73.
- Wang, T., Birsoy, K., Hughes, N. W., Krupczak, K. M., Post, Y., Wei, J. J., Lander, E. S., and Sabatini, D. M. (2015). Identification and characterization of essential genes in the human genome. *Science*, 350(6264):1096–1101.
- Wills, Z. P., Mandel-Brehm, C., Mardinly, A. R., McCord, A. E., Giger, R. J., and Greenberg, M. E. (2012). The nogo receptor family restricts synapse number in the developing hippocampus. *Neuron*, 73(3):466–481.
- Wojtowicz, W. M., Wu, W., Andre, I., Qian, B., Baker, D., and Zipursky, S. L. (2007). A vast repertoire of dscam binding specificities arises from modular interactions of variable ig domains. *Cell*, 130(6):1134–1145.
- Wright, G. J. (2009). Signal initiation in biological systems: the properties and detection of transient extracellular protein interactions. *Mol. Biosyst.*, 5(12):1405–1412.
- Wright, G. J., Cherwinski, H., Foster-Cuevas, M., Brooke, G., Puklavec, M. J., Bigler, M., Song, Y., Jenmalm, M., Gorman, D., McClanahan, T., Liu, M.-R., Brown, M. H., Sedgwick, J. D., Phillips, J. H., and Barclay, A. N. (2003). Characterization of the CD200 receptor family in mice and humans and their interactions with CD200. *J. Immunol.*, 171(6):3034–3046.

- Wright, G. J., Puklavec, M. J., Willis, A. C., Hoek, R. M., Sedgwick, J. D., Brown, M. H., and Barclay, A. N. (2000). Lymphoid/neuronal cell surface OX2 glycoprotein recognizes a novel receptor on macrophages implicated in the control of their function. *Immunity*, 13(2):233–242.
- Wuethrich, I., Peeters, J. G. C., Blom, A. E. M., Theile, C. S., Li, Z., Spooner, E., Ploegh, H. L., and Guimaraes, C. P. (2014). Site-specific chemoenzymatic labeling of aerolysin enables the identification of new aerolysin receptors. *PLoS One*, 9(10):e109883.
- Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., and Madden, T. L. (2012). Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*, 13:134.
- Young, M. T., Beckmann, R., Toye, A. M., and Tanner, M. J. (2000). Red-cell glycophorin a-band 3 interactions associated with the movement of band 3 to the cell surface. *Biochem. J*, 350 Pt 1:53–60.
- Yusa, K., Zhou, L., Li, M. A., Bradley, A., and Craig, N. L. (2011). A hyperactive piggybac transposase for mammalian applications. *Proc. Natl. Acad. Sci. U. S. A.*, 108(4):1531–1536.
- Zhang, N. and Li, L. (2004). Effects of common surfactants on protein digestion and matrix-assisted laser desorption/ionization mass spectrometric analysis of the digested peptides using two-layer sample preparation. *Rapid Commun. Mass Spectrom.*, 18(8):889–896.
- Zhu, D., Li, C., Swanson, A. M., Villalba, R. M., Guo, J., Zhang, Z., Matheny, S., Murakami, T., Stephenson, J. R., Daniel, S., Fukata, M., Hall, R. A., Olson, J. J., Neigh, G. N., Smith, Y., Rainnie, D. G., and Meir, E. G. V. (2015). BAI1 regulates spatial learning and synaptic plasticity in the hippocampus. *J. Clin. Invest.*, 125(4):1497–1508.
- Ziauddin, J. and Sabatini, D. M. (2001). Microarrays of cells expressing defined cDNAs. *Nature*, 411(6833):107–110.

