

Bibliography

Adams, M. D., Celniker, S. E., Holt, R. A., Evans, C. A., Gocayne, J. D., Amanatides, P. G., Scherer, S. E., Li, P. W., Hoskins, R. A., Galle, R. F., George, R. A., Lewis, S. E., Richards, S., Ashburner, M., Henderson, S. N., Sutton, G. G., Wortman, J. R., Yandell, M. D., Zhang, Q., Chen, L. X., Brandon, R. C., Rogers, Y. H., Blazej, R. G., Champe, M., Pfeiffer, B. D., Wan, K. H., Doyle, C., Baxter, E. G., Helt, G., Nelson, C. R., Gabor, G. L., Abril, J. F., Agbayani, A., An, H. J., Andrews-Pfannkoch, C., Baldwin, D., Ballew, R. M., Basu, A., Baxendale, J., Bayraktaroglu, L., Beasley, E. M., Beeson, K. Y., Benos, P. V., Berman, B. P., Bhandari, D., Bolshakov, S., Borkova, D., Botchan, M. R., Bouck, J., Brokstein, P., Brottier, P., Burtis, K. C., Busam, D. A., Butler, H., Cadieu, E., Center, A., Chandra, I., Cherry, J. M., Cawley, S., Dahlke, C., Davenport, L. B., Davies, P., de Pablos, B., Delcher, A., Deng, Z., Mays, A. D., Dew, I., Dietz, S. M., Dodson, K., Doup, L. E., Downes, M., Dugan-Rocha, S., Dunkov, Dunkov, B. C., Dunn, P., Durbin, K. J., Evangelista, C. C., Ferraz, C., Ferriera, S., Fleischmann, W., Fosler, C., Gabrielian, A. E., Garg, N. S., Gelbart, W. M., Glasser, K., Glodek, A., Gong, F., Gorrell, J. H., Gu, Z., Guan, P., Harris, M., Harris, N. L., Harvey, D., Heiman, T. J., Hernandez, J. R., Houck, J., Hostin, D., Houston, K. A., Howland, T. J., Wei, M. H., Ibegwam, C., Jalali, M., Kalush, F., Karpen, G. H., Ke, Z., Kennison, J. A., Ketchum, K. A., Kimmel, B. E., Kodira, C. D., Kraft, C., Kravitz, S., Kulp, D., Lai, Z., Lasko, P., Lei, Y., Levitsky, A. A., Li, J., Li, Z., Liang, Y., Lin, X., Liu, X., Mattei, B., McIntosh, T. C., McLeod, M. P., McPherson, D., Merkulov, G., Milshina, N. V., Mobarry, C., Morris, J., Moshrefi, A., Mount, S. M., Moy, M., Murphy, B., Murphy, L., Muzny, D. M., Nelson, D. L., Nelson, D. R., Nelson, K. A., Nixon, K., Nusskern, D. R., Pacleb, J. M., Palazzolo, M., Pittman, G. S., Pan, S., Pollard, J., Puri, V., Reese, M. G., Reinert, K., Remington, K., Saunders, R. D., Scheeler, F., Shen, H., Shue, B. C., Siden-Kiamos, I., Simpson, M., Skupski, M. P., Smith, T., Spier, E., Spradling, A. C., Stapleton, M., Strong, R., Sun, E., Svirskas, R., Tector, C., Turner, R., Venter, E., Wang, A. H., Wang, X., Wang, Z. Y., Wassarman, D. A., Weinstock, G. M., Weissenbach, J., Williams, S. M., WoodageT, Worley, K. C., Wu, D., Yang, S., Yao, Q. A., Ye, J., Yeh, R. F., Zaveri, J. S., Zhan, M., Zhang, G., Zhao, Q., Zheng, L., Zheng, X. H., Zhong, F. N., Zhong, W., Zhou, X., Zhu, S., Zhu, X., Smith, H. O., Gibbs, R. A., Myers, E. W., Rubin, G. M. and Venter, J. C. (2000) The genome sequence of *Drosophila melanogaster* *Science* 287 (5461): 2185-95

Adams, S. M., Helps, N. R., Sharp, M. G., Brammar, W. J., Walker, R. A. and Varley, J. M. (1992) Isolation and characterization of a novel gene with differential expression in benign and malignant human breast tumours *Hum Mol Genet* 1 (2): 91-6

Akao, Y. and Matsuda, Y. (1996) Identification and chromosome mapping of the mouse homologue of the human gene (DDX6) that encodes a putative RNA helicase of the DEAD box protein family *Cytogenet Cell Genet* 75 (1): 38-44

Albig, W. and Doenecke, D. (1997) The human histone gene cluster at the D6S105 locus *Hum Genet* 101 (3): 284-94

Albig, W., Drabent, B., Burmester, N., Bode, C. and Doenecke, D. (1998) The haemochromatosis candidate gene HFE (HLA-H) of man and mouse is located in syntenic regions within the histone gene cluster *J Cell Biochem* 69 (2): 117-26

- Allcock, R. J., Martin, A. M. and Price, P. (2000) The mouse as a model for the effects of MHC genes on human disease *Immunol Today* 21 (7): 328-32
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. and Lipman, D. J. (1990) Basic local alignment search tool *J Mol Biol* 215 (3): 403-10
- Amadou, C. (1996) Structure et évolution du bras court du chromosome 6 humain: la région de classe I du Complexe Majeur d'Histocompatibilité et sa partie distale. Ph.D thesis, Université Paul Sabatier, Toulouse III.
- Amadou, C. (1999) Evolution of the Mhc class I region: the framework hypothesis *Immunogenetics* 49 (4): 362-7
- Amadou, C., Kumanovics, A., Jones, E. P., Lambracht-Washington, D., Yoshino, M. and Lindahl, K. F. (1999) The mouse major histocompatibility complex: some assembly required *Immunol Rev* 167 211-21
- Ansari-Lari, M. A., Oeltjen, J. C., Schwartz, S., Zhang, Z., Muzny, D. M., Lu, J., Gorrell, J. H., Chinault, A. C., Belmont, J. W., Miller, W. and Gibbs, R. A. (1998) Comparative sequence analysis of a gene-rich cluster at human chromosome 12p13 and its syntenic region in mouse chromosome 6 *Genome Res* 8 (1): 29-40
- Armstrong, L. C., Saenz, A. J. and Bornstein, P. (1999) Metaxin 1 interacts with metaxin 2, a novel related protein associated with the mammalian mitochondrial outer membrane *J Cell Biochem* 74 (1): 11-22
- Asai, H., Kasai, H., Matsuda, Y., Yamazaki, N., Nagawa, F., Sakano, H. and Tsuboi, A. (1996) Genomic structure and transcription of a murine odorant receptor gene: differential initiation of transcription in the olfactory and testicular cells *Biochem Biophys Res Commun* 221 (2): 240-7
- Babbitt, B. P., Allen, P. M., Matsueda, G., Haber, E. and Unanue, E. R. (1985) Binding of immunogenic peptides to Ia histocompatibility molecules *Nature* 317 (6035): 359-61
- Baier, H. and Korschning, S. (1994) Olfactory glomeruli in the zebrafish form an invariant pattern and are identifiable across animals *J Neurosci* 14 (1): 219-30
- Bailey, J. A., Carrel, L., Chakravarti, A. and Eichler, E. E. (2000) Molecular evidence for a relationship between LINE-1 elements and X chromosome inactivation: the Lyon repeat hypothesis *Proc Natl Acad Sci U S A* 97 (12): 6634-9
- Bankier, A. T., Weston, K. M. and Barrell, B. G. (1987) Random cloning and sequencing by the M13/dideoxynucleotide chain termination method *Methods Enzymol* 155 51-93
- Barth, A. L., Dugas, J. C. and Ngai, J. (1997) Noncoordinate expression of odorant receptor genes tightly linked in the zebrafish genome *Neuron* 19 (2): 359-69
- Barth, A. L., Justice, N. J. and Ngai, J. (1996) Asynchronous onset of odorant receptor expression in the developing zebrafish olfactory system *Neuron* 16 (1): 23-34
- Bates, E. E., Ravel, O., Dieu, M. C., Ho, S., Guret, C., Bridon, J. M., Ait-Yahia, S., Briere, F., Caux, C., Banchereau, J. and Lebecque, S. (1997) Identification and analysis of a novel member

of the ubiquitin family expressed in dendritic cells and mature B cells *Eur J Immunol* 27 (10): 2471-7

Baxendale, S., Abdulla, S., Elgar, G., Buck, D., Berks, M., Micklem, G., Durbin, R., Bates, G., Brenner, S. and Beck, S. (1995) Comparative sequence analysis of the human and pufferfish Huntington's disease genes *Nat Genet* 10 (1): 67-76

Bellefroid, E. J., Marine, J. C., Ried, T., Lecocq, P. J., Riviere, M., Amemiya, C., Poncelet, D. A., Coulie, P. G., de Jong, P., Szpirer, C. and et al. (1993) Clustered organization of homologous KRAB zinc-finger genes with enhanced expression in human T lymphoid cells *Embo J* 12 (4): 1363-74

Bellefroid, E. J., Poncelet, D. A., Lecocq, P. J., Revelant, O. and Martial, J. A. (1991) The evolutionarily conserved Kruppel-associated box domain defines a subfamily of eukaryotic multifingered proteins *Proc Natl Acad Sci U S A* 88 (9): 3608-12

Belluscio, L., Gold, G. H., Nemes, A. and Axel, R. (1998) Mice deficient in G(olf) are anosmic *Neuron* 20 (1): 69-81

Ben-Arie, N., Lancet, D., Taylor, C., Khen, M., Walker, N., Ledbetter, D. H., Carrozzo, R., Patel, K., Sheer, D., Lehrach, H. and et al. (1994) Olfactory receptor gene cluster on human chromosome 17: possible duplication of an ancestral receptor repertoire *Hum Mol Genet* 3 (2): 229-35

Ben-Chetrit, E., Gandy, B. J., Tan, E. M. and Sullivan, K. F. (1989) Isolation and characterization of a cDNA clone encoding the 60-kD component of the human SS-A/Ro ribonucleoprotein autoantigen *J Clin Invest* 83 (4): 1284-92

Berghard, A. and Dryer, L. (1998) A novel family of ancient vertebrate odorant receptors *J Neurobiol* 37 (3): 383-92

Bernardi, G., Olofsson, B., Filipski, J., Zerial, M., Salinas, J., Cuny, G., Meunier-Rotival, M. and Rodier, F. (1985) The mosaic genome of warm-blooded vertebrates *Science* 228 (4702): 953-8

Bernardi, G. (2001) Misunderstandings about isochores. Part 1 *Gene* 276 (1-2): 3-13

Biamonti, G., Ruggiu, M., Saccone, S., Della Valle, G. and Riva, S. (1994) Two homologous genes, originated by duplication, encode the human hnRNP proteins A2 and A1 *Nucleic Acids Res* 22 (11): 1996-2002

Bignon, C., Roux-Dosseto, M., Zeigler, M. E., Mattei, M. G., Lissitzky, J. C., Wicha, M. S. and Martin, P. M. (1991) Genomic analysis of the 67-kDa laminin receptor in normal and pathological tissues: circumstantial evidence for retroposon features *Genomics* 10 (2): 481-5

Birnboim, H. C. and Doly, J. (1979) A rapid alkaline extraction procedure for screening recombinant plasmid DNA *Nucleic Acids Res* 7 (6): 1513-23

Bjorkman, P. J., Saper, M. A., Samraoui, B., Bennett, W. S., Strominger, J. L. and Wiley, D. C. (1987) The foreign antigen binding site and T cell recognition regions of class I histocompatibility antigens *Nature* 329 (6139): 512-8

- Blanpain, C., Lee, B., Vakili, J., Doranz, B. J., Govaerts, C., Migeotte, I., Sharron, M., Dupriez, V., Vassart, G., Doms, R. W. and Parmentier, M. (1999) Extracellular cysteines of CCR5 are required for chemokine binding, but dispensable for HIV-1 coreceptor activity *J Biol Chem* 274 (27): 18902-8
- Blow, J. J. and Laskey, R. A. (1988) A role for the nuclear envelope in controlling DNA replication within the cell cycle *Nature* 332 (6164): 546-8
- Bodmer, J. G., Marsh, S. G., Albert, E. D., Bodmer, W. F., Bontrop, R. E., Dupont, B., Erlich, H. A., Hansen, J. A., Mach, B., Mayr, W. R., Parham, P., Petersdorf, E. W., Sasazuki, T., Schreuder, G. M., Strominger, J. L., Svejgaard, A. and Terasaki, P. I. (1999) Nomenclature for factors of the HLA system, 1998 *Vox Sang* 77 (3): 164-91
- Bonfield, J. K., Smith, K. and Staden, R. (1995) A new DNA sequence assembly program *Nucleic Acids Res* 23 (24): 4992-9
- Brinon, J. G., Martinez-Guijarro, F. J., Bravo, I. G., Arevalo, R., Crespo, C., Okazaki, K., Hidaka, H., Aijon, J. and Alonso, J. R. (1999) Coexpression of neurocalcin with other calcium-binding proteins in the rat main olfactory bulb *J Comp Neurol* 407 (3): 404-14
- Britten, R. J., Baron, W. F., Stout, D. B. and Davidson, E. H. (1988) Sources and evolution of human Alu repeated sequences *Proc Natl Acad Sci U S A* 85 (13): 4770-4
- Brookfield, J. F. (2001) Selection on Alu sequences? *Curr Biol* 11 (22): R900-1
- Brunet, L. J., Gold, G. H. and Ngai, J. (1996) General anosmia caused by a targeted disruption of the mouse olfactory cyclic nucleotide-gated cation channel *Neuron* 17 (4): 681-93
- Brzustowicz, L. M., Lehner, T., Castilla, L. H., Penchaszadeh, G. K., Wilhelmsen, K. C., Daniels, R., Davies, K. E., Leppert, M., Ziter, F., Wood, D. and et al. (1990) Genetic mapping of chronic childhood-onset spinal muscular atrophy to chromosome 5q11.2-13.3 *Nature* 344 (6266): 540-1
- Buck, L. and Axel, R. (1991) A novel multigene family may encode odorant receptors: a molecular basis for odor recognition *Cell* 65 (1): 175-87
- Buck, L. B. (1992) The olfactory multigene family *Curr Opin Neurobiol* 2 (3): 282-8
- Buck, L. B. (1996) Information coding in the mammalian olfactory system *Cold Spring Harb Symp Quant Biol* 61 147-55
- Buck, L. B. (2000) The molecular architecture of odor and pheromone sensing in mammals *Cell* 100 (6): 611-8
- Bulger, M., Bender, M. A., van Doorninck, J. H., Wertman, B., Farrell, C. M., Felsenfeld, G., Groudine, M. and Hardison, R. (2000) Comparative structural and functional analysis of the olfactory receptor genes flanking the human and mouse beta-globin gene clusters *Proc Natl Acad Sci U S A* 97 (26): 14560-5
- Burge, C. and Karlin, S. (1997) Prediction of complete gene structures in human genomic DNA *J Mol Biol* 268 (1): 78-94

- Burge, C. B. and Karlin, S. (1998) Finding the genes in genomic DNA *Curr Opin Struct Biol* 8 (3): 346-54
- Byrd, C. A., Jones, J. T., Quattro, J. M., Rogers, M. E., Brunjes, P. C. and Vogt, R. G. (1996) Ontogeny of odorant receptor gene expression in zebrafish, *Danio rerio* *J Neurobiol* 29 (4): 445-58
- Carrier, A., Nguyen, C., Victorero, G., Granjeaud, S., Rocha, D., Bernard, K., Miazek, A., Ferrier, P., Malissen, M., Naquet, P., Malissen, B. and Jordan, B. R. (1999) Differential gene expression in CD3epsilon- and RAG1-deficient thymuses: definition of a set of genes potentially involved in thymocyte maturation *Immunogenetics* 50 (5-6): 255-70
- Carver, E. A., Issel-Tarver, L., Rine, J., Olsen, A. S. and Stubbs, L. (1998) Location of mouse and human genes corresponding to conserved canine olfactory receptor gene subfamilies *Mamm Genome* 9 (5): 349-54
- Cavin Perier, R., Junier, T. and Bucher, P. (1998) The Eukaryotic Promoter Database EPD *Nucleic Acids Res* 26 (1): 353-7
- Chess, A., Simon, I., Cedar, H. and Axel, R. (1994) Allelic inactivation regulates olfactory receptor gene expression *Cell* 78 (5): 823-34
- Chong, J. P., Thommes, P. and Blow, J. J. (1996) The role of MCM/P1 proteins in the licensing of DNA replication *Trends Biochem Sci* 21 (3): 102-6
- Christiansen, O. B., Riisom, K., Lauritsen, J. G. and Grunnet, N. (1989) No increased histocompatibility antigen-sharing in couples with idiopathic habitual abortions *Hum Reprod* 4 (2): 160-2
- Chu, W. M., Ballard, R., Carpick, B. W., Williams, B. R. and Schmid, C. W. (1998) Potential Alu function: regulation of the activity of double-stranded RNA-activated kinase PKR *Mol Cell Biol* 18 (1): 58-68
- Clancy, A. N., Coquelin, A., Macrides, F., Gorski, R. A. and Noble, E. P. (1984) Sexual behavior and aggression in male mice: involvement of the vomeronasal system *J Neurosci* 4 (9): 2222-9
- Conzelmann, S., Levai, O., Bode, B., Eiselt, U., Raming, K., Breer, H. and Strotmann, J. (2000) A novel brain receptor is expressed in a distinct population of olfactory sensory neurons *Eur J Neurosci* 12 (11): 3926-34
- Crowe, M. L., Perry, B. N. and Connerton, I. F. (1996) Olfactory receptor-encoding genes and pseudogenes are expressed in humans *Gene* 169 (2): 247-9
- Cuny, G., Soriano, P., Macaya, G. and Bernardi, G. (1981) The major components of the mouse and human genomes. 1. Preparation, basic properties and compositional heterogeneity *Eur J Biochem* 115 (2): 227-33
- Dalvi, A. and Rodgers, R. J. (1996) GABAergic influences on plus-maze behaviour in mice *Psychopharmacology (Berl)* 128 (4): 380-97

- Davies, B., Feo, S., Heard, E. and Fried, M. (1989) A strategy to detect and isolate an intron-containing gene in the presence of multiple processed pseudogenes *Proc Natl Acad Sci U S A* 86 (17): 6691-5
- Dawson, T. M., Arriza, J. L., Jaworsky, D. E., Borisy, F. F., Attramadal, H., Lefkowitz, R. J. and Ronnett, G. V. (1993) Beta-adrenergic receptor kinase-2 and beta-arrestin-2 as mediators of odorant-induced desensitization *Science* 259 (5096): 825-9
- Dayhoff, M. O. (1976) The origin and evolution of protein superfamilies *Fed Proc* 35 (10): 2132-8
- Dear, T. N., Campbell, K. and Rabbitts, T. H. (1991) Molecular cloning of putative odorant-binding and odorant-metabolizing proteins *Biochemistry* 30 (43): 10376-82
- Del Punta, K., Rothman, A., Rodriguez, I. and Mombaerts, P. (2000) Sequence diversity and genomic organization of vomeronasal receptor genes in the mouse *Genome Res* 10 (12): 1958-67
- den Hollander, A. I., Ghiani, M., de Kok, Y. J., Wijnholds, J., Ballabio, A., Cremers, F. P. and Broccoli, V. (2002) Isolation of Crb1, a mouse homologue of *Drosophila* crumbs, and analysis of its expression pattern in eye and brain *Mech Dev* 110 (1-2): 203-7
- Denzin, L. K. and Cresswell, P. (1995) HLA-DM induces CLIP dissociation from MHC class II alpha beta dimers and facilitates peptide loading *Cell* 82 (1): 155-65
- Dizhoor, A. M., Ray, S., Kumar, S., Niemi, G., Spencer, M., Brolley, D., Walsh, K. A., Philipov, P. P., Hurley, J. B. and Stryer, L. (1991) Recoverin: a calcium sensitive activator of retinal rod guanylate cyclase *Science* 251 (4996): 915-8
- Doherty, P. C. and Zinkernagel, R. M. (1975) Enhanced immunological surveillance in mice heterozygous at the H-2 gene complex *Nature* 256 (5512): 50-2
- Dong, X., Han, S., Zylka, M. J., Simon, M. I. and Anderson, D. J. (2001) A diverse family of GPCRs expressed in specific subsets of nociceptive sensory neurons *Cell* 106 (5): 619-32
- Down, T. A. and Hubbard, T. J. (2002) Computational detection and location of transcription start sites in mammalian genomic DNA *Genome Res* 12 (3): 458-61
- Dreyer, W. J. (1998) The area code hypothesis revisited: olfactory receptors and other related transmembrane receptors may function as the last digits in a cell surface code for assembling embryos *Proc Natl Acad Sci U S A* 95 (16): 9072-7
- Driscoll, J., Brown, M. G., Finley, D. and Monaco, J. J. (1993) MHC-linked LMP gene products specifically alter peptidase activities of the proteasome *Nature* 365 (6443): 262-4
- Drutel, G., Arrang, J. M., Diaz, J., Wisnewskey, C., Schwartz, K. and Schwartz, J. C. (1995) Cloning of OL1, a putative olfactory receptor and its expression in the developing rat heart *Receptors Channels* 3 (1): 33-40
- Duboule, D. (1998) Vertebrate hox gene regulation: clustering and/or colinearity? *Curr Opin Genet Dev* 8 (5): 514-8

- Duboule, D. and Dolle, P. (1989) The structural and functional organization of the murine HOX gene family resembles that of *Drosophila* homeotic genes *Embo J* 8 (5): 1497-505
- Dulac, C. and Axel, R. (1995) A novel family of genes encoding putative pheromone receptors in mammals *Cell* 83 (2): 195-206
- Ehlers, A., Beck, S., Forbes, S. A., Trowsdale, J., Volz, A., Younger, R. and Ziegler, A. (2000) MHC-linked olfactory receptor loci exhibit polymorphism and contribute to extended HLA/OR-haplotypes *Genome Res* 10 (12): 1968-78
- Eyre-Walker, A. and Hurst, L. D. (2001) The evolution of isochores *Nat Rev Genet* 2 (7): 549-55
- Fan, W., Cai, W., Parimoo, S., Schwarz, D. C., Lennon, G. G. and Weissman, S. M. (1996) Identification of seven new human MHC class I region genes around the HLA-F locus *Immunogenetics* 44 (2): 97-103
- Fan, W., Liu, Y. C., Parimoo, S. and Weissman, S. M. (1995) Olfactory receptor-like genes are located in the human major histocompatibility complex *Genomics* 27 (1): 119-23
- Feingold, E. A., Penny, L. A., Nienhuis, A. W. and Forget, B. G. (1999) An olfactory receptor gene is located in the extended human beta-globin gene cluster and is expressed in erythroid cells *Genomics* 61 (1): 15-23
- Felsenstein, J. (1985) Confidence-Limits On Phylogenies - an Approach Using the Bootstrap. *Evolution* 39(4):783-791.
- Felsenstein, J. (1989) PHYLIP -- Phylogeny Inference Package (Version 3.2). *Cladistics* 5: 164-166.
- Firestein, S. (2001) How the olfactory system makes sense of scents *Nature* 413 (6852): 211-8
- Fitch, D. H., Bailey, W. J., Tagle, D. A., Goodman, M., Sieu, L. and Slightom, J. L. (1991) Duplication of the gamma-globin gene mediated by L1 long interspersed repetitive elements in an early ancestor of simian primates *Proc Natl Acad Sci U S A* 88 (16): 7396-400
- Fitch, W. M. 1971. Toward defining the course of evolution: minimum change for a specified tree topology. *Systematic Zoology* 20: 406-416.
- Freitag, J., Krieger, J., Strotmann, J. and Breer, H. (1995) Two classes of olfactory receptors in *Xenopus laevis* *Neuron* 15 (6): 1383-92
- Friedrich, R. W. and Korschning, S. I. (1997) Combinatorial and chemotopic odorant coding in the zebrafish olfactory bulb visualized by optical imaging *Neuron* 18 (5): 737-52
- Galtier, N., Gouy, M. and Gautier, C. (1996) SEAVIEW and PHYLO_WIN: two graphic tools for sequence alignment and molecular phylogeny *Comput Appl Biosci* 12 (6): 543-8
- Gardiner-Garden, M., Ballesteros, M., Gordon, M. and Tam, P. P. (1998) Histone- and protamine-DNA association: conservation of different patterns within the beta-globin domain in human sperm *Mol Cell Biol* 18 (6): 3350-6

- Gatewood, J. M., Cook, G. R., Balhorn, R., Bradbury, E. M. and Schmid, C. W. (1987) Sequence-specific packaging of DNA in human sperm chromatin *Science* 236 (4804): 962-4
- Gaunt, S. J., Krumlauf, R. and Duboule, D. (1989) Mouse homeo-genes within a subfamily, Hox-1.4, -2.6 and -5.1, display similar anteroposterior domains of expression in the embryo, but show stage- and tissue-dependent differences in their regulation *Development* 107 (1): 131-41
- Gautier-Courteille, C., Salanova, M. and Conti, M. (1998) The olfactory adenylyl cyclase III is expressed in rat germ cells during spermiogenesis *Endocrinology* 139 (5): 2588-99
- Gehlsen, K. R., Dillner, L., Engvall, E. and Ruoslahti, E. (1988) The human laminin receptor is a member of the integrin family of cell adhesion receptors *Science* 241 (4870): 1228-9
- Gilad, Y., Segre, D., Skorecki, K., Nachman, M. W., Lancet, D. and Sharon, D. (2000) Dichotomy of single-nucleotide polymorphism haplotypes in olfactory receptor genes and pseudogenes *Nat Genet* 26 (2): 221-4
- Gimelbrant, A. A. and McClintock, T. S. (1997) A nuclear matrix attachment region is highly homologous to a conserved domain of olfactory receptors *J Mol Neurosci* 9 (1): 61-3
- Giorgi, D., Friedman, C., Trask, B. J. and Rouquier, S. (2000) Characterization of nonfunctional V1R-like pheromone receptor sequences in human *Genome Res* 10 (12): 1979-85
- Glusman, G., Bahar, A., Sharon, D., Pilpel, Y., White, J. and Lancet, D. (2000) The olfactory receptor gene superfamily: data mining, classification, and nomenclature *Mamm Genome* 11 (11): 1016-23
- Glusman, G., Clifton, S., Roe, B. and Lancet, D. (1996) Sequence analysis in the olfactory receptor gene cluster on human chromosome 17: recombinatorial events affecting receptor diversity *Genomics* 37 (2): 147-60
- Glusman, G., Sosinsky, A., Ben-Asher, E., Avidan, N., Sonkin, D., Bahar, A., Rosenthal, A., Clifton, S., Roe, B., Ferraz, C., Demaille, J. and Lancet, D. (2000) Sequence, structure, and evolution of a complete human olfactory receptor gene cluster *Genomics* 63 (2): 227-45
- Glusman, G., Yanai, I., Rubin, I. and Lancet, D. (2001) The complete human olfactory subgenome *Genome Res* 11 (5): 685-702
- Gobin, S. J. and van den Elsen, P. J. (2000) Transcriptional regulation of the MHC class Ib genes HLA-E, HLA-F, and HLA-G *Hum Immunol* 61 (11): 1102-7
- Goei, V. L., Choi, J., Ahn, J., Bowlus, C. L., Raha-Chowdhury, R. and Gruen, J. R. (1998) Human gamma-aminobutyric acid B receptor gene: complementary DNA cloning, expression, chromosomal location, and genomic organization *Biol Psychiatry* 44 (8): 659-66
- Gonnet, G. H., Cohen, M. A. and Benner, S. A. (1992) Exhaustive matching of the entire protein sequence database *Science* 256 (5062): 1443-5
- Gordon, D., Abajian, C. and Green, P. (1998) Consed: a graphical tool for sequence finishing *Genome Res* 8 (3): 195-202

- Graham, A., Papalopulu, N. and Krumlauf, R. (1989) The murine and Drosophila homeobox gene complexes have common features of organization and expression *Cell* 57 (3): 367-78
- Graham, F. L., Smiley, J., Russell, W. C. and Nairn, R. (1977) Characteristics of a human cell line transformed by DNA from human adenovirus type 5 *J Gen Virol* 36 (1): 59-74
- Gregory, S. G., Howell, G. H. and Bentley, D. R. (1997) Genome Mapping by Fluorescent Fingerprinting *Genome Research* 7 1162-1168
- Grifa, A., Totaro, A., Rommens, J. M., Carella, M., Roetto, A., Borgato, L., Zelante, L. and Gasparini, P. (1998) GABA (gamma-amino-butyric acid) neurotransmission: identification and fine mapping of the human GABAB receptor gene *Biochem Biophys Res Commun* 250 (2): 240-5
- Gross-Isseroff, R., Ophir, D., Bartana, A., Voet, H. and Lancet, D. (1992) Evidence for genetic determination in human twins of olfactory thresholds for a standard odorant *Neurosci Lett* 141 (1): 115-8
- Grosveld, F. (1999) Activation by locus control regions? *Curr Opin Genet Dev* 9 (2): 152-7
- Grosveld, F., van Assendelft, G. B., Greaves, D. R. and Kollias, G. (1987) Position-independent, high-level expression of the human beta-globin gene in transgenic mice *Cell* 51 (6): 975-85
- Gruen, J. R., Goei, V. L., Summers, K. M., Capossela, A., Powell, L., Halliday, J., Zoghbi, H., Shukla, H. and Weissman, S. M. (1992) Physical and genetic mapping of the telomeric major histocompatibility complex region in man and relevance to the primary hemochromatosis gene (HFE) *Genomics* 14 (2): 232-40
- Guigo, R., Agarwal, P., Abril, J. F., Burset, M. and Fickett, J. W. (2000) An assessment of gene prediction accuracy in large DNA sequences *Genome Res* 10 (10): 1631-42
- Gunther, E. and Walter, L. (2001) *The major histocompatibility complex of the rat (Rattus norvegicus)* *Immunogenetics* 7 (53): 520-42
- Haino, M., Hayashida, H., Miyata, T., Shin, E. K., Matsuda, F., Nagaoka, H., Matsumura, R., Takaishi, S., Fukita, Y., Fujikura, J. and et al. (1994) Comparison and evolution of human immunoglobulin VH segments located in the 3' 0.8-megabase region. Evidence for unidirectional transfer of segmental gene sequences *J Biol Chem* 269 (4): 2619-26
- Hall, L., Williams, K., Perry, A. C., Frayne, J. and Jury, J. A. (1998) The majority of human glutathione peroxidase type 5 (GPX5) transcripts are incorrectly spliced: implications for the role of GPX5 in the male reproductive tract *Biochem J* 333 (Pt 1) 5-9
- Hallberg, E., Wozniak, R. W. and Blobel, G. (1993) An integral membrane protein of the pore membrane domain of the nuclear envelope contains a nucleoporin-like region *J Cell Biol* 122 (3): 513-21
- Hanke, J., Brett, D., Zastrow, I., Aydin, A., Delbruck, S., Lehmann, G., Luft, F., Reich, J. and Bork, P. (1999) Alternative splicing of human genes: more the rule than the exception? *Trends Genet* 15 (10): 389-90

- Hardison, R. C., Oeltjen, J. and Miller, W. (1997) Long human-mouse sequence alignments reveal novel regulatory elements: a reason to sequence the mouse genome *Genome Res* 7 (10): 959-66
- Hellmann-Blumberg, U., Hintz, M. F., Gatewood, J. M. and Schmid, C. W. (1993) Developmental differences in methylation of human Alu repeats *Mol Cell Biol* 13 (8): 4523-30
- Henikoff, S. and Henikoff, J. G. (1992) Amino acid substitution matrices from protein blocks *Proc Natl Acad Sci U S A* 89 (22): 10915-9
- Henry, J., Mather, I. H., McDermott, M. F. and Pontarotti, P. (1998) B30.2-like domain proteins: update and new insights into a rapidly expanding family of proteins *Mol Biol Evol* 15 (12): 1696-705
- Herrada, G. and Dulac, C. (1997) A novel family of putative pheromone receptors in mammals with a topographically organized and sexually dimorphic distribution *Cell* 90 (4): 763-73
- Hildebrand, J. G. and Shepherd, G. M. (1997) Mechanisms of olfactory discrimination: converging evidence for common principles across phyla *Annu Rev Neurosci* 20 595-631
- Hillier, L. D., Lennon, G., Becker, M., Bonaldo, M. F., Chiapelli, B., Chissoe, S., Dietrich, N., DuBuque, T., Favella, A., Gish, W., Hawkins, M., Hultman, M., Kucaba, T., Lacy, M., Le, M., Le, N., Mardis, E., Moore, B., Morris, M., Parsons, J., Prange, C., Rifkin, L., Rohlfing, T., Schellenberg, K., Marra, M. and et al. (1996) Generation and analysis of 280,000 human expressed sequence tags *Genome Res* 6 (9): 807-28
- Horikawa, I., Tanaka, H., Yuasa, Y., Suzuki, M. and Oshimura, M. (1995) Molecular cloning of a novel human cDNA on chromosome 1q21 and its mouse homolog encoding a nuclear protein with DNA-binding ability *Biochem Biophys Res Commun* 208 (3): 999-1007
- Horton, R., Niblett, D., Milne, S., Palmer, S., Tubby, B., Trowsdale, J. and Beck, S. (1998) Large-scale sequence comparisons reveal unusually high levels of variation in the HLA-DQB1 locus in the class II region of the human MHC *J Mol Biol* 282 (1): 71-97
- Howe, K., Bateman, A. and Durbin, R. (unpublished) QuickTree: A program for building huge Neighbor-Joining trees of protein sequences
- Hubbard, T., Barker, D., Birney, E., Cameron, G., Chen, Y., Clark, L., Cox, T., Cuff, J., Curwen, V., Down, T., Durbin, R., Eyras, E., Gilbert, J., Hammond, M., Huminiecki, L., Kasprzyk, A., Lehvaslaiho, H., Lijnzaad, P., Melsopp, C., Mongin, E., Pettett, R., Pocock, M., Potter, S., Rust, A., Schmidt, E., Searle, S., Slater, G., Smith, J., Spooner, W., Stabenau, A., Stalker, J., Stupka, E., Ureta-Vidal, A., Vastrik, I. and Clamp, M. (2002) The Ensembl genome database project *Nucleic Acids Res* 30 (1): 38-41
- Huebner, K., Cannizzaro, L. A., Nakamura, T., Hilova, J., Mariage-Samson, R., Hecht, F., Hill, M. and Croce, C. M. (1988) A rearranged transforming gene, tre, is made up of human sequences derived from chromosome regions 5q, 17q and 18q *Oncogene* 3 (4): 449-55
- Hughes, A. L. and Nei, M. (1988) Pattern of nucleotide substitution at major histocompatibility complex class I loci reveals overdominant selection *Nature* 335 (6186): 167-70

- Ikematsu, N., Yoshida, Y., Kawamura-Tsuzuku, J., Ohsugi, M., Onda, M., Hirai, M., Fujimoto, J. and Yamamoto, T. (1999) Tob2, a novel anti-proliferative Tob/BTG1 family member, associates with a component of the CCR4 transcriptional regulatory complex capable of binding cyclin-dependent kinases *Oncogene* 18 (52): 7432-41
- Isomura, T., Tamiya-Koizumi, K., Suzuki, M., Yoshida, S., Taniguchi, M., Matsuyama, M., Ishigaki, T., Sakuma, S. and Takahashi, M. (1992) RFP is a DNA binding protein associated with the nuclear matrix *Nucleic Acids Res* 20 (20): 5305-10
- Isono, K., McIninch, J. D. and Borodovsky, M. (1994) Characteristic features of the nucleotide sequences of yeast mitochondrial ribosomal protein genes as analyzed by computer program GeneMark *DNA Res* 1 (6): 263-9
- Issel-Tarver, L. and Rine, J. (1996) Organization and expression of canine olfactory receptor genes *Proc Natl Acad Sci U S A* 93 (20): 10897-902
- Issel-Tarver, L. and Rine, J. (1997) The evolution of mammalian olfactory receptor genes *Genetics* 145 (1): 185-95
- Iwata, Y., Nakayama, A., Murakami, H., Iida, K., Iwashita, T., Asai, N. and Takahashi, M. (1999) Characterization of the promoter region of the human RFP gene *Biochem Biophys Res Commun* 261 (2): 381-4
- Jabbari, K. and Bernardi, G. (1998) CpG doublets, CpG islands and Alu repeats in long human DNA sequences from different isochore families *Gene* 224 (1-2): 123-7
- Jack, L. J. and Mather, I. H. (1990) Cloning and analysis of cDNA encoding bovine butyrophilin, an apical glycoprotein expressed in mammary tissue and secreted in association with the milk-fat globule membrane during lactation *J Biol Chem* 265 (24): 14481-6
- Jareborg, N., Birney, E. and Durbin, R. (1999) Comparative analysis of noncoding regions of 77 orthologous mouse and human gene pairs *Genome Res* 9 (9): 815-24
- Johns, M. A., Feder, H. H., Komisaruk, B. R. and Mayer, A. D. (1978) Urine-induced reflex ovulation in anovulatory rats may be a vomeronasal effect *Nature* 272 (5652): 446-8
- Jones, D. T., Taylor, W. R. and Thornton, J. M. (1992) The rapid generation of mutation data matrices from protein sequences *Comput Appl Biosci* 8 (3): 275-82
- Jones, E. P., Kumanovics, A., Yoshino, M. and Fischer Lindahl, K. (1999) Mhc class I and non-class I gene organization in the proximal H2-M region of the mouse *Immunogenetics* 49 (3): 183-95
- Jones, E. P., Xiao, H., Schultz, R. A., Flaherty, L., Trachtulec, Z., Vincek, V., Larin, Z., Lehrach, H. and Lindahl, K. F. (1995) MHC class I gene organization in > 1.5-Mb YAC contigs from the H2-M region *Genomics* 27 (1): 40-51
- Jurka, J. (2000) Repbase update: a database and an electronic journal of repetitive elements *Trends Genet* 16 (9): 418-20

- Jurka, J. and Smith, T. (1988) A fundamental division in the Alu family of repeated sequences *Proc Natl Acad Sci U S A* 85 (13): 4775-8
- Jurka, J., Kapitonov, V. V., Klonowski, P., Walichiewicz, J. and Smit, A. F. (1996) Identification of new medium reiteration frequency repeats in the genomes of Primates, Rodentia and Lagomorpha *Genetica* 98 (3): 235-47
- Kagie, M. J., Kleijer, W. J., Huijmans, J. G., Maaswinkel-Mooy, P. and Kanhai, H. H. (1992) beta-Glucuronidase deficiency as a cause of fetal hydrops *Am J Med Genet* 42 (5): 693-5
- Kamitani, T., Kito, K., Nguyen, H. P., Fukuda-Kamitani, T. and Yeh, E. T. (1998) Characterization of a second member of the sentrin family of ubiquitin-like proteins *J Biol Chem* 273 (18): 11349-53
- Kapitonov, V. and Jurka, J. (1996) The age of Alu subfamilies *J Mol Evol* 42 (1): 59-65
- Kapitonov, V. V., Holmquist, G. P. and Jurka, J. (1998) L1 repeat is a basic unit of heterochromatin satellites in cetaceans *Mol Biol Evol* 15 (5): 611-2
- Kasture, S. B., Mandhane, S. N. and Chopde, C. T. (1996) Baclofen-induced catatonia: modification by serotonergic agents *Neuropharmacology* 35 (5): 595-8
- Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T. J., Tanaka, T. and Page, D. C. (1998) A map of 75 human ribosomal protein genes *Genome Res* 8 (5): 509-23
- Kent, W. J. and Haussler, D. (2001) Assembly of the working draft of the human genome with GigAssembler *Genome Res* 11 (9): 1541-8
- Kerr, D. I. and Ong, J. (1995) GABAB receptors *Pharmacol Ther* 67 (2): 187-246
- Keverne, E. B. (1999) The vomeronasal organ *Science* 286 (5440): 716-20
- Kleene, S. J. and Gesteland, R. C. (1991) Calcium-activated chloride conductance in frog olfactory cilia *J Neurosci* 11 (11): 3624-9
- Klingenhoff, A., Frech, K., Quandt, K. and Werner, T. (1999) Functional promoter modules can be detected by formal models independent of overall nucleotide sequence similarity *Bioinformatics* 15 (3): 180-6
- Knowles, B. B., Solter, D., Trinchieri, G., Maloney, K. M., Ford, S. R. and Aden, D. P. (1977) Complement-mediated antiserum cytotoxic reactions to human chromosome 7 coded antigen(s): immunoselection of rearranged human chromosome 7 in human-mouse somatic cell hybrids *J Exp Med* 145 (2): 314-26
- Kobilka, B. (1992) Adrenergic receptors as models for G protein-coupled receptors *Annu Rev Neurosci* 15 87-114
- Kobilka, B. K., Kobilka, T. S., Daniel, K., Regan, J. W., Caron, M. G. and Lefkowitz, R. J. (1988) Chimeric alpha 2-beta 2-adrenergic receptors: delineation of domains involved in effector coupling and ligand binding specificity *Science* 240 (4857): 1310-6

- Kochanek, S., Renz, D. and Doerfler, W. (1993) DNA methylation in the Alu sequences of diploid and haploid primary human cells *Embo J* 12 (3): 1141-51
- Koop, B. F. and Hood, L. (1994) Striking sequence similarity over almost 100 kilobases of human and mouse T-cell receptor DNA *Nat Genet* 7 (1): 48-53
- Koshimoto, H., Katoh, K., Yoshihara, Y., Nemoto, Y. and Mori, K. (1994) Immunohistochemical demonstration of embryonic expression of an odor receptor protein and its zonal distribution in the rat olfactory epithelium *Neurosci Lett* 169 (1-2): 73-6
- Kratz, E., Dugas, J. C. and Ngai, J. (2002) Odorant receptor gene regulation: implications from genomic organization *Trends Genet* 18 (1): 29-34
- Krautwurst, D., Yau, K. W. and Reed, R. R. (1998) Identification of ligands for olfactory receptors by functional expression of a receptor library *Cell* 95 (7): 917-26
- Ku, N. O., Wright, T. L., Terrault, N. A., Gish, R. and Omary, M. B. (1997) Mutation of human keratin 18 in association with cryptogenic cirrhosis *J Clin Invest* 99 (1): 19-23
- Kubick, S., Strotmann, J., Andreini, I. and Breer, H. (1997) Subfamily of olfactory receptors characterized by unique structural features and expression patterns *J Neurochem* 69 (2): 465-75
- Kulski, J. K. and Dawkins, R. L. (1999) The P5 multicopy gene family in the MHC is related in sequence to human endogenous retroviruses HERV-L and HERV-16 *Immunogenetics* 49 (5): 404-12
- Kurahashi, T. and Menini, A. (1997) Mechanism of odorant adaptation in the olfactory receptor cell *Nature* 385 (6618): 725-9
- Ladenburger, E. M., Fackelmayer, F. O., Hameister, H. and Knippers, R. (1997) MCM4 and PRKDC, human genes encoding proteins MCM4 and DNA-PKcs, are close neighbours located on chromosome 8q12-->q13 *Cytogenet Cell Genet* 77 (3-4): 268-70
- Laity, J. H., Lee, B. M. and Wright, P. E. (2001) Zinc finger proteins: new insights into structural and functional diversity *Curr Opin Struct Biol* 11 (1): 39-46
- Lancet, D., Ben-Arie, N., Cohen, S., Gat, U., Gross-Isseroff, R., Horn-Saban, S., Khen, M., Lehrach, H., Natochin, M., North, M. and et al. (1993) Olfactory receptors: transduction, diversity, human psychophysics and genome analysis *Ciba Found Symp* 179 131-41; discussion 141-6
- Landsman, D., Soares, N., Gonzalez, F. J. and Bustin, M. (1986) Chromosomal protein HMG-17. Complete human cDNA sequence and evidence for a multigene family *J Biol Chem* 261 (16): 7479-84
- Lane, R. P., Cutforth, T., Young, J., Athanasiou, M., Friedman, C., Rowen, L., Evans, G., Axel, R., Hood, L. and Trask, B. J. (2001) Genomic analysis of orthologous mouse and human olfactory receptor loci *Proc Natl Acad Sci U S A* 98 (13): 7390-5

- Lane, R. P., Roach, J. C., Lee, I. Y., Boysen, C., Smit, A., Trask, B. J. and Hood, L. (2002) Genomic Analysis of the Olfactory Receptor Region of the Mouse and Human T-Cell Receptor alpha/delta Loci *Genome Res* 12 (1): 81-7
- Lapenta, V., Chiurazzi, P., van der Spek, P., Pizzuti, A., Hanaoka, F. and Brahe, C. (1997) SMT3A, a human homologue of the *S. cerevisiae* SMT3 gene, maps to chromosome 21qter and defines a novel gene family *Genomics* 40 (2): 362-6
- Lapidot, M., Pilpel, Y., Gilad, Y., Falcovitz, A., Sharon, D., Haaf, T. and Lancet, D. (2001) Mouse-human orthology relationships in an olfactory receptor gene cluster *Genomics* 3 (71): 296-306
- Latif, F., Tory, K., Gnarra, J., Yao, M., Duh, F. M., Orcutt, M. L., Stackhouse, T., Kuzmin, I., Modi, W., Geil, L. and et al. (1993) Identification of the von Hippel-Lindau disease tumor suppressor gene *Science* 260 (5112): 1317-20
- Le Gouill, C., Parent, J. L., Rola-Pleszczynski, M. and Stankova, J. (1997) Role of the Cys90, Cys95 and Cys173 residues in the structure and function of the human platelet-activating factor receptor *FEBS Lett* 402 (2-3): 203-8
- Leibovici, M., Lapointe, F., Aletta, P. and Ayer-Le Lievre, C. (1996) Avian olfactory receptors: differentiation of olfactory neurons under normal and experimental conditions *Dev Biol* 175 (1): 118-31
- Levy, N. S., Bakalyar, H. A. and Reed, R. R. (1991) Signal transduction in olfactory neurons *J Steroid Biochem Mol Biol* 39 (4B): 633-7
- Lewis, E. B. (1978) A gene complex controlling segmentation in *Drosophila* *Nature* 276 (5688): 565-70
- Lieber, M. (1996) Immunoglobulin diversity: rearranging by cutting and repairing *Curr Biol* 6 (2): 134-6
- Lin, H. and Grosschedl, R. (1995) Failure of B-cell differentiation in mice lacking the transcription factor EBF *Nature* 376 (6537): 263-7
- Linardopoulou, E., Mefford, H. C., Nguyen, O., Friedman, C., van den Engh, G., Farwell, D. G., Coltrera, M. and Trask, B. J. (2001) Transcriptional activity of multiple copies of a subtelomERICALLY located olfactory receptor gene that is polymorphic in number and location *Hum Mol Genet* 10 (21): 2373-83
- Lindahl, K. F., Byers, D. E., Dabhi, V. M., Hovik, R., Jones, E. P., Smith, G. P., Wang, C. R., Xiao, H. and Yoshino, M. (1997) H2-M3, a full-service class Ib histocompatibility antigen *Annu Rev Immunol* 15 851-79
- Ling, K., Wang, P., Zhao, J., Wu, Y. L., Cheng, Z. J., Wu, G. X., Hu, W., Ma, L. and Pei, G. (1999) Five-transmembrane domains appear sufficient for a G protein-coupled receptor: functional five-transmembrane domain chemokine receptors *Proc Natl Acad Sci U S A* 96 (14): 7922-7

- Liu, W. M., Chu, W. M., Choudary, P. V. and Schmid, C. W. (1995) Cell stress and translational inhibitors transiently increase the abundance of mammalian SINE transcripts *Nucleic Acids Res* 23 (10): 1758-65
- Liu, Y. C., Pan, J., Zhang, C., Fan, W., Collinge, M., Bender, J. R. and Weissman, S. M. (1999) A MHC-encoded ubiquitin-like protein (FAT10) binds noncovalently to the spindle assembly checkpoint protein MAD2 *Proc Natl Acad Sci U S A* 96 (8): 4313-8
- Lomas, D. E. and Keverne, E. B. (1982) Role of the vomeronasal organ and prolactin in the acceleration of puberty in female mice *J Reprod Fertil* 66 (1): 101-7
- Loveland, B., Wang, C. R., Yonekawa, H., Hermel, E. and Lindahl, K. F. (1990) Maternally transmitted histocompatibility antigen of mice: a hydrophobic peptide of a mitochondrially encoded protein *Cell* 60 (6): 971-80
- Lyon, M. F. (1998) X-chromosome inactivation: a repeat hypothesis *Cytogenet Cell Genet* 80 (1-4): 133-7
- Lyon, M. F. (2000) LINE-1 elements and X chromosome inactivation: a function for "junk" DNA? *Proc Natl Acad Sci U S A* 97 (12): 6248-9
- Macaya, G., Thiery, J. P. and Bernardi, G. (1976) An approach to the organization of eukaryotic genomes at a macromolecular level *J Mol Biol* 108 (1): 237-54
- Magram, J., Chada, K. and Costantini, F. (1985) Developmental regulation of a cloned adult beta-globin gene in transgenic mice *Nature* 315 (6017): 338-40
- Malfray, L., Roth, M. P., Carrington, M., Borot, N., Volz, A., Ziegler, A. and Coppin, H. (1997) Heterogeneity in rates of recombination in the 6-Mb region telomeric to the human major histocompatibility complex *Genomics* 43 (2): 226-31
- Malnic, B., Hirono, J., Sato, T. and Buck, L. B. (1999) Combinatorial receptor codes for odors *Cell* 96 (5): 713-23
- Mardis, E. R. (1994) High-throughput detergent extraction of M13 subclones for fluorescent DNA sequencing *Nucleic Acids Res* 22 (11): 2173-5
- Matarazzo, V., Tirard, A., Renucci, M., Belaich, A. and Clement, J. L. (1998) Isolation of putative olfactory receptor sequences from pig nasal epithelium *Neurosci Lett* 249 (2-3): 87-90
- Matsuda, S., Kawamura-Tsuzuku, J., Ohsugi, M., Yoshida, M., Emi, M., Nakamura, Y., Onda, M., Yoshida, Y., Nishiyama, A. and Yamamoto, T. (1996) Tob, a novel protein that interacts with p185erbB2, is associated with anti-proliferative activity *Oncogene* 12 (4): 705-13
- Matsunami, H. and Buck, L. B. (1997) A multigene family encoding a diverse array of putative pheromone receptors in mammals *Cell* 90 (4): 775-84
- Maxam, A. M. and Gilbert, W. (1977) A new method for sequencing DNA *Proc Natl Acad Sci U S A* 74 (2): 560-4

- Maxson R., Cohn R., Kedes L. and Mohun T. (1983) Expression and organization of histone genes. *Annu Rev Genet.* 17:239-77.
- McClintock, M. K. (1971) Menstrual synchrony and suppression *Nature* 229 (5282): 244-5
- McClintock, T. S., Landers, T. M., Gimelbrant, A. A., Fuller, L. Z., Jackson, B. A., Jayawickreme, C. K. and Lerner, M. R. (1997) Functional expression of olfactory-adrenergic receptor chimeras and intracellular retention of heterologously expressed olfactory receptors *Brain Res Mol Brain Res* 48 (2): 270-8
- McPherson, J. D., Marra, M., Hillier, L., Waterston, R. H., Chinwalla, A., Wallis, J., Sekhon, M., Wylie, K., Mardis, E. R., Wilson, R. K., Fulton, R., Kucaba, T. A., Wagner-McPherson, C., Barbazuk, W. B., Gregory, S. G., Humphray, S. J., French, L., Evans, R. S., Bethel, G., Whittaker, A., Holden, J. L., McCann, O. T., Dunham, A., Soderlund, C., Scott, C. E., Bentley, D. R., Schuler, G., Chen, H. C., Jang, W., Green, E. D., Idol, J. R., Maduro, V. V., Montgomery, K. T., Lee, E., Miller, A., Emerling, S., Kucherlapati, Gibbs, R., Scherer, S., Gorrell, J. H., Sodergren, E., Clerc-Blankenburg, K., Tabor, P., Naylor, S., Garcia, D., de Jong, P. J., Catanese, J. J., Nowak, N., Osoegawa, K., Qin, S., Rowen, L., Madan, A., Dors, M., Hood, L., Trask, B., Friedman, C., Massa, H., Cheung, V. G., Kirsch, I. R., Reid, T., Yonescu, R., Weissenbach, J., Bruls, T., Heilig, R., Branscomb, E., Olsen, A., Doggett, N., Cheng, J. F., Hawkins, T., Myers, R. M., Shang, J., Ramirez, L., Schmutz, J., Velasquez, O., Dixon, K., Stone, N. E., Cox, D. R., Haussler, D., Kent, W. J., Furey, T., Rogic, S., Kennedy, S., Jones, S., Rosenthal, A., Wen, G., Schilhabel, M., Gloeckner, G., Nyakatura, G., Siebert, R., Schlegelberger, B., Korenberg, J., Chen, X. N., Fujiyama, A., Hattori, M., Toyoda, A., Yada, T., Park, H. S., Sakaki, Y., Shimizu, N., Asakawa, S., Kawasaki, K., Sasaki, T., Shintani, A., Shimizu, A., Shibuya, K., Kudoh, J., Minoshima, S., Ramser, J., Seranski, P., Hoff, C., Poustka, A., Reinhardt, R. and Lehrach, H. (2001) A physical map of the human genome *Nature* 409 (6822): 934-41
- Menco, B. P. and Jackson, J. E. (1997) A banded topography in the developing rat's olfactory epithelial surface *J Comp Neurol* 388 (2): 293-306
- Meng, E. C. and Bourne, H. R. (2001) Receptor activation: what does the rhodopsin structure tell us? *Trends Pharmacol Sci* 22 (11): 587-93
- Meredith, M. (1986) Vomeronasal organ removal before sexual experience impairs male hamster mating behavior *Physiol Behav* 36 (4): 737-43
- Meunier-Rotival, M., Soriano, P., Cuny, G., Strauss, F. and Bernardi, G. (1982) Sequence organization and genomic distribution of the major family of interspersed repeats of mouse DNA *Proc Natl Acad Sci U S A* 79 (2): 355-9
- Meyerhans, A., Vartanian, J. P. and Wain-Hobson, S. (1990) DNA recombination during PCR *Nucleic Acids Res* 18 (7): 1687-91
- Mironov, A. A., Fickett, J. W. and Gelfand, M. S. (1999) Frequent alternative splicing of human genes *Genome Res* 9 (12): 1288-93
- Moller, S., Croning, M. D. and Apweiler, R. (2001) Evaluation of methods for the prediction of membrane spanning regions *Bioinformatics* 17 (7): 646-53
- Mombaerts, P. (1996) Targeting olfaction *Curr Opin Neurobiol* 6 (4): 481-6

- Mombaerts, P. (1999) Molecular biology of odorant receptors in vertebrates *Annu Rev Neurosci* 22 487-509
- Mombaerts, P. (1999) Odorant receptor genes in humans *Curr Opin Genet Dev* 9 (3): 315-20
- Mombaerts, P., Wang, F., Dulac, C., Chao, S. K., Nemes, A., Mendelsohn, M., Edmondson, J. and Axel, R. (1996a) Visualizing an olfactory sensory map *Cell* 87 (4): 675-86
- Mombaerts, P., Wang, F., Dulac, C., Vassar, R., Chao, S. K., Nemes, A., Mendelsohn, M., Edmondson, J. and Axel, R. (1996b) The molecular biology of olfactory perception *Cold Spring Harb Symp Quant Biol* 61 135-45
- Mondadori, C., Jaekel, J. and Preiswerk, G. (1993) CGP 36742: the first orally active GABAB blocker improves the cognitive performance of mice, rats, and rhesus monkeys *Behav Neural Biol* 60 (1): 62-8
- Monnot, C., Weber, V., Stinnakre, J., Bihoreau, C., Teutsch, B., Corvol, P. and Clauser, E. (1991) Cloning and functional characterization of a novel mas-related gene, modulating intracellular angiotensin II actions *Mol Endocrinol* 5 (10): 1477-87
- Murrell, J. R. and Hunter, D. D. (1999) An olfactory sensory neuron line, odora, properly targets olfactory proteins and responds to odorants *J Neurosci* 19 (19): 8260-70
- Musahl, C., Schulte, D., Burkhart, R. and Knippers, R. (1995) A human homologue of the yeast replication protein Cdc21. Interactions with other Mcm proteins *Eur J Biochem* 230 (3): 1096-101
- Nagase, T., Ishikawa, K., Suyama, M., Kikuno, R., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N. and Ohara, O. (1998) Prediction of the coding sequences of unidentified human genes. XI. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro *DNA Res* 5 (5): 277-86
- Nahm, D. H., Lee, Y. E., Yim, E. J., Park, H. S., Yim, H., Kang, Y. and Kim, J. K. (2002) Identification of cytokeratin 18 as a bronchial epithelial autoantigen associated with nonallergic asthma *Am J Respir Crit Care Med* 165 (11): 1536-9
- Nakamura, T., Hillova, J., Mariage-Samson, R. and Hill, M. (1988) Molecular cloning of a novel oncogene generated by DNA recombination during transfection *Oncogene Res* 2 (4): 357-70
- Navarro, M. and Gull, K. (2001) A pol I transcriptional body associated with VSG mono-allelic expression in *Trypanosoma brucei* *Nature* 414 (6865): 759-63
- Nef, P., Hermans-Borgmeyer, I., Artieres-Pin, H., Beasley, L., Dionne, V. E. and Heinemann, S. F. (1992) Spatial pattern of receptor expression in the olfactory epithelium *Proc Natl Acad Sci U S A* 89 (19): 8948-52
- Nef, S. and Nef, P. (1997) Olfaction: transient expression of a putative odorant receptor in the avian notochord *Proc Natl Acad Sci U S A* 94 (9): 4766-71

- Nei, M. and Kumar, S. *Molecular Evolution and Phylogenetics* (Oxford University Press, New York, 2000).
- Neitzel, H., Kalscheuer, V., Henschel, S., Digweed, M. and Sperling, K. (1998) Beta-heterochromatin in mammals: evidence from studies in *Microtus agrestis* based on the extensive accumulation of L1 and non-L1 retroposons in the heterochromatin *Cytogenet Cell Genet* 80 (1-4): 165-72
- Ngai, J., Chess, A., Dowling, M. M., Necles, N., Macagno, E. R. and Axel, R. (1993a) Coding of olfactory information: topography of odorant receptor expression in the catfish olfactory epithelium *Cell* 72 (5): 667-80
- Ngai, J., Dowling, M. M., Buck, L., Axel, R. and Chess, A. (1993b) The family of genes encoding odorant receptors in the channel catfish *Cell* 72 (5): 657-66
- Nomura, N., Miyajima, N., Sazuka, T., Tanaka, A., Kawarabayasi, Y., Sato, S., Nagase, T., Seki, N., Ishikawa, K. and Tabata, S. (1994) Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1 (supplement) *DNA Res* 1 (1): 47-56
- O'Hara, P. J., Sheppard, P. O., Thogersen, H., Venezia, D., Haldeman, B. A., McGrane, V., Houamed, K. M., Thomsen, C., Gilbert, T. L. and Mulvihill, E. R. (1993) The ligand-binding domain in metabotropic glutamate receptors is related to bacterial periplasmic binding proteins *Neuron* 11 (1): 41-52
- Ober, C., Elias, S., Kostyu, D. D. and Hauck, W. W. (1992) Decreased fecundability in Hutterite couples sharing HLA-DR *Am J Hum Genet* 50 (1): 6-14
- Okazaki, S., Tanase, S., Choudhury, B. K., Setoyama, K., Miura, R., Ogawa, M. and Setoyama, C. (1994) A novel nuclear protein with zinc fingers down-regulated during early mammalian cell differentiation *J Biol Chem* 269 (9): 6900-7
- Olavesen, M. G., Bentley, E., Mason, R. V., Stephens, R. J. and Ragoussis, J. (1997) Fine mapping of 39 ESTs on human chromosome 6p23-p25 *Genomics* 46 (2): 303-6
- O'Leary, D. D., Yates, P. A. and McLaughlin, T. (1999) Molecular development of sensory maps: representing sights and smells in the brain *Cell* 96 (2): 255-69
- Ortmann, B., Androlewicz, M. J. and Cresswell, P. (1994) MHC class I/beta 2-microglobulin complexes associate with TAP transporters before peptide binding *Nature* 368 (6474): 864-7
- Palczewski, K., Kumasaka, T., Hori, T., Behnke, C. A., Motoshima, H., Fox, B. A., Le Trong, I., Teller, D. C., Okada, T., Stenkamp, R. E., Yamamoto, M. and Miyano, M. (2000) Crystal structure of rhodopsin: A G protein-coupled receptor *Science* 289 (5480): 739-45
- Panning, B. and Smiley, J. R. (1993) Activation of RNA polymerase III transcription of human Alu repetitive elements by adenovirus type 5: requirement for the E1b 58-kilodalton protein and the products of E4 open reading frames 3 and 6 *Mol Cell Biol* 13 (6): 3231-44

- Parmentier, M., Libert, F., Schurmans, S., Schiffmann, S., Lefort, A., Eggerickx, D., Ledent, C., Mollereau, C., Gerard, C., Perret, J. and et al. (1992) Expression of members of the putative olfactory receptor gene family in mammalian germ cells *Nature* 355 (6359): 453-5
- Pennesi, G., Brioli, G., Lulli, P., Mariani, B., Morellini, M., Nicotra, M. and Trabace, S. (1998) HLA and complement factors alleles sharing in Italian couples with recurrent spontaneous abortions *Hum Immunol* 59 (6): 382-6
- Peters, H. C., Kammer, G., Volz, A., Kaupmann, K., Ziegler, A., Bettler, B., Epplen, J. T., Sander, T. and Riess, O. (1998) Mapping, genomic structure, and polymorphisms of the human GABAR₁ receptor gene: evaluation of its involvement in idiopathic generalized epilepsy *Neurogenetics* 2 (1): 47-54
- Petruzzella, V., Tiranti, V., Fernandez, P., Ianna, P., Carrozzo, R. and Zeviani, M. (1998) Identification and characterization of human cDNAs specific to BCS1, PET112, SCO1, COX15, and COX11, five genes involved in the formation and function of the mitochondrial respiratory chain *Genomics* 54 (3): 494-504
- Pham-Dinh, D., Della Gaspera, B., Kerlero de Rosbo, N. and Dautigny, A. (1995) Structure of the human myelin/oligodendrocyte glycoprotein gene and multiple alternative spliced isoforms *Genomics* 29 (2): 345-52
- Pham-Dinh, D., Mattei, M. G., Nussbaum, J. L., Roussel, G., Pontarotti, P., Roeckel, N., Mather, I. H., Artzt, K., Lindahl, K. F. and Dautigny, A. (1993) Myelin/oligodendrocyte glycoprotein is a member of a subset of the immunoglobulin superfamily encoded within the major histocompatibility complex *Proc Natl Acad Sci U S A* 90 (17): 7990-4
- Pilpel, Y. and Lancet, D. (1999) The variable and conserved interfaces of modeled olfactory receptor proteins *Protein Sci* 8 (5): 969-77
- Porteros, A., Brinon, J. G., Crespo, C., Okazaki, K., Hidaka, H., Aijon, J. and Alonso, J. R. (1996) Neurocalcin immunoreactivity in the rat accessory olfactory bulb *Brain Res* 729 (1): 82-9
- Potts, W. K. and Wakeland, E. K. (1990) The maintenance of MHC polymorphism *Immunol Today* 11 (2): 39-40
- Potts, W. K. and Wakeland, E. K. (1993) Evolution of MHC genetic diversity: a tale of incest, pestilence and sexual preference *Trends Genet* 9 (12): 408-12
- Prober, J. M., Trainor, G. L., Dam, R. J., Hobbs, F. W., Robertson, C. W., Zagursky, R. J., Cocuzza, A. J., Jensen, M. A. and Baumeister, K. (1987) A system for rapid DNA sequencing with fluorescent chain-terminating dideoxynucleotides *Science* 238 (4825): 336-41
- Qasba, P. and Reed, R. R. (1998) Tissue and zonal-specific expression of an olfactory receptor transgene *J Neurosci* 18 (1): 227-36
- Quandt, K., Frech, K., Karas, H., Wingender, E. and Werner, T. (1995) MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data *Nucleic Acids Res* 23 (23): 4878-84

- Raasi, S., Schmidtke, G. and Groettrup, M. (2001) The ubiquitin-like protein FAT10 forms covalent conjugates and induces apoptosis *J Biol Chem* 276 (38): 35334-43
- Raming, K., Konzelmann, S. and Breer, H. (1998) Identification of a novel G-protein coupled receptor expressed in distinct brain regions and a defined olfactory zone *Receptors Channels* 6 (2): 141-51
- Raming, K., Krieger, J., Strotmann, J., Boekhoff, I., Kubick, S., Baumstark, C. and Breer, H. (1993) Cloning and expression of odorant receptors *Nature* 361 (6410): 353-6
- Rana, B. K., Shiina, T. and Insel, P. A. (2001) Genetic variations and polymorphisms of G protein-coupled receptors: functional and therapeutic implications *Annu Rev Pharmacol Toxicol* 41 593-624
- Ressler, K. J., Sullivan, S. L. and Buck, L. B. (1993) A zonal organization of odorant receptor gene expression in the olfactory epithelium *Cell* 73 (3): 597-609
- Ressler, K. J., Sullivan, S. L. and Buck, L. B. (1994) Information coding in the olfactory system: evidence for a stereotyped and highly organized epitope map in the olfactory bulb *Cell* 79 (7): 1245-55
- Reynolds, J. and Keverne, E. B. (1979) The accessory olfactory system and its role in the pheromonally mediated suppression of oestrus in grouped mice *J Reprod Fertil* 57 (1): 31-5
- Rhodes, D. A., Stammers, M., Malcherek, G., Beck, S. and Trowsdale, J. (2001) The cluster of BTN genes in the extended major histocompatibility complex *Genomics* 71 (3): 351-62
- Rice, P., Longden, I. and Bleasby, A. (2000) EMBOSS: the European Molecular Biology Open Software Suite *Trends Genet* 16 (6): 276-7
- Richards, S., Bucan, M., Brorson, K., Kiefer, M. C., Hunt, S. W., 3rd, Lehrach, H. and Lindahl, K. F. (1989) Genetic and molecular mapping of the Hmt region of mouse *Embo J* 8 (12): 3749-57
- Roetto, A., Sbaiz, L., Bosio, S., Piperno, A., Fargion, S., Carella, M., Totaro, A., Grifa, A., Gasparini, P. and Camaschella, C. (1997) A recombination event close to HFE gene in hereditary hemochromatosis *Ann Genet* 40 (3): 150-3
- Rogers, J. and Bradley, A. (2001) The mouse genome sequence: status and prospects *Genomics* 77 (3): 117-8
- Roth, M. P., Malfroy, L., Offer, C., Sevin, J., Enault, G., Borot, N., Pontarotti, P. and Coppin, H. (1995) The human myelin oligodendrocyte glycoprotein (MOG) gene: complete nucleotide sequence and structural characterization *Genomics* 28 (2): 241-50
- Rouquier, S., Taviaux, S., Trask, B. J., Brand-Arpon, V., van den Engh, G., Demaille, J. and Giorgi, D. (1998) Distribution of olfactory receptor genes in the human genome *Nat Genet* 18 (3): 243-50
- Rubin, C. M., VandeVoort, C. A., Teplitz, R. L. and Schmid, C. W. (1994) Alu repeated DNAs are differentially methylated in primate germ cells *Nucleic Acids Res* 22 (23): 5121-7

- Ruddy, D. A., Kronmal, G. S., Lee, V. K., Mintier, G. A., Quintana, L., Domingo, R., Jr., Meyer, N. C., Irrinki, A., McClelland, E. E., Fullan, A., Mapa, F. A., Moore, T., Thomas, W., Loeb, D. B., Harmon, C., Tsuchihashi, Z., Wolff, R. K., Schatzman, R. C. and Feder, J. N. (1997) A 1.1-Mb transcript map of the hereditary hemochromatosis locus *Genome Res* 7 (5): 441-56
- Ryba, N. J. and Tirindelli, R. (1997) A new multigene family of putative pheromone receptors *Neuron* 19 (2): 371-9
- Sachidanandam, R., Weissman, D., Schmidt, S. C., Kakol, J. M., Stein, L. D., Marth, G., Sherry, S., Mullikin, J. C., Mortimore, B. J., Willey, D. L., Hunt, S. E., Cole, C. G., Coggill, P. C., Rice, C. M., Ning, Z., Rogers, J., Bentley, D. R., Kwok, P. Y., Mardis, E. R., Yeh, R. T., Schultz, B., Cook, L., Davenport, R., Dante, M., Fulton, L., Hillier, L., Waterston, R. H., McPherson, J. D., Gilman, B., Schaffner, S., Van Etten, W. J., Reich, D., Higgins, J., Daly, M. J., Blumenstiel, B., Baldwin, J., Stange-Thomann, N., Zody, M. C., Linton, L., Lander, E. S. and Altshuler, D. (2001) A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms *Nature* 409 (6822): 928-33
- Saitou, N. and Nei, M. (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees *Mol Biol Evol* 4 (4): 406-25
- Sakmar, T. P. (2002) Structure of rhodopsin and the superfamily of seven-helical receptors: the same and not the same *Curr Opin Cell Biol* 14 (2): 189-95
- Sanger, F., Nicklen, S. and Coulson, A. R. (1977) DNA sequencing with chain-terminating inhibitors *Proc Natl Acad Sci U S A* 74 (12): 5463-7
- Satoh, T., Tsuruga, H., Yabuta, N., Ishidate, M., Jr. and Nojima, H. (1997) Assignment of the human CDC21 (MCM4) gene to chromosome 8q11.2 *Genomics* 46 (3): 525-6
- Scherf, M., Klingenhoff, A. and Werner, T. (2000) Highly specific localization of promoter regions in large genomic sequences by PromoterInspector: a novel context analysis approach *J Mol Biol* 297 (3): 599-606
- Schleicher, S., Boekhoff, I., Arriza, J., Lefkowitz, R. J. and Breer, H. (1993) A beta-adrenergic receptor kinase-like enzyme is involved in olfactory signal termination *Proc Natl Acad Sci U S A* 90 (4): 1420-4
- Schmid, C. W. (1998) Does SINE evolution preclude Alu function? *Nucleic Acids Res* 26 (20): 4541-50
- Scholler, N., Hayden-Ledbetter, M., Hellstrom, K. E., Hellstrom, I. and Ledbetter, J. A. (2001) CD83 is a sialic acid-binding Ig-like lectin (Siglec) adhesion receptor that binds monocytes and a subset of activated CD8+ T cells *J Immunol* 166 (6): 3865-72
- Schurmans, S., Muscatelli, F., Miot, F., Mattei, M. G., Vassart, G. and Parmentier, M. (1993) The OLFR1 gene encoding the HGMP07E putative olfactory receptor maps to the 17p13-->p12 region of the human genome and reveals an MspI restriction fragment length polymorphism *Cytogenet Cell Genet* 63 (3): 200-4

- Schwartz, S., Zhang, Z., Frazer, K. A., Smit, A., Riemer, C., Bouck, J., Gibbs, R., Hardison, R. and Miller, W. (2000) PipMaker--a web server for aligning two genomic DNA sequences *Genome Res* 10 (4): 577-86
- Schwarz, D. A., Barry, G., Eliasof, S. D., Petroski, R. E., Conlon, P. J. and Maki, R. A. (2000) Characterization of gamma -aminobutyric acid receptor GABAB(1e), a GABAB(1) splice variant encoding a truncated receptor *J Biol Chem* 275 (41): 32174-81
- Seki, N., Ohira, M., Nagase, T., Ishikawa, K., Miyajima, N., Nakajima, D., Nomura, N. and Ohara, O. (1997) Characterization of cDNA clones in size-fractionated cDNA libraries from human brain *DNA Res* 4 (5): 345-9
- Selbie, L. A., Townsend-Nicholson, A., Iismaa, T. P. and Shine, J. (1992) Novel G protein-coupled receptors: a gene family of putative human olfactory receptor sequences *Brain Res Mol Brain Res* 13 (1-2): 159-63
- Senger, G., Ragoussis, J., Trowsdale, J., and Sheer, D. (1993) Fine mapping of the human MHC class II region within chromosome band 6p21 and evaluation of probe ordering using interphase fluorescence in situ hybridization *Cytogenet Cell Genet.* 64 (1):49-53
- Sharon, D., Gilad, Y., Glusman, G., Khen, M., Lancet, D. and Kalush, F. (2000) Identification and characterization of coding single-nucleotide polymorphisms within a human olfactory receptor gene cluster *Gene* 260 (1-2): 87-94
- Sharon, D., Glusman, G., Pilpel, Y., Horn-Saban, S. and Lancet, D. (1998) Genome dynamics, evolution, and protein modeling in the olfactory receptor gene superfamily *Ann NY Acad Sci* 855 182-93
- Sharon, D., Glusman, G., Pilpel, Y., Khen, M., Gruetzner, F., Haaf, T., and Lancet, D. (1999) Primate evolution of an olfactory receptor cluster: diversification by gene conversion and recent emergence of pseudogenes *Genomics* 61 (1): 24-36
- Shiina, T., Tamiya, G., Oka, A., Takishima, N., Yamagata, T., Kikkawa, E., Iwata, K., Tomizawa, M., Okuaki, N., Kuwano, Y., Watanabe, K., Fukuzumi, Y., Itakura, S., Sugawara, C., Ono, A., Yamazaki, M., Tashiro, H., Ando, A., Ikemura, T., Soeda, E., Kimura, M., Bahram, S. and Inoko, H. (1999) Molecular dynamics of MHC genesis unraveled by sequence analysis of the 1,796,938-bp HLA class I region *Proc Natl Acad Sci U S A* 96 (23): 13282-7
- Simon, M., Le Mignon, L., Fauchet, R., Yaouanq, J., David, V., Edan, G. and Bourel, M. (1987) A study of 609 HLA haplotypes marking for the hemochromatosis gene: (1) mapping of the gene near the HLA-A locus and characters required to define a heterozygous population and (2) hypothesis concerning the underlying cause of hemochromatosis-HLA association *Am J Hum Genet* 41 (2): 89-105
- Singer, M. S., Hughes, T. E., Shepherd, G. M. and Greer, C. A. (1998) Identification of olfactory receptor mRNA sequences from the rat olfactory bulb glomerular layer *Neuroreport* 9 (16): 3745-8
- Sinnarajah, S., Dessauer, C. W., Sri Kumar, D., Chen, J., Yuen, J., Yilma, S., Dennis, J. C., Morrison, E. E., Vodyanoy, V. and Kehrl, J. H. (2001) RGS2 regulates signal transduction in olfactory neurons by attenuating activation of adenylyl cyclase III *Nature* 409 (6823): 1051-5

- Slade, R. W. and McCallum, H. I. (1992) Overdominant vs. frequency-dependent selection at MHC loci *Genetics* 132 (3): 861-4
- Sloan, V. S., Cameron, P., Porter, G., Gammon, M., Amaya, M., Mellins, E. and Zaller, D. M. (1995) Mediation by HLA-DM of dissociation of peptides from HLA-DR *Nature* 375 (6534): 802-6
- Sly, W. S., Quinton, B. A., McAlister, W. H. and Rimoin, D. L. (1973) Beta glucuronidase deficiency: report of clinical, radiologic, and biochemical features of a new mucopolysaccharidosis *J Pediatr* 82 (2): 249-57
- Smaglik, P. and Abbott, A. (2000) Project offers free mouse sequence *Nature* 407 (6805): 663-4
- Smit, A. F. (1996) The origin of interspersed repeats in the human genome *Curr Opin Genet Dev* 6 (6): 743-8
- Smit, A. F. (1999) Interspersed repeats and other mementos of transposable elements in mammalian genomes *Curr Opin Genet Dev* 9 (6): 657-63
- Smith, L. M., Sanders, J. Z., Kaiser, R. J., Hughes, P., Dodd, C., Connell, C. R., Heiner, C., Kent, S. B. and Hood, L. E. (1986) Fluorescence detection in automated DNA sequence analysis *Nature* 321 (6071): 674-9
- Smith, T. F. and Waterman, M. S. (1981) Identification of common molecular subsequences *J Mol Biol* 147 (1): 195-7
- Snyder, S. H., Sklar, P. B. and Pevsner, J. (1988) Molecular mechanisms of olfaction *J Biol Chem* 263 (28): 13971-4
- Soderlund, C., Humphray, S., Dunham, A. and French, L. (2000) Contigs built with fingerprints, markers, and FPC V4.7 *Genome Res* 10 (11): 1772-87
- Soderlund, C., Longden, I. and Mott, R. (1997) FPC: a system for building contigs from restriction fingerprinted clones *Comput Appl Biosci* 13 (5): 523-35
- Solovyev, V. and Salamov, A. (1997) The Gene-Finder computer tools for analysis of human and model organisms genome sequences *Proc Int Conf Intell Syst Mol Biol* 5 294-302
- Sonnhammer, E. L. and Durbin, R. (1995) A dot-matrix program with dynamic threshold control suited for genomic DNA and protein sequence analysis *Gene* 167 (1-2): GC1-10
- Soriano, P., Macaya, G. and Bernardi, G. (1981) The major components of the mouse and human genomes. 2. Reassociation kinetics *Eur J Biochem* 115 (2): 235-9
- Soriano, P., Meunier-Rotival, M. and Bernardi, G. (1983) The distribution of interspersed repeats is nonuniform and conserved in the mouse and human genomes *Proc Natl Acad Sci U S A* 80 (7): 1816-20
- Sosinsky, A., Glusman, G. and Lancet, D. (2000) The genomic structure of human olfactory receptor genes *Genomics* 70 (1): 49-61

- Speleman, F., Vervoort, R., van Roy, N., Liebaers, I., Sly, W. S. and Lissens, W. (1996) Localization by fluorescence in situ hybridization of the human functional beta-glucuronidase gene (GUSB) to 7q11.21 --> q11.22 and two pseudogenes to 5p13 and 5q13 *Cytogenet Cell Genet* 72 (1): 53-5
- Staden, R., Beal, K. F. and Bonfield, J. K. (2000) The Staden package, 1998 *Methods Mol Biol* 132 115-30
- Staden, R., Judge, D. P. and Bonfield, J. K. (2001) Sequence assembly and finishing methods *Methods Biochem Anal* 43 303-22
- Stephens, R., Horton, R., Humphray, S., Rowen, L., Trowsdale, J. and Beck, S. (1999) Gene organisation, sequence variation and isochore structure at the centromeric boundary of the human MHC *J Mol Biol* 291 (4): 789-99
- Stoessner, G., Baker, W., van den Broek, A., Camon, E., Garcia-Pastor, M., Kanz, C., Kulikova, T., Leinonen, R., Lin, Q., Lombard, V., Lopez, R., Redaschi, N., Stoehr, P., Tuli, M. A., Tzouvara, K. and Vaughan, R. (2002) The EMBL Nucleotide Sequence Database *Nucleic Acids Res* 30 (1): 21-6
- Strotmann, J., Beck, A., Kubick, S. and Breer, H. (1995) Topographic patterns of odorant receptor expression in mammals: a comparative study *J Comp Physiol [A]* 177 (6): 659-66
- Strotmann, J., Wanner, I., Helfrich, T., Beck, A. and Breer, H. (1994a) Rostro-caudal patterning of receptor-expressing olfactory neurones in the rat nasal cavity *Cell Tissue Res* 278 (1): 11-20
- Strotmann, J., Wanner, I., Helfrich, T., Beck, A., Meinken, C., Kubick, S. and Breer, H. (1994b) Olfactory neurones expressing distinct odorant receptor subtypes are spatially segregated in the nasal neuroepithelium *Cell Tissue Res* 276 (3): 429-38
- Strotmann, J., Wanner, I., Krieger, J., Raming, K. and Breer, H. (1992) Expression of odorant receptors in spatially restricted subsets of chemosensory neurones *Neuroreport* 3 (12): 1053-6
- Sullivan, S. L., Adamson, M. C., Ressler, K. J., Kozak, C. A. and Buck, L. B. (1996) The chromosomal distribution of mouse odorant receptor genes *Proc Natl Acad Sci U S A* 93 (2): 884-8
- Sullivan, S. L., Bohm, S., Ressler, K. J., Horowitz, L. F. and Buck, L. B. (1995) Target-independent pattern specification in the olfactory epithelium *Neuron* 15 (4): 779-89
- Sulston, J., Mallett, F., Staden, R., Durbin, R., Horsnell, T. and Coulson, A. (1988) Software for genome mapping by fingerprinting techniques *Comput Appl Biosci* 4 (1): 125-32
- Szpirer, C., Szpirer, J., Riviere, M., Tazi, R. and Pontarotti, P. (1997) Mapping of the Olf89 and Rfp genes to the rat genome: comparison with the mouse and human and new insights into the evolution of the rodent genome *Cytogenet Cell Genet* 78 (2): 137-9
- Takahashi, K., Tsuchida, K., Tanabe, Y., Masu, M. and Nakanishi, S. (1993) Role of the large extracellular domain of metabotropic glutamate receptors in agonist selectivity determination *J Biol Chem* 268 (26): 19341-5

Takahata, N. and Nei, M. (1990) Allelic genealogy under overdominant and frequency-dependent selection and polymorphism of major histocompatibility complex loci *Genetics* 124 (4): 967-78

Tang, M. X., Redemann, C. T. and Szoka, F. C., Jr. (1996) In vitro gene delivery by degraded polyamidoamine dendrimers *Bioconjug Chem* 7 (6): 703-14

Tegoni, M., Pelosi, P., Vincent, F., Spinelli, S., Campanacci, V., Grolli, S., Ramoni, R. and Cambillau, C. (2000) Mammalian odorant binding proteins *Biochim Biophys Acta* 1482 (1-2): 229-40

The International Human Genome Sequencing Consortium [IHGSC]: Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., Funke, R., Gage, D., Harris, K., Heaford, A., Howland, J., Kann, L., Lehoczky, J., LeVine, R., McEwan, P., McKernan, K., Meldrim, J., Mesirov, J. P., Miranda, C., Morris, W., Naylor, J., Raymond, C., Rosetti, M., Santos, R., Sheridan, A., Sougnez, C., Stange-Thomann, N., Stojanovic, N., Subramanian, A., Wyman, D., Rogers, J., Sulston, J., Ainscough, R., Beck, S., Bentley, D., Burton, J., Clee, C., Carter, N., Coulson, A., Deadman, R., Deloukas, P., Dunham, A., Dunham, I., Durbin, R., French, L., Grafham, D., Gregory, S., Hubbard, T., Humphray, S., Hunt, A., Jones, M., Lloyd, C., McMurray, A., Matthews, L., Mercer, S., Milne, S., Mullikin, J. C., Mungall, A., Plumb, R., Ross, M., Shownkeen, R., Sims, S., Waterston, R. H., Wilson, R. K., Hillier, L. W., McPherson, J. D., Marra, M. A., Mardis, E. R., Fulton, L. A., Chinwalla, A. T., Pepin, K. H., Gish, W. R., Chissoe, S. L., Wendl, M. C., Delehaunty, K. D., Miner, T. L., Delehaunty, A., Kramer, J. B., Cook, L. L., Fulton, R. S., Johnson, D. L., Minx, P. J., Clifton, S. W., Hawkins, T., Branscomb, E., Predki, P., Richardson, P., Wenning, S., Slezak, T., Doggett, N., Cheng, J. F., Olsen, A., Lucas, S., Elkin, C., Uberbacher, E., Frazier, M., Gibbs, R. A., Muzny, D. M., Scherer, S. E., Bouck, J. B., Sodergren, E. J., Worley, K. C., Rives, C. M., Gorrell, J. H., Metzker, M. L., Naylor, S. L., Kucherlapati, R. S., Nelson, D. L., Weinstock, G. M., Sakaki, Y., Fujiyama, A., Hattori, M., Yada, T., Toyoda, A., Itoh, T., Kawagoe, C., Watanabe, H., Totoki, Y., Taylor, T., Weissenbach, J., Heilig, R., Saurin, W., Artiguenave, F., Brottier, P., Bruls, T., Pelletier, E., Robert, C., Wincker, P., Smith, D. R., Doucette-Stamm, L., Rubenfield, M., Weinstock, K., Lee, H. M., Dubois, J., Rosenthal, A., Platzer, M., Nyakatura, G., Taudien, S., Rump, A., Yang, H., Yu, J., Wang, J., Huang, G., Gu, J., Hood, L., Rowen, L., Madan, A., Qin, S., Davis, R. W., Federspiel, N. A., Abola, A. P., Proctor, M. J., Myers, R. M., Schmutz, J., Dickson, M., Grimwood, J., Cox, D. R., Olson, M. V., Kaul, R., Shimizu, N., Kawasaki, K., Minoshima, S., Evans, G. A., Athanasiou, M., Schultz, R., Roe, B. A., Chen, F., Pan, H., Ramser, J., Lehrach, H., Reinhardt, R., McCombie, W. R., de la Bastide, M., Dedhia, N., Blocker, H., Hornischer, K., Nordsiek, G., Agarwala, R., Aravind, L., Bailey, J. A., Bateman, A., Batzoglou, S., Birney, E., Bork, P., Brown, D. G., Burge, C. B., Cerutti, L., Chen, H. C., Church, D., Clamp, M., Copley, R. R., Doerks, T., Eddy, S. R., Eichler, E. E., Furey, T. S., Galagan, J., Gilbert, J. G., Harmon, C., Hayashizaki, Y., Haussler, D., Hermjakob, H., Hokamp, K., Jang, W., Johnson, L. S., Jones, T. A., Kasif, S., Kaspryzk, A., Kennedy, S., Kent, W. J., Kitts, P., Koonin, E. V., Korf, I., Kulp, D., Lancet, D., Lowe, T. M., McLysaght, A., Mikkelsen, T., Moran, J. V., Mulder, N., Pollara, V. J., Ponting, C. P., Schuler, G., Schultz, J., Slater, G., Smit, A. F., Stupka, E., Szustakowski, J., Thierry-Mieg, D., Thierry-Mieg, J., Wagner, L., Wallis, J., Wheeler, R., Williams, A., Wolf, Y. I., Wolfe, K. H., Yang, S. P., Yeh, R. F., Collins, F., Guyer, M. S., Peterson, J., Felsenfeld, A., Wetterstrand, K. A., Patrinos, A., Morgan, M. J. and Szustakowski, J. (2001) Initial sequencing and analysis of the human genome *Nature* 409 (6822): 860-921

- The MHC sequencing consortium (1999) Complete sequence and gene map of a human major histocompatibility complex. *Nature* 401 (6756): 921-3
- Theodosiou, A. M., Morrison, K. E., Nesbit, A. M., Daniels, R. J., Campbell, L., Francis, M. J., Christodoulou, Z. and Davies, K. E. (1994) Complex repetitive arrangements of gene sequence in the candidate region of the spinal muscular atrophy gene in 5q13. *Am J Hum Genet* 55 (6): 1209-17
- Thomas, M. B., Haines, S. L. and Akeson, R. A. (1996) Chemoreceptors expressed in taste, olfactory and male reproductive tissues. *Gene* 178 (1-2): 1-5
- Thompson, J. D., Higgins, D. G. and Gibson, T. J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22 (22): 4673-80
- Thomson, G. (1995) HLA disease associations: models for the study of complex human genetic disorders. *Crit Rev Clin Lab Sci* 32 (2): 183-219
- Tiwari, J. L. and Terasaki, P. I. *HLA and Disease Associations* (Springer-Verlag, New York, 1985).
- Tohgo, A., Takasawa, S., Munakata, H., Yonekura, H., Hayashi, N. and Okamoto, H. (1994) Structural determination and characterization of a 40 kDa protein isolated from rat 40 S ribosomal subunit. *FEBS Lett* 340 (1-2): 133-8
- Trappe, R., Doenecke, D. and Albig, W. (1999) The expression of human H2A-H2B histone gene pairs is regulated by multiple sequence elements in their joint promoters. *Biochim Biophys Acta* 1446 (3): 341-51
- Trask, B. J., Friedman, C., Martin-Gallardo, A., Rowen, L., Akinbami, C., Blankenship, J., Collins, C., Giorgi, D., Iadonato, S., Johnson, F., Kuo, W. L., Massa, H., Morrish, T., Naylor, S., Nguyen, O. T., Rouquier, S., Smith, T., Wong, D. J., Youngblom, J. and van den Engh, G. (1998) Members of the olfactory receptor gene family are contained in large blocks of DNA duplicated polymorphically near the ends of human chromosomes. *Hum Mol Genet* 7 (1): 13-26
- Trask, B. J., Massa, H., Brand-Arpon, V., Chan, K., Friedman, C., Nguyen, O. T., Eichler, E., van den Engh, G., Rouquier, S., Shizuya, H. and Giorgi, D. (1998) Large multi-chromosomal duplications encompass many members of the olfactory receptor gene family in the human genome. *Hum Mol Genet* 7 (13): 2007-20
- Twist, C. J., Beier, D. R., Disteche, C. M., Edelhoff, S. and Tedder, T. F. (1998) The mouse Cd83 gene: structure, domain organization, and chromosome localization. *Immunogenetics* 48 (6): 383-93
- Tzachanis, D., Freeman, G. J., Hirano, N., van Puijenbroek, A. A., Delfs, M. W., Berezovskaya, A., Nadler, L. M. and Boussiotis, V. A. (2001) Tob is a negative regulator of activation that is expressed in anergic and quiescent T cells. *Nat Immunol* 2 (12): 1174-82
- Uberbacher, E. C. and Mural, R. J. (1991) Locating protein-coding regions in human DNA sequences by a multiple sensor-neural network approach. *Proc Natl Acad Sci U S A* 88 (24): 11261-5

- Vanderhaeghen, P., Schurmans, S., Vassart, G. and Parmentier, M. (1993) Olfactory receptors are displayed on dog mature sperm cells *J Cell Biol* 123 (6 Pt 1): 1441-52
- Vanderhaeghen, P., Schurmans, S., Vassart, G. and Parmentier, M. (1997a) Molecular cloning and chromosomal mapping of olfactory receptor genes expressed in the male germ line: evidence for their wide distribution in the human genome *Biochem Biophys Res Commun* 237 (2): 283-7
- Vanderhaeghen, P., Schurmans, S., Vassart, G. and Parmentier, M. (1997b) Specific repertoire of olfactory receptor genes in the male germ cells of several mammalian species *Genomics* 39 (3): 239-46
- Vassar, R., Chao, S. K., Sitcheran, R., Nunez, J. M., Vosshall, L. B. and Axel, R. (1994) Topographic organization of sensory projections to the olfactory bulb *Cell* 79 (6): 981-91
- Vassar, R., Ngai, J. and Axel, R. (1993) Spatial segregation of odorant receptor expression in the mammalian olfactory epithelium *Cell* 74 (2): 309-18
- Velten, F., Rogel-Gaillard, C., Renard, C., Pontarotti, P., Tazi-Ahnini, R., Vaiman, M. and Chardon, P. (1998) A first map of the porcine major histocompatibility complex class I region *Tissue Antigens* 51 (2): 183-94
- Vernet, P., Faure, J., Dufaure, J. P. and Drevet, J. R. (1997) Tissue and developmental distribution, dependence upon testicular factors and attachment to spermatozoa of GPX5, a murine epididymis-specific glutathione peroxidase *Mol Reprod Dev* 47 (1): 87-98
- Vogt, R. G., Lindsay, S. M., Byrd, C. A. and Sun, M. (1997) Spatial patterns of olfactory neurons expressing specific odor receptor genes in 48-hour-old embryos of zebrafish *Danio rerio J Exp Biol* 200 (Pt 3) 433-43
- Volz, A., Ehlers, A., Younger, R. M., Forbes, S., Trowsdale, J., Beck, S. and Ziegler, A. (unpublished) Complex transcriptional control of MHC-linked olfactory receptor genes includes long distance and extensive alternative splicing, exon sharing and premature polyadenylation
- Volz, A. and Ziegler, A. (1996) Physical mapping of a 6 Mbp region directly telomeric of the HLA-complex *DNA Seq* 7 (1): 61-2
- Volz, A., Fonatsch, C. and Ziegler, A. (1992) Regional mapping of the gene for autosomal dominant spinocerebellar ataxia (SCA1) by localizing the closely linked D6S89 locus to 6p24.2--p23.05 *Cytogenet Cell Genet* 60 (1): 37-9
- Walensky, L. D., Roskams, A. J., Lefkowitz, R. J., Snyder, S. H. and Ronnett, G. V. (1995) Odorant receptors and desensitization proteins colocalize in mammalian sperm *Mol Med* 1 (2): 130-41
- Walensky, L. D., Ruat, M., Bakin, R. E., Blackshaw, S., Ronnett, G. V. and Snyder, S. H. (1998) Two novel odorant receptor families expressed in spermatids undergo 5'-splicing *J Biol Chem* 273 (16): 9378-87
- Wang, C. R., Loveland, B. E. and Lindahl, K. F. (1991) H-2M3 encodes the MHC class I molecule presenting the maternally transmitted antigen of the mouse *Cell* 66 (2): 335-45

- Wang, D. G., Fan, J. B., Siao, C. J., Berno, A., Young, P., Sapolisky, R., Ghandour, G., Perkins, N., Winchester, E., Spencer, J., Kruglyak, L., Stein, L., Hsie, L., Topaloglou, T., Hubbell, E., Robinson, E., Mittmann, M., Morris, M. S., Shen, N., Kilburn, D., Rioux, J., Nusbaum, C., Rozen, S., Hudson, T. J., Lander, E. S. and et al. (1998) Large-scale identification, mapping, and genotyping of single-nucleotide polymorphisms in the human genome *Science* 280 (5366): 1077-82
- Wang, F., Nemes, A., Mendelsohn, M. and Axel, R. (1998) Odorant receptors govern the formation of a precise topographic map *Cell* 93 (1): 47-60
- Wang, M. M. and Reed, R. R. (1993) Molecular cloning of the olfactory neuronal transcription factor Olf-1 by genetic selection in yeast *Nature* 364 (6433): 121-6
- Warner, J. R. and Nierras, C. R. (1998) Trapping human ribosomal protein genes *Genome Res* 8 (5): 419-21
- Weber, D. A., Evavold, B. D. and Jensen, P. E. (1996) Enhanced dissociation of HLA-DR-bound peptides in the presence of HLA-DM *Science* 274 (5287): 618-20
- Weighardt, F., Biamonti, G. and Riva, S. (1995) Nucleo-cytoplasmic distribution of human hnRNP proteins: a search for the targeting domains in hnRNP A1 *J Cell Sci* 108 (Pt 2) 545-55
- Wellerdieck, C., Oles, M., Pott, L., Korschning, S., Gisselmann, G. and Hatt, H. (1997) Functional expression of odorant receptors of the zebrafish Danio rerio and of the nematode *C. elegans* in HEK293 cells *Chem Senses* 22 (4): 467-76
- Weth, F., Nadler, W. and Korschning, S. (1996) Nested expression domains for odorant receptors in zebrafish olfactory epithelium *Proc Natl Acad Sci U S A* 93 (23): 13321-6
- Wetzel, C. H., Oles, M., Wellerdieck, C., Kuczkowiak, M., Gisselmann, G. and Hatt, H. (1999) Specificity and sensitivity of a human olfactory receptor functionally expressed in human embryonic kidney 293 cells and *Xenopus Laevis* oocytes *J Neurosci* 19 (17): 7426-33
- Wingender, E., Chen, X., Hehl, R., Karas, H., Liebich, I., Matys, V., Meinhardt, T., Pruss, M., Reuter, I. and Schacherer, F. (2000) TRANSFAC: an integrated system for gene expression regulation *Nucleic Acids Res* 28 (1): 316-9
- Wolfsberg, T. G. and Landsman, D. (1997) A comparison of expressed sequence tags (ESTs) to human genomic sequences *Nucleic Acids Res* 25 (8): 1626-32
- Womble, D. D. (2000) GCG: The Wisconsin Package of sequence analysis programs *Methods Mol Biol* 132 3-22
- Wong, S. T., Trinh, K., Hacker, B., Chan, G. C., Lowe, G., Gaggar, A., Xia, Z., Gold, G. H. and Storm, D. R. (2000) Disruption of the type III adenylyl cyclase gene leads to peripheral and behavioral anosmia in transgenic mice *Neuron* 27 (3): 487-97
- Wysocki, C. J. and Beauchamp, G. K. (1984) Ability to smell androstenone is genetically determined *Proc Natl Acad Sci U S A* 81 (15): 4899-902

- Wysocki, C. J., Dorries, K. M. and Beauchamp, G. K. (1989) Ability to perceive androstenone can be acquired by ostensibly anosmic people *Proc Natl Acad Sci U S A* 86 (20): 7976-8
- Xu, Y., Mural, R., Shah, M. and Uberbacher, E. (1994) Recognizing exons in genomic sequence using GRAIL II *Genet Eng (N Y)* 16 241-53
- Yamagata, K., Goto, K., Kuo, C. H., Kondo, H. and Miki, N. (1990) Visinin: a novel calcium binding protein expressed in retinal cone cells *Neuron* 4 (3): 469-76
- Yamaguchi, M., Yamazaki, K., Beauchamp, G. K., Bard, J., Thomas, L. and Boyse, E. A. (1981) Distinctive urinary odors governed by the major histocompatibility locus of the mouse *Proc Natl Acad Sci U S A* 78 (9): 5817-20
- Yamazaki, K., Beauchamp, G. K., Kupniewski, D., Bard, J., Thomas, L. and Boyse, E. A. (1988) Familial imprinting determines H-2 selective mating preferences *Science* 240 (4857): 1331-2
- Yamazaki, K., Beauchamp, G. K., Wysocki, C. J., Bard, J., Thomas, L. and Boyse, E. A. (1983) Recognition of H-2 types in relation to the blocking of pregnancy in mice *Science* 221 (4606): 186-8
- Yamazaki, K., Boyse, E. A., Mike, V., Thaler, H. T., Mathieson, B. J., Abbott, J., Boyse, J., Zayas, Z. A. and Thomas, L. (1976) Control of mating preferences in mice by genes in the major histocompatibility complex *J Exp Med* 144 (5): 1324-35
- Yoon, S. J., LeBlanc-Straceski, J., Ward, D., Krauter, K. and Kucherlapati, R. (1994) Organization of the human keratin type II gene cluster at 12q13 *Genomics* 24 (3): 502-8
- Yoshino, M., Xiao, H., Amadou, C., Jones, E. P. and Lindahl, K. F. (1998a) BAC clones and STS markers near the distal breakpoint of the fourth t-inversion, In(17)4d, in the H2-M region on mouse chromosome 17 *Mamm Genome* 9 (3): 186-92
- Yoshino, M., Xiao, H., Jones, E. P. and Fischer Lindahl, K. (1998b) BAC/YAC contigs from the H2-M region of mouse Chr 17 define gene order as Znf173-Tctex5-mog-D17Tu42-M3-M2 *Immunogenetics* 47 (5): 371-80
- Yoshino, M., Xiao, H., Jones, E. P., Kumanovics, A., Amadou, C. and Fischer Lindahl, K. (1997) Genomic evolution of the distal Mhc class I region on mouse Chr 17 *Hereditas* 127 (1-2): 141-8
- Young, J. M. and Trask, B. J. (2002) The sense of smell: genomics of vertebrate odorant receptors *Hum Mol Genet* 11 (10): 1153-60
- Young, J. M., Friedman, C., Williams, E. M., Ross, J. A., Tonnes-Priddy, L. and Trask, B. J. (2002) Different evolutionary processes shaped the mouse and human olfactory receptor gene families *Hum Mol Genet* 11 (5): 535-46
- Younger, R. M., Amadou, C., Bethel, G., Ehlers, A., Lindahl, K. F., Forbes, S., Horton, R., Milne, S., Mungall, A. J., Trowsdale, J., Volz, A., Ziegler, A. and Beck, S. (2001) Characterization of clustered MHC-linked olfactory receptor genes in human and mouse *Genome Res* 4 (11): 519-30

- Zeng, F. Y. and Wess, J. (1999) Identification and molecular characterization of m3 muscarinic receptor dimers *J Biol Chem* 274 (27): 19487-97
- Zerial, M., Salinas, J., Filipski, J. and Bernardi, G. (1986) Gene distribution and nucleotide sequence organization in the human genome *Eur J Biochem* 160 (3): 479-85
- Zhang, M. Q. (1997) Identification of protein coding regions in the human genome by quadratic discriminant analysis *Proc Natl Acad Sci U S A* 94 (2): 565-8
- Zhang, Q. H., Ye, M., Wu, X. Y., Ren, S. X., Zhao, M., Zhao, C. J., Fu, G., Shen, Y., Fan, H. Y., Lu, G., Zhong, M., Xu, X. R., Han, Z. G., Zhang, J. W., Tao, J., Huang, Q. H., Zhou, J., Hu, G. X., Gu, J., Chen, S. J. and Chen, Z. (2000) Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34+ hematopoietic stem/progenitor cells *Genome Res* 10 (10): 1546-60
- Zhang, X. and Firestein, S. (2002) The olfactory receptor gene superfamily of the mouse *Nat Neurosci* 5 (2): 124-33
- Zhao, H. and Firestein, S. (1999) Vertebrate odorant receptors *Cell Mol Life Sci* 56 (7-8): 647-59
- Zhao, H., Ivic, L., Otaki, J. M., Hashimoto, M., Mikoshiba, K. and Firestein, S. (1998) Functional expression of a mammalian odorant receptor *Science* 279 (5348): 237-42
- Zhou, L. J., Schwarting, R., Smith, H. M. and Tedder, T. F. (1992) A novel cell-surface molecule expressed by human interdigitating reticulum cells, Langerhans cells, and activated lymphocytes is a new member of the Ig superfamily *J Immunol* 149 (2): 735-42
- Zhou, Q., Hinkle, G., Sogin, M. L. and Dionne, V. E. (1997) Phylogenetic analysis of olfactory receptor genes from mudpuppy (*Necturus maculosus*) *Biol Bull* 193 (2): 248-50
- Ziegler, A., Dohr, G. and Uchanska-Ziegler, B. (2002) Possible roles for products of polymorphic MHC and linked olfactory receptor genes during selection processes in reproduction *Am J Reprod Immunol* 48 (1): 34-42
- Ziegler, A., Beck, S., Ehlers, A., Younger, R. and Volz, A. (2002) *Testicular transcripts of HLA-linked olfactory receptor genes exhibit unorthodox features* In Hansen, J.A. and Dupont, B. (Ed) *HLA 2002 Immunobiology of the Human MHC*
- Ziegler, A., Ehlers, A., Forbes, S., Trowsdale, J., Volz, A., Younger, R. and Beck, S. (2000) Polymorphisms in olfactory receptor genes: a cautionary note *Hum Immunol* 61 (12): 1281-4
- Ziegler, A., Muller, C., Heinig, J., Radka, S. F., Kompf, J. and Fonatsch, C. (1985) Monosomy 6 in a human lymphoma line induced by selection with a monoclonal antibody *Immunobiology* 169 (5): 455-60
- Zozulya, S., Echeverri, F. and Nguyen, T. (2001) The human olfactory receptor repertoire *Genome Biol* 2 (6): RESEARCH0018